



Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Free Radical Biology and Medicine

journal homepage: www.elsevier.com/locate/freeradbiomed



Corrigendum

Corrigendum to “Low molar excess of 4-oxo-2-nonenal and 4-hydroxy-2-nonenal promote oligomerization of alpha-synuclein through different pathways” [Free Rad. Biol. Med. (2017) 421–431]



Leire Almandoz-Gil^a, Hedvig Welander^a, Elisabeth Ihse^a, Payam Emami Khoonsari^b,
Sravani Musunuri^b, Christofer Lendel^c, Jessica Sigvardson^d, Mikael Karlsson^e, Martin Ingelsson^a,
Kim Kultima^{b,1}, Joakim Bergström^{a,*,1}

^a Department of Public Health and Caring Sciences, Molecular Geriatrics, Uppsala University, SE-751 85 Uppsala, Sweden

^b Department of Medical Sciences, Uppsala University, SE-751 85 Uppsala, Sweden

^c Department of Chemistry, KTH Royal Institute of Technology, SE-100 44, Stockholm, Sweden

^d BioArctic AB, Warfväges väg 35, SE-112 51 Stockholm, Sweden

^e Department of Engineering Sciences, Uppsala University, SE-751 21 Uppsala, Sweden

The authors regret the omission of the concentration of 140 μM on page 422 and the concentration of 840 μM on page 423 in the section “2.4. Production of unmodified alpha-synuclein oligomers through

lyophilization”

The authors would like to apologise for any inconvenience caused.

DOI of original article: <http://dx.doi.org/10.1016/j.freeradbiomed.2017.07.004>

* Corresponding author.

¹ These authors contributed equally to this work.

E-mail address: Joakim.bergstrom@pubcare.uu.se (J. Bergström).

<https://doi.org/10.1016/j.freeradbiomed.2018.02.007>

Available online 16 February 2018

0891-5849/ © 2018 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).