

Anger frequency and risk of cardiovascular morbidity and mortality

Olga E. Titova ^{1,*}, John A. Baron^{1,2,3}, Karl Michaëlsson ¹,
and Susanna C. Larsson ^{1,4,*}

¹Department of Surgical Sciences, Unit of Medical Epidemiology, Uppsala University, Uppsala, Sweden; ²Department of Medicine, University of North Carolina School of Medicine, Chapel Hill, NC, USA; ³Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC, USA; and ⁴Unit of Cardiovascular and Nutritional Epidemiology, Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

Received 16 May 2022; revised 22 July 2022; accepted 31 July 2022; online publish-ahead-of-print 6 August 2022

Handling Editor: Karolina Szummer

Aims

Anger may increase the risk of cardiovascular diseases (CVDs) but previous findings are inconclusive and large prospective studies are needed. We investigated whether frequency of strong anger is associated with the incidence of specific CVDs and CVD mortality, and if sex, age, and cardiometabolic risk factors modify these associations.

Methods and results

We used data from a population-based cohort of 47 077 Swedish adults (56–94 years of age) who completed questionnaires regarding their experience of anger, lifestyle habits, and health characteristics. Participants were followed for incident cardiovascular outcomes and death up to 9 years through linkage to the Swedish National Patient and Death Registers. Hazard ratios and confidence intervals adjusted for potential confounders were assessed.

In multivariable analyses, frequent episodes of strong anger were associated with an increased risk of heart failure, atrial fibrillation, and CVD mortality [hazard ratios (95% confidence intervals) = 1.19 (1.04–1.37), 1.16 (1.06–1.28), and 1.23 (1.09–1.40), respectively]. The link between anger frequency and heart failure was more pronounced in men and participants with a history of diabetes. No evidence of an independent association of anger frequency with risk of myocardial infarction, aortic valve stenosis, and abdominal aortic aneurysm was found.

Conclusion


Our findings indicate that anger may contribute to the development of specific CVDs and CVD mortality, especially heart failure in men and in those with diabetes.

* Corresponding authors. Emails: olga.titova@surgsci.uu.se (O.E.T.); susanna.larsson@surgsci.uu.se (S.C.L.)

© The Author(s) 2022. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Graphical Abstract



Keywords

Cardiovascular disease • Cardiovascular mortality • Anger • Sex • Cohort • Diabetes

Introduction

As the major cause of severe long-term disability and pre-mature death, cardiovascular diseases (CVDs) impose a large burden on healthcare systems and society. In addition to traditional risk factors for CVDs (e.g. obesity, Type 2 diabetes, hypertension, and smoking), psychological factors, such as anxiety, depression, and stress, have been associated with an increased risk of some CVDs.¹⁻³ Negative emotions, including strong anger, may also influence the risk of CVDs. A meta-analysis of nine case-crossover studies found an increased risk of myocardial infarction (MI), acute coronary syndromes, stroke, and ventricular arrhythmia in the 2 h following outbursts of anger.⁴ While there is evidence that strong anger may trigger acute cardiovascular events, the long-term influence of anger on CVD events has scarcely been investigated and the scarce available data are inconsistent. Several cohort studies have indicated that anger (higher anger expression, frequent episodes of anger, or trait anger) is linked to higher risk of total CVD,⁵ atrial fibrillation (AF),⁶ coronary heart disease (CHD),^{7,8} heart failure (HF),⁹ and CVD mortality,¹⁰ but other cohort studies and meta-analyses showed no such associations^{11,12} or demonstrated an inverse relationship with some CVDs such as MI and stroke.¹³ There appear to be no cohort studies of anger and the incidence of aortic valve stenosis (AVS) or abdominal aortic aneurysm (AAA). Moreover, it remains unclear whether age, sex, and modifiable factors for CVDs (e.g. metabolic disorders) modify these associations.

Here we report the findings from a population-based cohort study of 47 077 middle-aged and elderly men and women in which we investigated whether the self-reported frequency of episodes of strong anger is associated with the later occurrence of specific CVDs, including MI, HF, AF, AVS, AAA, as well as with overall CVD mortality. In addition, we investigated whether age, sex, and a history of cardiometabolic CVD risk factors modify these associations.

Methods

Study population

We used data from the national research infrastructure SIMPLER (Swedish Infrastructure for Medical Population-based Life-course Environmental Research) that include the Cohort of Swedish Men (COSM) and the Swedish Mammography Cohort (SMC). COSM and SMC are population-based longitudinal cohorts which were designed to investigate the link between dietary exposures, lifestyle, and disease outcomes.¹⁴ Details of the study cohorts have been reported elsewhere.¹⁴ Briefly, the SMC was established between 1987 and 1990, when all women living in Västmanland and Uppsala counties and born between 1914 and 1948 received a questionnaire on dietary habits and other characteristics (90 303 women invited, 74% response rate). The COSM was established in 1997 and included men living in Västmanland and Örebro counties, born between 1918 and 1952 (100 303 men invited, 49% response rate).¹⁴ In 1997, the 56 030 participants of SMC who were still alive and resided in the study area received an expanded diet and lifestyle questionnaire similar to one sent to COSM participants (70%

respond rate).¹⁴ Questionnaire sent to the participants in 1997 did not include questions related to experience of anger. Therefore, information on lifestyle and health characteristics for the present study was obtained with structured questionnaires in 2008 and 2009 supplemented with national registry information. In the present analysis, we excluded individuals who died or had a specific CVD endpoint prior to 1 January 2009, as verified through linkage to the Swedish National Patient and Death Registers. In addition, we excluded those who had missing information on anger (see [Supplementary material online, Figure S1](#)). This left 47 077 eligible participants (21 442 women and 25 635 men) with a mean baseline age of 70 (56–94) years. In the statistical analysis of each CVD outcome, we excluded individuals with a diagnosis of the corresponding specific CVD before start of a follow up (e.g. those who received diagnosis of AF before baseline were excluded from the analysis of AF), as determined through linkage to the Swedish National Patient Register and based on the International Classification of Diseases (ICD) codes. The number of prevalent CVD cases excluded in each analysis is shown in [Supplementary material online, Figure S1](#). The study was conducted following the Helsinki declaration, and has been approved by the Swedish Ethical Review Authority. Participants provided written informed consent.

Exposure and cardiometabolic risk factor assessment

In 2008–9, participants completed questionnaires that included information on how often they experienced strong anger. Frequency of feelings of strong anger was assessed by a single question without a specified period of time: ‘How often do you feel strong anger?’ A similar single question has been used in previous studies to evaluate anger frequency.^{10,13} Participants indicated how many times per day or week they generally experience strong anger. ‘Never’ or ‘seldom’ was also possible responses. Individuals were divided into two groups: those who never or seldom felt strong anger and those who indicated generally having feelings of strong anger at least once a week. Educational attainment, employment status, smoking, alcohol consumption, weight, height, physical activity, and history of diabetes, depression, hypertension, and hypercholesterolaemia were assessed with structured questionnaires. A history of diabetes was defined based on the first available record of any type of diabetes mellitus (based on the ICD 8th, 9th, and 10th revision codes) before start of baseline, self-reported diabetes at baseline, glucose-lowering drug prescription (based on the Swedish Prescribed Drug Register), and information from the National Diabetes Register.

Identification of incident cardiovascular disease outcomes, cardiovascular disease-related mortality and follow up

Information regarding specific CVDs and death were obtained by linkage to the Swedish National Patient Register and the Cause of Death Register using the unique personal identification number assigned to all Swedish residents. The Patient Register contains information on all inpatient diagnosis since 1987 and outpatient specialist care since 2001.¹⁵ Incident cases were classified according to the ICD 10th revision codes as follows: acute MI (I21), HF (I50 and I11.0), AF (I48), AVS (I35.0 and I35.2), AAA (I71.3 and I71.4), and CVD mortality (I00–I99, primary cause of death). Participants were followed up from 1 January 2009 to the date of diagnosis of CVD

or CVD mortality, death from any cause, or 31 December 2017, whichever occurred first.

Statistical analysis

Descriptive data are presented as mean (standard deviation) or as percentages. Cox proportional hazards regression models were used to estimate hazard ratios (HRs) and corresponding 95% confidence intervals (CIs) for specific CVDs. The main exposure variable, frequency of the experience of strong anger, was treated as a binary variable (at least once a week vs. never/seldom). A basic model incorporated age (as the time scale) and sex (as a stratification variable). In the first multivariable model, we added social and lifestyle factors as covariates including education (less than high school, high school, or university), employment status (working or not working), cigarette smoking (never, former, or current smokers), alcohol consumption (never drinkers; past or current drinkers of <1 drink/week; <7 drinks/week; 7 to <15 drinks/week; 15–21 drinks/week; or >21 drinks/week), walking/bicycling (never/seldom; <20 min/day; 20–40 min/day; or >40 min/day), and exercise (almost never; <1 h/week; 1 h/week; 2–3 h/week; 4–5 h/week; or ≥5 h/week). In the second multivariable model, we additionally adjusted for body mass index (weight divided by the square of height; <22.5, 22.5–24.9, 25.0–29.9, or ≥30 kg/m²), and history of hypercholesterolaemia (no/yes), depression (no/yes), diabetes (no/yes), and hypertension (no/yes). Potential confounders were selected using directed acyclic graphs¹⁶ based on our knowledge of the relationships among potential confounders, intermediate variables, exposure, and outcome variables, as well as on existing information regarding factors associated with CVD and anger.

Proportional hazard assumptions were assessed by Schoenfeld's test. We examined multiplicative interactions of the experience of strong anger with sex, age, and a history of metabolic risk factors for CVDs. The proportion of missing data on the potential confounders in the analysis was small (≤3%), and a separate category for variables containing missing values was created. With regard to CVD mortality, we performed a sensitivity analysis additionally excluding individuals who were diagnosed with any CVD before the start of a follow up ($n = 7625$). All statistical tests were two sided, and the analyses were performed using Stata version 15.1 (StataCorp, College Station, TX, USA).

Results

In total, 4266 participants (9%) reported experiencing strong anger once a week or more often. Baseline characteristics of study participants according to frequency of anger episodes are shown in [Table 1](#) and [Supplementary material online, Table S1](#). Compared with participants who seldom or never experienced strong anger, those who indicated frequent episodes were more likely to report a history of depression, but there were only modest differences in age, sex, educational level, employment status, smoking status, alcohol intake, body stature, leisure time physical activity, prevalence of diabetes, and hypercholesterolaemia ([Table 1](#)).

The numbers of incident CVD events during up to 9 years of follow up can be found in [Figures 1](#) and [2](#) and see [Supplementary material online, Table S2](#). For example, there were 2640 incident

Table 1 Baseline characteristics of the study participants according to reports of strong anger

Characteristics	Frequency of strong anger		
	All	Never/seldom	At least once a week
Number of participants	47 077	42 811	4266
Age at baseline, years, mean (SD)	70.0 (8.1)	70.2 (8.1)	67.9 (7.4)
Men, %	54.5	54.8	51.3
Education >12 years, %	21.3	20.8	26.1
Full-time or part-time employed, %	26.7	26.1	33.4
Cigarette smoking, %			
Former smokers	36.6	36.1	41.1
Current smokers	8.5	8.4	9.6
Alcohol intake ≥ 15 drinks/week, %	3.4	3.3	4.4
Walking/bicycling >40 min/day, %	33.3	33.5	31.6
Exercise ≥ 2 h/week, %	15.0	15.0	15.6
Body mass index, kg/m ² , %			
25.0–29.9	41.6	41.6	41.0
≥ 30.0	13.3	13.0	15.9
History of depression, %	8.4	7.5	17.3
History of hypertension, %	42.1	42.1	42.3
History of hypercholesterolaemia, %	25.4	25.2	27.0
History of diabetes, %	10.8	10.7	11.8

SD, standard deviation.

HF events identified in this cohort during 377 406 person-years of follow up. In the final multivariable model, frequent episodes of strong anger were associated with a 19% higher risk of HF (HR, 1.19; 95% CI, 1.04–1.37). In addition, a 16% higher risk of AF and 23% higher risk of CVD mortality were found in participants who reported frequent episodes of strong anger (HR, 1.16; 95% CI, 1.06–1.28; and HR, 1.23; 95% CI, 1.09–1.40, respectively; [Figure 1](#), see [Supplementary material online, Table S2](#) and Graphical abstract). We found no link between frequency of self-reported anger and risk of MI, AVS, or AAA ([Figure 1](#), see [Supplementary material online, Table S2](#) and Graphical abstract).

No interaction between frequent experience of strong anger and age was found for any CVD outcome (P for interaction >0.60 , basic model; in this model, age was added as a covariate, not as the time scale). However, a statistically significant interaction with sex was observed in relation to incidence of HF only (P for interaction = 0.04, basic model; in this model, sex was added as a covariate). Subsequent multivariable analyses revealed a significant association of anger frequency and incidence of HF in men (HR, 1.30; 95% CI, 1.10–1.54) but not among women (HR, 1.02; 95% CI, 0.81–1.29; [Figure 2](#)). In addition, there was a significant interaction of anger with diabetes history in relation to HF (P for interaction = 0.048, basic model). The association was more pronounced in participants with a history of diabetes (multivariable HR, 1.39; 95% CI, 1.06–1.84) than in those without a diabetes history ([Figure 2](#)). No significant interaction of strong anger with BMI, history of hypertension, or hypercholesterolaemia in relation to any CVD outcome was observed ($P > 0.10$, basic model).

In a sensitivity analysis, the findings for CVD mortality were similar after additional exclusion of participants with any CVD before

baseline (multivariable HR, 1.28; 95% CI, 1.08–1.53; $n = 39\,452$, number of cases = 1811).

Discussion

In our large cohort study, we found positive associations of the frequency of the experience of strong anger with the incidence of HF, AF, and CVD mortality. There was some evidence that the link between frequent episodes of strong anger at baseline and risk of HF varied by sex and history of diabetes. Thus, our findings demonstrate that men as well as participants with a history of diabetes at baseline who experienced frequently strong anger are at particular risk of HF. No evidence of an association of anger frequency with risk of MI, AVS, and AAA was found.

Our findings of a positive association between anger and incident HF agrees with the results from a cohort of 13 171 middle-aged men and women in the USA, which indicated a 26% increased risk of HF in men with high trait anger compared with low trait anger. No evidence of the association of anger-proneness and risk of HF was observed in women in this study.⁹ In contrast, a multiethnic population-based study of 6782 initially healthy individuals (242 total incident HF cases) did not find an association of anger and other psychosocial factors with risk of incident HF.¹¹ The observed inconsistency of these findings with the others could be related to the smaller sample size and relatively small number of cases in the latter study.¹¹

Prospective cohort studies evaluating the association of anger and risk of AF are limited and inconsistent. Our analysis indicates that frequent episodes of anger may increase the risk of AF. Another cohort study of 3873 men and women (age range 18–77 years) also




Figure 1 Multivariable hazard ratio (95% confidence interval) of cardiovascular diseases and cardiovascular disease mortality according to reports of frequency of strong anger. The Cox proportional hazards regression model was adjusted for age (underlying time scale), sex (as a stratification variable), education, employment status, cigarette smoking, alcohol intake, walking/bicycling, exercise, body mass index, and history of hypercholesterolaemia, hypertension, diabetes, and depression. CI, confidence interval; HR, hazard ratio.

demonstrated a relationship of measures of anger and hostility and increased risk of AF in men.⁶ However, no association of anger and AF risk was observed in a multiethnic study of 6644 participants.¹⁷ Several studies investigated the link of different anger-related measures, mainly hostility, and all-cause mortality risk, indicating a positive association^{6,18,19} or null findings.^{10,20} In addition, a recent cohort study of 17 352 men reported that frequent episodes of anger are associated with a 17% increased risk of CVD death over the 20-year follow-up period.¹⁰ Our study, which included women as well as men, confirms these findings.

Our results on anger and MI support findings of other cohort studies that did not find a significant association of measures of anger and risk of MI^{7,21} but our estimates cannot rule out modestly strong associations. Other cohort studies in men have reported that high levels of hostility or anger are associated with a considerably elevated risk of acute MI.^{22,23} Interestingly, a study of 23 522 male health professionals (aged 50–85 years) demonstrated a reduced risk of non-fatal MI during a 2-year follow up in participants with moderate levels of anger-out expression (outwardly expressive behaviour in response to feeling angry), whereas no association of high levels of anger expression and incidence of MI was observed.¹³ The conceptual difference between anger, cynical hostility, and aggression used as exposure variables in different studies could explain discrepancies

between findings of the studies' in relation to MI. For example, hostility is considered as a cognitive characteristic, a negative aggressive attitude towards other people consisting of denigration, enmity, and ill will.²⁴ Anger, on the other hand, is a negative emotion of different intensity and as a personality trait, it refers to the tendency to experience frequent episodes of this emotion.²⁴ Aggression is described as a verbal and physical behaviour involving attacking and destructive actions.²⁴ We are unaware of any study of anger measures in relation to risk of AVS and AAA.

In the available literature, anger-related measures were evaluated with various instruments (e.g. the Spielberger Trait Anger Scale or Framingham scales^{6,10,17} or with a single question of anger frequency^{10,13}). The discrepancies in the results of different studies that investigated the relationship with the risk of specific CVDs or CVD mortality may be at least partially explained by differences in methodological tools or small number of cases. In addition, a number of studies had a case-crossover design^{25,26} or were performed on populations with pre-existing CVDs,^{27–29} and so were open to reporting bias or reverse causation. Another consideration is that the associations of anger with CVDs may vary by ethnicity.⁹

The sex-specific association of anger with HF observed in our study may be related to different ways of anger expression and coping style among women and men,³⁰ biological (e.g. levels of sex




Figure 2 Multivariable hazard ratio (95% confidence interval) of heart failure according to anger frequency stratified by sex and history of diabetes. The Cox proportional hazards regression model was adjusted for age (underlying time scale), sex (as a stratification variable, analysis by diabetes history), education, employment status, cigarette smoking, alcohol intake, walking/bicycling, exercise, body mass index, and history of hypercholesterolaemia, hypertension, diabetes (analysis by sex), and depression. CI, confidence interval; HR, hazard ratio; DM, diabetes mellitus.

hormones)³¹ and lifestyle differences, and comorbidities. Several previous studies and meta-analyses have also reported sex-specific relationship of anger measures with CVD events and cardiometabolic risk factors, and demonstrated a stronger association in men than in women.^{32,33}

There is still uncertainty in the findings of anger and specific CVDs risk, and factors other than sex and age may influence these associations. Besides sex differences, we have found that the association of anger frequency and risk of HF is stronger in participants with a history of diabetes, suggesting that diabetes may modify this association. Evidence suggests that the prevalence of anxiety and depression is higher in individuals with diabetes mellitus than in healthy controls,^{34,35} and mood changes associated with fluctuations of blood glucose levels in participants with insulin-dependent diabetes mellitus have been previously reported.³⁶ Mood changes such as anger and hostility often accompany depression including patients with diabetes.³⁷ However, these relationships are likely bidirectional, and psychosocial factors may increase the risk of diabetes.³⁸ In addition, diabetes is associated with a higher risk of CVDs and CVD mortality.³⁹ For example, epidemiological and clinical studies report a high prevalence of HF in patients with prediabetes or diabetes.⁴⁰

Although the exact biological mechanism by which anger may increase the risk of specific CVDs remains unknown, several potential pathways may explain the role of anger in the development of CVDs such as exposure to stress and physiological response to stress (e.g.

increased heart rate and blood pressure), inflammation, unfavourable lifestyle, and behaviour choices. Irritability and anger can appear as a consequence of acute or chronic stress. Psychosocial stress activates neuroendocrine stress–response systems, the sympathetic–adrenal–medullary system, and hypothalamus–pituitary–adrenal–cortical axis system, which results in release of adrenaline, noradrenaline, and cortisol as well as increased heart rate and blood pressure.⁴¹ For example, anger and hostility have been linked to elevated cortisol levels,⁴² which have been associated with cardiometabolic risk factors such as metabolic syndrome components⁴³ as well as with AF,⁴⁴ CVD morbidity⁴⁵ and mortality.⁴⁶

Strength and limitations

There are several strengths of our study, including a large sample size providing a large number of cases of a broad range of CVD outcomes, complete case identification with no loss to follow up, adjustment for potential confounders, and inclusion of both women and men. We cannot exclude that pharmacological agents used to treat metabolic conditions and not considered in the present analysis, could affect the expression of anger and risk of CVDs. In addition, as anger was measured with a single question, we cannot distinguish between ‘anger-in’ (anger suppression and direction it towards themselves) and ‘anger-out’ (anger expression directed towards

others) behaviours.⁴⁷ Furthermore, although we excluded individuals with prevalent specific CVDs based on the ICD codes, participants did not undergo clinical diagnostic investigations at baseline, and therefore we cannot rule out any pre-existing but undiagnosed forms of CVD. We conducted several interaction analyses for several outcomes, and we cannot rule out that the observed interactions are chance findings. Since our study included women and men of mainly northern European origin, our findings might be not generalizable to other ethnic groups. Finally, due to the observational nature of this study, we cannot exclude residual and unmeasured confounding.

Conclusion

In the present study, frequent experience of strong anger was associated with increased risk of HF, AF, and CVD mortality in the initially healthy middle-aged and older individuals. A stronger association of anger frequency with risk of HF was found in men and participants with a history of diabetes. Our findings extend the recent biological and epidemiological findings that anger may contribute to the development of specific CVDs and CVD mortality.

Lead author biography



Susanna Larsson is currently a senior lecturer at the Department of Surgical Sciences, Uppsala University, Sweden and affiliated with the Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden. She has a PhD degree (2006) from Karolinska Institutet, and she became an associate professor of epidemiology in 2011. Her research is focused on identifying the causes of cardiovascular diseases and other chronic diseases.

Author contributions

O.E.T. and S.C.L. contributed to the conception and design of the study; O.E.T. acquired the data, performed the statistical analysis, and drafted the manuscript. All authors contributed to the interpretation of the results and critical revision of the manuscript for important intellectual content and approved the final version of the manuscript.

Data availability

The data that support findings of this prospective cohort study are available upon application to the Swedish Infrastructure for Medical Population-based Life-course Environmental Research (SIMPLER; <https://www.simpler4health.se>).

Supplementary material

Supplementary material is available at *European Heart Journal Open* online.

Acknowledgements

The authors thank the national research infrastructure SIMPLER for provisioning of facilities and experimental support. The computations were performed on resources provided by the Swedish National Infrastructure for Computing (www.snic.se) support for sensitive data SNIC-SENS through the Uppsala Multidisciplinary Center for Advanced Computational Science (UPPMAX) under Project SIMP2019004.

Funding

Work of the authors is supported by grants from the Börjeson, Emil and Ragna Foundation (to O.E.T.), PO Zetterlings stiftelse (to O.E.T.); the Swedish Research Council for Health, Working Life and Welfare (Forte; grant number 2018-00123) (to S.C.L.), the Swedish Research Council (Vetenskapsrådet; grant number 2016-01042 and 2019-00977) (to S.C.L.), and the Swedish Heart-Lung Foundation (Hjärt-Lungfonden; grant number 20210351) (to S.C.L.). The study was also supported by additional grants from the Swedish Research Council (<https://www.vr.se>; grant numbers 2015-03257, 2017-00644, and 2017-06100 to K.M.). SIMPLER receives funding through the Swedish Research Council under the grant number 2017-00644 (to Uppsala University and K.M.). SNIC is financially supported by the Swedish Research Council. The funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Conflict of interest: None declared.

References

- Emdin CA, Odotayo A, Wong CX, Tran J, Hsiao AJ, Hunn BH. Meta-analysis of anxiety as a risk factor for cardiovascular disease. *Am J Cardiol* 2016;**118**:511–519.
- Nicholson A, Kuper H, Hemingway H. Depression as an aetiologic and prognostic factor in coronary heart disease: a meta-analysis of 6362 events among 146 538 participants in 54 observational studies. *Eur Heart J* 2006;**27**:2763–2774.
- Öhlin B, Nilsson PM, Nilsson JA, Berglund G. Chronic psychosocial stress predicts long-term cardiovascular morbidity and mortality in middle-aged men. *Eur Heart J* 2004;**25**:867–873.
- Mostofsky E, Penner EA, Mittleman MA. Outbursts of anger as a trigger of acute cardiovascular events: a systematic review and meta-analysis. *Eur Heart J* 2014;**35**:1404–1410.
- Tezuka K, Kubota Y, Ohira T, Shimizu Y, Yamagishi K, Umesawa M, Sankai T, Imano H, Okada T, Kiyama M, Iso H, Investigators C. Anger expression and the risk of cardiovascular disease among urban and rural Japanese residents: the circulatory risk in communities study. *Psychosom Med* 2020;**82**:215–223.
- Eaker ED, Sullivan LM, Kelly-Hayes M, D'Agostino RB Sr, Benjamin EJ. Anger and hostility predict the development of atrial fibrillation in men in the Framingham offspring study. *Circulation* 2004;**109**:1267–1271.
- Kawachi I, Sparrow D, Spiro A 3rd, Vokonas P, Weiss ST. A prospective study of anger and coronary heart disease. The normative aging study. *Circulation* 1996;**94**:2090–2095.
- Williams JE, Paton CC, Siegler IC, Eigenbrodt ML, Nieto FJ, Tyroler HA. Anger proneness predicts coronary heart disease risk: prospective analysis from the atherosclerosis risk in communities (ARIC) study. *Circulation* 2000;**101**:2034–2039.
- Kucharska-Newton AM, Williams JE, Chang PP, Stearns SC, Sueta CA, Blecker SB, Mosley TH. Anger proneness, gender, and the risk of heart failure. *J Card Fail* 2014;**20**:1020–1026.
- Trudel-Fitzgerald C, Reduron LR, Kawachi I, Kubzansky LD. Specificity in associations of anger frequency and expression with different causes of mortality over 20 years. *Psychosom Med* 2021;**83**:402–409.
- Ogilvie RP, Everson-Rose SA, Longstreth WT Jr, Rodriguez CJ, Diez-Roux AV, Lutsey PL. Psychosocial factors and risk of incident heart failure: the multi-ethnic study of atherosclerosis. *Circ Heart Fail* 2016;**9**:e002243.
- Fu Y, He W, Ma J, Wei B. Relationship between psychological factors and atrial fibrillation: a meta-analysis and systematic review. *Medicine (Baltimore)* 2020;**99**:e19615.

13. Eng PM, Fitzmaurice G, Kubzansky LD, Rimm EB, Kawachi I. Anger expression and risk of stroke and coronary heart disease among male health professionals. *Psychosom Med* 2003;**65**:100–110.
14. Harris H, Håkansson N, Olofsson C, Stackelberg O, Julin B, Åkesson A, Wolk A. The Swedish mammography cohort and the cohort of Swedish men: study design and characteristics of 2 population-based longitudinal cohorts. *OA Epidemiol* 2013;**1**:16.
15. Ludvigsson JF, Andersson E, Ekblom A, Feychting M, Kim J-L, Reuterwall C, Heurgren M, Olausson PO. External review and validation of the Swedish national inpatient register. *BMC Public Health* 2011;**11**:450.
16. Textor J, Hardt J, Knüppel S. DAGitty: a graphical tool for analyzing causal diagrams. *Epidemiology* 2011;**22**:745.
17. Garg PK, O'Neal WT, Diez-Roux AV, Alonso A, Soliman EZ, Heckbert S. Negative affect and risk of atrial fibrillation: MESA. *J Am Heart Assoc* 2019;**8**:e010603.
18. Appleton KM, Woodside JV, Arweiler D, Haas B, Amouyel P, Montaye M, Ferrieres J, Ruidavets JB, Yarnell JW, Kee F, Evans A, Bingham A, Ducimetiere P, Patterson CC. A role for behavior in the relationships between depression and hostility and cardiovascular disease incidence, mortality, and all-cause mortality: the prime study. *Ann Behav Med* 2016;**50**:582–591.
19. Boscarino JA, Figley CR. The impact of repression, hostility, and post-traumatic stress disorder on all-cause mortality: a prospective 16-year follow-up study. *J Nerv Ment Dis* 2009;**197**:461–466.
20. Hearn MD, Murray DM, Luepker RV. Hostility, coronary heart disease, and total mortality: a 33-year follow-up study of university students. *J Behav Med* 1989;**12**:105–121.
21. Stürmer T, Hasselbach P, Amelang M. Personality, lifestyle, and risk of cardiovascular disease and cancer: follow-up of population based cohort. *BMJ* 2006;**332**:1359.
22. Everson SA, Kauhanen J, Kaplan GA, Goldberg DE, Julkunen J, Tuomilehto J, Salonen JT. Hostility and increased risk of mortality and acute myocardial infarction: the mediating role of behavioral risk factors. *Am J Epidemiol* 1997;**146**:142–152.
23. Chang PP, Ford DE, Meoni LA, Wang NY, Klag MJ. Anger in young men and subsequent premature cardiovascular disease: the precursors study. *Arch Intern Med* 2002;**162**:901–906.
24. Smith TW, Glazer K, Ruiz JM, Gallo LC. Hostility, anger, aggressiveness, and coronary heart disease: an interpersonal perspective on personality, emotion, and health. *J Pers* 2004;**72**:1217–1270.
25. Moller J, Hallqvist J, Diderichsen F, Theorell T, Reuterwall C, Ahlbom A. Do episodes of anger trigger myocardial infarction? A case-crossover analysis in the Stockholm Heart Epidemiology Program (SHEEP). *Psychosom Med* 1999;**61**:842–849.
26. Mostofsky E, Maclure M, Tofler GH, Muller JE, Mittleman MA. Relation of outbursts of anger and risk of acute myocardial infarction. *Am J Cardiol* 2013;**112**:343–348.
27. Denollet J, Gidron Y, Vrints CJ, Conraads VM. Anger, suppressed anger, and risk of adverse events in patients with coronary artery disease. *Am J Cardiol* 2010;**105**:1555–1560.
28. Lampert R, Burg MM, Jamner LD, Dziura J, Brandt C, Li F, Donovan T, Soufer R. Effect of beta-blockers on triggering of symptomatic atrial fibrillation by anger or stress. *Heart Rhythm* 2019;**16**:1167–1173.
29. Frasure-Smith N, Lesperance F. Depression and other psychological risks following myocardial infarction. *Arch Gen Psychiatry* 2003;**60**:627–636.
30. Davidson K, Hall P. What does potential for hostility measure? Gender differences in the expression of hostility. *J Behav Med* 1995;**18**:233–247.
31. dos Santos RL, da Silva FB, Ribeiro RF Jr, Stefanon I. Sex hormones in the cardiovascular system. *Horm Mol Biol Clin Investig* 2014;**18**:89–103.
32. Doster JA, Purdum MB, Martin LA, Govens AJ, Moorefield R. Gender differences, anger expression, and cardiovascular risk. *J Nerv Ment Dis* 2009;**197**:552–554.
33. Chida Y, Steptoe A. The association of anger and hostility with future coronary heart disease: a meta-analytic review of prospective evidence. *J Am Coll Cardiol* 2009;**53**:936–946.
34. Roy T, Lloyd CE. Epidemiology of depression and diabetes: a systematic review. *J Affect Disord* 2012;**142**:S8–S21.
35. Khuwaja AK, Lalani S, Dhanani R, Azam IS, Rafique G, White F. Anxiety and depression among outpatients with type 2 diabetes: a multi-centre study of prevalence and associated factors. *Diabetol Metab Syndr* 2010;**2**:72.
36. Gonder-Frederick LA, Cox DJ, Bobbitt SA, Pennebaker JW. Mood changes associated with blood glucose fluctuations in insulin-dependent diabetes mellitus. *Health Psychol* 1989;**8**:45–59.
37. Penckofer S, Quinn L, Byrn M, Ferrans C, Miller M, Strange P. Does glycemic variability impact mood and quality of life? *Diabetes Technol Ther* 2012;**14**:303–310.
38. Mezuk B, Eaton WW, Albrecht S, Golden SH. Depression and type 2 diabetes over the lifespan: a meta-analysis. *Diabetes Care* 2008;**31**:2383–2390.
39. Hackett RA, Steptoe A. Psychosocial factors in diabetes and cardiovascular risk. *Curr Cardiol Rep* 2016;**18**:95.
40. Lehrke M, Marx N. Diabetes Mellitus and heart failure. *Am J Cardiol* 2017;**120**:S37–S47.
41. Ulrich-Lai YM, Herman JP. Neural regulation of endocrine and autonomic stress responses. *Nat Rev Neurosci* 2009;**10**:397–409.
42. Moons WG, Eisenberger NI, Taylor SE. Anger and fear responses to stress have different biological profiles. *Brain Behav Immun* 2010;**24**:215–219.
43. Fraser R, Ingram MC, Anderson NH, Morrison C, Davies E, Connell JM. Cortisol effects on body mass, blood pressure, and cholesterol in the general population. *Hypertension* 1999;**33**:1364–1368.
44. Larsson SC, Lee W-H, Burgess S, Allara E. Plasma cortisol and risk of atrial fibrillation: a Mendelian randomization study. *J Clin Endocrinol Metab* 2021;**106**:e2521–e2526.
45. Crawford AA, Soderberg S, Kirschbaum C, Murphy L, Eliasson M, Ebrahim S, Davey Smith G, Olsson T, Sattar N, Lawlor DA, Timpson NJ, Reynolds RM, Walker BR. Morning plasma cortisol as a cardiovascular risk factor: findings from prospective cohort and Mendelian randomization studies. *Eur J Endocrinol* 2019;**181**:429–438.
46. Vogelzangs N, Beekman ATF, Milaneschi Y, Bandinelli S, Ferrucci L, Penninx BWJH. Urinary cortisol and six-year risk of all-cause and cardiovascular mortality. *J Clin Endocrinol Metab* 2010;**95**:4959–4964.
47. Spielberger CD, Reheiser EC, Sydeman SJ. Measuring the experience, expression, and control of anger. *Issues Compr Pediatr Nurs* 1995;**18**:207–232.