



<http://www.diva-portal.org>

Preprint

This is the submitted version of a paper published in *Upsala Journal of Medical Sciences*.

Citation for the original published paper (version of record):

Wang, X., Welsh, N. (2014)

Bcl-2 maintains the mitochondrial membrane potential, but fails to affect production of reactive oxygen species and endoplasmic reticulum stress, in sodium palmitate-induced beta-cell death.

*Upsala Journal of Medical Sciences*, 119(4): 306-315

<http://dx.doi.org/10.3109/03009734.2014.962714>

Access to the published version may require subscription.

N.B. When citing this work, cite the original published paper.

Permanent link to this version:

<http://urn.kb.se/resolve?urn=urn:nbn:se:uu:diva-240449>

**Bcl-2 maintains the mitochondrial membrane potential, but fails to affect production of reactive oxygen species and endoplasmic reticulum stress, in sodium palmitate-induced  $\beta$ -cell death**

**Running title: Bcl-2 maintains the  $\Delta\psi_m$  in palmitate-induced  $\beta$ -cell death**

**Xuan Wang and Nils Welsh**

**Dept. of Medical Cell Biology, Uppsala University, Uppsala, Sweden**

Address correspondence to: Xuan Wang, Dept. of Medical Cell Biology, Uppsala University, Biomedicum, P.O. Box 571, SE-751 23, Uppsala, Sweden. Telephone +46-18 471-42 12, Telefax: +46-18 471 40 59

Email: [xuan.wang@mcb.uu.se](mailto:xuan.wang@mcb.uu.se)

## **Abstract**

**Background** Sodium palmitate causes apoptosis of  $\beta$ -cells and the anti-apoptotic protein Bcl-2 has been shown to counteract this event. However, the exact mechanisms that underlie palmitate-induced pancreatic  $\beta$ -cell apoptosis and through which pathway Bcl-2 executes the protective effect are still unclear.

**Methods** A stable Bcl-2 overexpressing RINm5F cell clone (BMG) and its negative control (B45) were exposed to palmitate for up to 8h and cell viability, mitochondrial membrane potential ( $\Delta\psi_m$ ), reactive oxygen species (ROS) generation, ER stress and NF- $\kappa$ B activation were studied in time course experiments.

**Results** Palmitate exposure for 8h resulted in increased cell death rates, and this event was partially counteracted by Bcl-2. Bcl-2 overexpression promoted in parallel also a delayed induction of GADD153/CHOP and a weaker phosphorylation of BimEL in palmitate exposed cells. At earlier time points (2-4h) palmitate exposure resulted in increased generation of ROS, a decrease in mitochondrial membrane potential ( $\Delta\psi_m$ ) and a modest increase in the phosphorylation of eIF-2 $\alpha$  and IRE1 $\alpha$ . BMG cells produced similar amounts of ROS and displayed the same eIF-2 $\alpha$  and IRE1 $\alpha$  phosphorylation rates as B45 cells. However, the palmitate-induced dissipation of  $\Delta\psi_m$  was partially counteracted by Bcl-2. In addition, basal NF- $\kappa$ B activity was increased in BMG cells.

**Conclusions** Our results indicate that Bcl-2 counteracts palmitate-induced  $\beta$ -cell death by maintaining mitochondrial membrane integrity and augmenting NF- $\kappa$ B activity, but not by affecting ROS production and ER stress.

**Keywords** Bcl-2, mitochondrial membrane integrity, NF- $\kappa$ B, palmitate, pancreatic beta cell death, reactive oxygen species

## Introduction

Type 2 diabetes is often associated with obesity and increased plasma free fatty acid (FFA) levels. When pancreatic insulin producing cells are exposed to FFA for prolonged time periods,  $\beta$ -cell apoptosis can be observed [1]. The exact mechanisms that underlie palmitate-induced pancreatic  $\beta$ -cell apoptosis are still unclear. It may be that palmitate induces several different events, including endoplasmic reticulum (ER) stress [2], mitochondrial perturbations [3], generation of reactive oxygen species [3], ceramide and NO [4], altered  $\text{Ca}^{2+}$  homeostasis [5] and others that together or individually promote  $\beta$ -cell death.

The anti-apoptotic protein Bcl-2 has been found to be localized at the nucleus, mitochondria and ER and may play an important role in the regulation of cell death induced by palmitate [6]. In non- $\beta$ -cells, palmitate-induced apoptosis was associated with dissipation of the mitochondrial membrane potential ( $\Delta\psi_m$ ), two events that were counteracted by Bcl-2 overexpression [6]. FFAs may, at least in part, decrease  $\Delta\psi_m$  by acting as uncouplers of mitochondrial respiration. The uncoupling effect of FFAs is associated with not only a decreased  $\Delta\psi_m$ , but also increased respiration, enhanced ROS generation, lower ATP levels and mitochondrial swelling [3].

Besides acting at the mitochondrial level, Bcl-2 may also affect ER stress-induced apoptosis since ER-targeted Bcl-2 expression protected against tunicamycin-induced ER stress in renal tubular cells [7], possibly via modulation of the permeability of the ER membrane to  $\text{Ca}^{2+}$  [8]. The ER of the pancreatic  $\beta$ -cell is highly active due to the engagement in synthesis and release of insulin. When ER homeostasis is lost,  $\beta$ -cells will initiate the unfolded protein response (UPR) to balance the protein synthesis demand against the ER protein-folding capacity. PKR-like ER kinase (PERK), inositol requiring ER-to-nucleus signal kinase 1 (IRE1) and activating transcription factor 6 (ATF6) are three ER membrane-located ER stress transducers activated by palmitate [2]. Active PERK phosphorylates the eukaryotic translation initiation factor (eIF-2 $\alpha$ ), thereby generally attenuating protein translation and decreasing the functional demand on the ER. Meanwhile, phosphorylation of eIF-2 $\alpha$  induces activating transcription factor 4 (ATF4) expression. ATF4 is a transcription factor that regulates expression of C/EBP homologous protein (CHOP), also named growth arrest and DNA damage inducible protein 153 (GADD153). Active IRE1 splices X-

box binding protein 1 (XBP1) mRNA. Spliced XBP1 (XBP1s) activates many UPR genes required for protein folding, maturation and degradation. Active ATF6 augments the expression of ER chaperones, such as BiP/GRP78 and GRP94, to increase the protein folding capacity. If the UPR fails to alleviate the ER stress, cells undergo apoptosis, characterized by induction of CHOP and c-Jun N-terminal kinase (JNK), activation of caspase 12, downregulation of Bcl-2, activation of Bax/Bak, Bim, NOXA, PUMA and p53 [9].

In view of the general uncertainty as to how FFAs kill  $\beta$ -cells, and that Bcl-2 intervenes in specific steps in FFA-induced cell death, we have delineated the early time course of palmitate-induced mitochondrial and ER stress events, and how Bcl-2 interferes in these pathways, with the aim to better understand primary palmitate-induced signaling events leading to  $\beta$ -cell death.

## **Materials and methods**

### **Materials**

The chemicals were obtained from the following sources: RPMI 1640, fetal bovine serum, L-glutamine, benzylpenicillin, streptomycin, Hanks' balanced salt solution, trypsin, carbonyl cyanide p-(trifluoromethoxy)phenyl-hydrazone (FCCP), bisbenzimidazole (Hoechst 33342), propidium iodide (PI), 5, 5', 6, 6'-tetrachloro-1, 1', 3, 3'-tetraethylbenzimidazolcarbocyanine iodide (JC-1), sodium palmitate, dichlorofluorescein-diacetate (DCFH-DA), alkaline phosphatase (from bovine intestinal mucosa) and albumin bovine serum (fatty acid and endotoxin free) from Sigma-Aldrich; Bcl-2 and GADD153 (F-168) antibodies from Santa Cruz Biotechnology; phospho-IRE1 $\alpha$  from Affinity BioReagents, phospho-eIF2 $\alpha$ , cleaved caspase 3 and Bim from Cell Signaling and a horseradish-peroxidase-labeled donkey anti-rabbit or anti-mouse antibody from GE Healthcare. All other chemicals of analytical grade were obtained from E. Merck.

### **Cell culture**

RINm5F cells overexpressing Bcl-2 were generated previously [10] by stable liposome-mediated transfection with an episomally maintained multicopy bovine

papilloma virus (BPV)-derived BMG neo vector [11] containing the mouse bcl-2 cDNA [12] inserted downstream of a metallothionein promoter. Stable clones with increased Bcl-2 expression (BMG cells) were isolated and RINm5F cells transfected with an empty BPV-derived neo-containing vector were used as control (B45 cells). The cell clones were cultured in RPMI 1640 supplemented with 10% fetal bovine serum, 2 mM L-glutamine and benzylpenicillin (100 U/ml), streptomycin (0.1 mg/ml) at 37° C in humidified air with 5% CO<sub>2</sub>.

### **FFA preparation and cell treatment**

Palmitate was dissolved in 50% ethanol, heated to 45° C and used at a 1:100 dilution. During FFA exposure, cell were cultured in the RPMI 1640 medium supplemented with 0.5 mM palmitate, 0.5% BSA or 1% BSA (FFA:BSA: molar ratio of 6.6:1 and 3.3:1, respectively). The control condition contained the same amount of vehicle. All cultures were supplemented with 1% FBS.

### **Determination of cell viability**

After exposure to 0.5 mM palmitate (0.5% BSA or 1% BSA), or 1 µg/ml carbonyl cyanide p-(trifluoromethoxy)phenyl-hydrazone (FCCP) (solubilized in 50% ethanol and used in a 1:100 dilution) for 8 h, BMG and B45 cells were vital stained with 20 µg/ml bisbenzimidazole and 10 µg/ml propidium iodide for 15 min at 37° C. Free-floating cells and cell attached to the culture dish were then washed separately with PBS, pooled and examined by inverted fluorescence microscopy (excitation at 365 nm for HO and at 546 nm for PI). In each experimental group 3000-5000 cells were examined and red fluorescence (propidium iodide) and blue fluorescence (bisbenzimidazole) was quantified using Adobe Photoshop software and the ratio red over blue was calculated as a relative measure of cell viability.

### **Determination of the mitochondrial membrane potential ( $\Delta\psi_m$ ) by JC-1 fluorescence**

The mitochondrial membrane potential was semiquantitatively determined using the fluorescent probe JC-1. BMG and B45 cells were incubated for 1, 2, 3, 4, 5, 6, 7, 8 h at 37° C with 0.5 mM palmitate (0.5% BSA). FCCP (1 µg/ml) was also used in the incubation as a positive control of mitochondrial uncoupling. During the last 30 min of incubation, each group of the cells was incubated in the same medium as described

before with the additional supplementation of 4  $\mu$ M JC-1 (solubilized in DMSO, diluted 1:100). After careful washings of the cells, the fluorescence was determined using the Kodak Image Station 4000 MM with excitation wavelength at 465 nm and emission wavelength at 530 nm (green, JC-1 monomers at depolarized membrane potential) and 585 nm (red, J-aggregates at hyperpolarized membrane potential), respectively. The fluorescence-ratio of red over green was used to monitor changes in mitochondrial membrane potential [13].

### **Determination of ROS by DCFH-DA fluorescence**

The production of ROS was estimated using the oxidation-sensitive fluorescent probe DCFH-DA. BMG and B45 cells were cultured for 1, 2, 3 h at 37° C with 0.5 mM palmitate (0.5% BSA). At the end of incubation, 10  $\mu$ M DCFH-DA was applied to each group of cells for 20 min, followed by flow cytometry analysis with BD FACSCalibur (BD Biosciences, San Jose, CA). Fluorescence emission from DCF (green) was detected at a wavelength of 530 nm.

### **Immunoblot analysis**

After incubation with palmitate for 1-8 h, BMG or B45 cells were washed in cold PBS and lysed on ice in SDS-sample buffer (2% SDS, 0.15 M Tris, pH 6.8, 10% glycerol, 5%  $\beta$ -mercaptoethanol, bromphenol blue and 2 mM phenylmethylsulfonyl fluoride). The samples were boiled for 5 minutes and separated on 10% or 12% SDS-PAGE gels. Proteins were transferred to Hybond-P membranes (GE Healthcare). The membranes were pre-blocked in 2.5% BSA for 1h and then incubated with Bcl-2 (1:200), cleaved caspase 3 (1:1000), GADD153 (1:200), phospho-IRE1 $\alpha$  (1:1000), phospho-eIF2 $\alpha$  (1:1000), methyl-Lys221-p65, Bim (1:1000) antibodies over-night at 4° C. After secondary antibody incubation, the bound antibodies were visualized with the Kodak Image Station 4000 MM using ECL plus (GE Healthcare). The band intensities were calculated using Kodak Molecular Imaging Software 4.5.1 SE. Total protein loading and transfer onto the membranes was visualized by amidoblack staining.

### **Statistical analysis**

Data are presented as means  $\pm$  S.E.M. Statistical significance between two experimental conditions was analyzed using Student's paired t-test. Statistical significance: \*P<0.05, #P<0.01.

## **Results**

### **Overexpression of Bcl-2 in RINm5F cells**

To confirm the possibility that overexpression of Bcl-2 might increase resistance to palmitate-induced  $\beta$ -cells death and to investigate through which mechanism Bcl-2 overexpression might execute its protective effect, a bcl-2 transfected insulin-producing rat pancreatic RINm5F cell line BMG was used in subsequent experiments [10]. BMG cells came from the stable clones of RINm5F cells overexpressing Bcl-2 protein 3-4 fold, as assessed by Western blot analysis (Fig. 1A). B45 cells, which were transfected with an empty BPV-derived neo-containing vector and expressed low levels of Bcl-2, were used as control.

### **Palmitate-induced cell death was partially counteracted by Bcl-2 overexpression.**

To investigate whether Bcl-2 protects against saturated FFA-induced cell death, B45 and BMG cells were incubated with 0.5 mM palmitate complexed with 0.5% BSA or 1% BSA (FFA:BSA: molar ratio of 6.6:1 and 3.3:1, respectively) for 8 h. Relative measurements of cell death rate given by bisbenzimidazole and propidium iodide staining showed that 0.5 mM palmitate complexed with both 0.5% BSA or 1% BSA caused increased cell death. Palmitate:BSA at the ratio of 3.3:1 induced less cell death than the ratio of 6.6:1, which might be due to the higher toxicity of unbound free fatty acid. Bcl-2 overexpression promoted a partial protection against both 0.5 mM palmitate (0.5% BSA) (P=0.025) and 0.5 mM palmitate (1% BSA) treatments (P=0.029) (Fig. 1B). Bcl-2 overexpression tended to protect against the uncoupler FCCP, but this did not reach statistical significance. The overexpression of Bcl-2 was maintained through the 8-h-incubation with 0.5mM palmitate complexed with 0.5% BSA (Fig. 1C). The Bcl-2 overexpression-induced partial protection against 0.5 mM palmitate (0.5% BSA) was further confirmed by analysis of cleaved caspase 3 activation (Fig. 1D-E).

### **Palmitate-induced GADD153/CHOP induction was delayed by Bcl-2 overexpression.**

As an important event of palmitate-induced  $\beta$ -cell death, levels of the transcription factor GADD153/CHOP were analyzed at an interval of 2 hour during the 8 hours of palmitate exposure. The induction of GADD153/CHOP protein levels, which occurred after 6 h of palmitate exposure, was markedly delayed or counteracted by Bcl-2 (Fig. 2).

**Palmitate induced BimEL phosphorylation, which was attenuated by Bcl-2 overexpression.**

Palmitate has been shown to induce the proapoptotic BH3-only protein Bcl-2-like 11 (Bim) up-regulation [14] and Bim activity may be controlled by several posttranscriptional phosphorylation events at multiple serine and threonine residues [15]. For instance, BimEL, the longest splice form of Bim, can be phosphorylated by c-Jun N-terminal kinase (JNK) or MAPK-p38 at serine 65 to increase its proapoptotic activity [16, 17]. Using an anti-Bim antibody, we observed two bands of BimEL (Fig. 3B). The upper band is probably the phosphorylated form of BimEL as verified by phosphatase treatment (Fig. 3C). When the ratio of BimEL upper (phosphorylated) to lower (non-phosphorylated) bands was calculated, a significant decrease of BimEL phosphorylation was observed in BMG cells at 4 and 8 h of palmitate incubation (Fig. 3A).

**Palmitate induced a gradually decrease of mitochondrial membrane potential ( $\Delta\psi_m$ ), which was counteracted by Bcl-2.**

Partial mitochondrial uncoupling is induced by palmitate in pancreatic islets [3] and Bcl-2 overexpression can stabilize the  $\Delta\psi_m$  in cytokine treated RINm5F cells [18]. We therefore investigated whether Bcl-2 overexpression affected palmitate-induced alterations in JC-1 fluorescence. A 60 min exposure to palmitate promoted a small increase in  $\Delta\psi_m$  in both B45 and BMG cells (Fig. 4A), possibly due to an increase in mitochondrial fatty acid oxidation. However, at 3 - 8 h, the  $\Delta\psi_m$  was instead gradually decreased in response to palmitate. Bcl-2 overexpression counteracted the palmitate-induced loss of  $\Delta\psi_m$  at 3 and 8 h of palmitate exposure.

**Bcl-2 overexpression failed to counteract palmitate-induced ROS generation.**

The magnitude of the mitochondrial membrane potential is known to influence rates of mitochondrial ROS production [19]. We therefore analyzed ROS levels in response

to palmitate and Bcl-2 overexpression. Using the hydrogen peroxide and the hydroxyl radical sensor DCFH-DA, we observed increased ROS generation in response to a 2 h palmitate exposure and that this effect was even stronger at 3h in both B45 and BMG cells (Fig. 4B). Bcl-2 overexpression failed to counteract palmitate-induced ROS generation at all time points.

#### **Effects of Bcl-2 overexpression on palmitate-induced ER stress signaling.**

The Bcl-2 protein is known to modulate ER stress responses [20]. We therefore investigated the initiation of palmitate-induced ER stress in B45 and BMG cells. The activation of two ER stress sensors eIF-2 $\alpha$  and IRE1 $\alpha$  was analyzed by immunoblotting with phospho-specific antibodies at an interval of 1 or 2h during the palmitate exposure. We observed that the phosphorylation of IRE1 $\alpha$  was only weakly increased by palmitate and that this event was not affected by Bcl-2 overexpression (Fig. 5A-B). The phosphorylation of eIF2 $\alpha$  was also not significantly affected by Bcl-2 overexpression (Fig. 5C-D).

#### **Effects of palmitate on Lys221-p65 methylation in B45 and BMG cells.**

Recently, palmitate has been shown to induce NF- $\kappa$ B activation in human islets [21] and a mouse  $\beta$  cell line [22]. The activation of most p65-dependent genes has been reported to rely on lysine methylation of the p65 subunit [23, 24]. Therefore we performed a time course study of Lys221-p65 methylation to monitor palmitate-induced NF- $\kappa$ B activation in B45 and BMG cells. Cytokine incubation, IL-1 $\beta$  (20 ng/ml) and IFN- $\gamma$  (20 ng/ml) for 1 h, was used here as a positive control for NF- $\kappa$ B activation. Palmitate promoted in B45 cells, but not in BMG cells, a modest but significant increase in p65 methylation at 2, 4, 6 and 8 hour ( $P < 0.05$  using paired Student's t-test when comparing to time zero). A significant increase in BMG cell p65 methylation was observed at basal conditions and after 2 h of palmitate exposure when compared to B45 cells, which indicated an enhanced NF- $\kappa$ B activation through Bcl-2 overexpression. The augmented NF- $\kappa$ B pathway in BMG cells was still capable to be further activated as we observed higher methylation rates of p65 when exposed to cytokines for 1 h (Fig. 6).

## Discussion

We have presently investigated the impact of Bcl-2 on early mitochondrial and ER events induced by sodium palmitate in insulin-producing RINm5F cells. Because  $\beta$ -cell levels of Bcl-2 decrease in response to prolonged exposure periods of high palmitate concentrations [4] and because factors that protect against lipotoxicity, i.e. GLP-1 and hepatocyte growth factor, increase Bcl-2 expression [25, 26], it is likely that the anti-apoptotic Bcl-2 protein mediates a protective action against FFA-induced  $\beta$ -cell dysfunction and death. Indeed, we presently observed partial resistance against palmitate-induced cell death in cells overexpressing Bcl-2. However, the mechanisms by which Bcl-2 promotes this beneficial effect are not well characterized. Bcl-2 is known to primarily counteract proapoptotic mitochondrial events leading to a loss of  $\Delta\psi_m$ , mitochondrial permeability transition (MPT) and release of cytochrome c. Moreover, Bcl-2 is also found at the ER where it reduces ER stress-mediated apoptosis, possibly via counteraction of proapoptotic Bax/Bak and modulation of ER calcium homeostasis [20]. To address these possibilities, we investigated early palmitate-induced alterations in mitochondrial membrane potential and observed that Bcl-2 overexpression mediated protection against the gradual decrease in  $\Delta\psi_m$  starting at 3 h of palmitate exposure. Palmitate-induced loss of  $\Delta\psi_m$  was first reported by us in 1999 [3] and, more recently, it has been observed that palmitate, depending on dose and time of exposure, promotes both limited MPT opening, associated with impaired insulin secretion, and full MPT opening, associated with apoptosis [27]. Since Bcl-2 overexpression protected both at an early (3h) and at a late time point (8h) against palmitate-induced MPT, it is possible that Bcl-2 acts via inhibition of both limited and full MPT pore opening.

Sodium palmitate promotes  $\beta$ -cell production of ROS [3] and antioxidants protect, at least in part, against FFA-induced  $\beta$ -cell dysfunction in vitro and in vivo [28]. In the present investigation ROS production was stimulated already at 2h after addition of palmitate, and Bcl-2 overexpressing cells produced similar amounts of ROS as control cells. The source of palmitate-induced oxygen free radicals is not known, but may be of mitochondrial [29] and/or plasma membrane (NADPH-oxidase) [30] origin. In either case, our results suggest that palmitate-induced ROS production occurs prior and independently of the gradual decrease in mitochondrial  $\Delta\psi_m$ , indicating that the loss of  $\Delta\psi_m$  may have resulted from the increased ROS production. Interestingly, it

has been observed that Bcl-2 overexpression protects against AMPK-induced  $\beta$ -cell apoptosis without affecting production of oxygen radicals [31]. Furthermore, ROS are recognized activators of cytochrome c release and MPT [32]. Thus, it is possible that palmitate-induced ROS production causes opening of mitochondrial pores, and that this latter event is counteracted by Bcl-2.

We presently also report that Bcl-2 overexpression did not affect early palmitate-induced eIF-2 $\alpha$  and IRE1 $\alpha$  phosphorylation events. This argues against a protective role of Bcl-2 at the  $\beta$ -cell ER membrane and instead for the notion that ER stress occurs as a consequence of augmented ROS production, or some other upstream event. The exact role of palmitate-induced ER stress in  $\beta$ -cell death has presently not been investigated, but the finding that BMG cells were partially protected against palmitate-induced cell death, but not against early ER stress events, diminishes the putative importance of the ER-stress pathway in  $\beta$ -cell apoptosis.

GADD153/CHOP is thought to play a direct role in palmitate-induced  $\beta$ -cell apoptosis as siRNA-mediated knockdown of GADD153/CHOP has been observed to lower apoptosis rates [9]. We presently observed that Bcl-2 overexpression delayed or counteracted palmitate-induced GADD153/CHOP induction at 6-8h. This is a somewhat surprising finding because increased eIF-2 $\alpha$  phosphorylation is generally associated with subsequent expression of GADD153/CHOP [9]. However, it may also be that the palmitate-induced expression of GADD153/CHOP is to a large extent controlled by non-ER stress related pathways. For example, it has long been known that ceramide, Fas and JNK all promote GADD153/CHOP induction [33,34]. Furthermore, it has been reported that the anti-apoptotic transcription factor NF- $\kappa$ B acts as a repressor of the GADD153 gene [35]. Thus, as we presently observed enhanced NF- $\kappa$ B signaling in BMG cells, both at basal conditions and when treated with palmitate for 2h, it is possible that Bcl-2 attenuates GADD153/CHOP induction, at least in part, by maintaining a higher activity of NF- $\kappa$ B, and not by modulating ER stress events.

Bim has also been shown to be involved in palmitate-induced  $\beta$ -cell apoptosis [14], and we presently observed increased Bim phosphorylation in response to palmitate at 4-8h. It appears that JNK phosphorylates Bim at amino acid ser65 and that this event is associated with increased apoptosis [36]. As overexpression of Bcl-2 has been

shown to suppress JNK activation [37], it is possible that palmitate-induced BimEL phosphorylation is mediated, at least in part, by JNK activation. In similarity with GADD153/CHOP as discussed above, Bcl-2 may suppress Bim activation in  $\beta$ -cells by maintaining mitochondrial integrity and/or upregulating NF- $\kappa$ B activity.

In summary, we propose a hypothetical model in which palmitate induces the following chain of events; a) increased ROS production (2h), b) decreased mitochondrial membrane potential and increased ER stress (3-4h), c) augmented expression of GADD153/CHOP and increased Bim phosphorylation (4-8h), and d) cell death (8h). In addition, our results are compatible with the view that Bcl-2 protects  $\beta$ -cells against palmitate by maintaining mitochondrial membrane integrity and upregulating NF- $\kappa$ B activity, and not by affecting early ROS production and ER-stress signaling events. Thus, mitochondria may play a pivotal role in the control of  $\beta$ -cell survival and function in Type 2 diabetes.

### **Acknowledgements**

This work was supported in part by the Swedish Diabetes Association, the family Ernfors Fund, and the Novo-Nordisk Foundation.

**Declaration of interests:** The authors declare no conflicts of interests.

### **References**

1. Cnop M, Hannaert JC, Hoorens A, Eizirik DL, Pipeleers DG. Inverse relationship between cytotoxicity of free fatty acids in pancreatic islet cells and cellular triglyceride accumulation. *Diabetes*. 2001; 50:1771-7.
2. Cunha DA, Hekerman P, Ladriere L, Bazarra-Castro A, Ortis F, Wakeham MC, et al. Initiation and execution of lipotoxic ER stress in pancreatic beta-cells. *J Cell Sci*. 2008; 121:2308-18.
3. Carlsson C, Borg LA, Welsh N. Sodium palmitate induces partial mitochondrial uncoupling and reactive oxygen species in rat pancreatic islets in vitro. *Endocrinology*. 1999; 140:3422-8.
4. Lupi R, Dotta F, Marselli L, Del Guerra S, Masini M, Santangelo C et al. Prolonged exposure to free fatty acids has cytostatic and pro-apoptotic effects on human pancreatic islets: evidence that beta-cell death is caspase mediated, partially dependent on ceramide pathway, and Bcl-2 regulated. *Diabetes*. 2002; 51:1437-42.
5. Gwiazda KS, Yang TL, Lin Y, Johnson JD. Effects of palmitate on ER and cytosolic Ca<sup>2+</sup> homeostasis in beta-cells. *Am J Physiol Endocrinol Metab*. 2009; 296:E690-701.

6. de Pablo MA, Susin SA, Jacotot E, Larochette N, Costantini P, Ravagnan L, et al. Palmitate induces apoptosis via a direct effect on mitochondria. *Apoptosis*. 1999; 4:81-7.
7. Bhatt K, Feng L, Pabla N, Liu K, Smith S, Dong Z. Effects of targeted Bcl-2 expression in mitochondria or endoplasmic reticulum on renal tubular cell apoptosis. *Am J Physiol Renal Physiol*. 2008; 294:F499-507.
8. Lam M, Dubyak G, Chen L, Nunez G, Miesfeld RL, Distelhorst CW. Evidence that BCL-2 represses apoptosis by regulating endoplasmic reticulum-associated Ca<sup>2+</sup> fluxes. *Proc Natl Acad Sci U S A*. 1994; 91:6569-73.
9. Eizirik DL, Cardozo AK, Cnop M. The role for endoplasmic reticulum stress in diabetes mellitus. *Endocr Rev*. 2008; 29:42-61.
10. Saldeen J. Cytokines induce both necrosis and apoptosis via a common Bcl-2-inhibitable pathway in rat insulin-producing cells. *Endocrinology*. 2000; 141:2003-10.
11. Karasuyama H, Melchers F. Establishment of mouse cell lines which constitutively secrete large quantities of interleukin 2, 3, 4 or 5, using modified cDNA expression vectors. *Eur J Immunol*. 1988; 18:97-104.
12. Deng G, Podack ER. Suppression of apoptosis in a cytotoxic T-cell line by interleukin 2-mediated gene transcription and deregulated expression of the protooncogene bcl-2. *Proc Natl Acad Sci U S A*. 1993; 90:2189-93.
13. Smiley ST, Reers M, Mottola-Hartshorn C, Lin M, Chen A, Smith TW, et al. Intracellular heterogeneity in mitochondrial membrane potentials revealed by a J-aggregate-forming lipophilic cation JC-1. *Proc Natl Acad Sci U S A*. 1991; 88:3671-5.
14. Akazawa Y, Cazanave S, Mott JL, Elmi N, Bronk SF, Kohno S, et al. Palmitoleate attenuates palmitate-induced Bim and PUMA up-regulation and hepatocyte lipoapoptosis. *J Hepatol*. 2010; 52:586-93.
15. Hubner A, Barrett T, Flavell RA, Davis RJ. Multisite phosphorylation regulates Bim stability and apoptotic activity. *Molecular cell*. 2008; 30:415-25.
16. Becker EB, Howell J, Kodama Y, Barker PA, Bonni A. Characterization of the c-Jun N-terminal kinase-BimEL signaling pathway in neuronal apoptosis. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2004; 24:8762-70.
17. Cai B, Chang SH, Becker EB, Bonni A, Xia Z. p38 MAP kinase mediates apoptosis through phosphorylation of BimEL at Ser-65. *J Biol Chem*. 2006; 281:25215-22.
18. Barbu A, Welsh N, Saldeen J. Cytokine-induced apoptosis and necrosis are preceded by disruption of the mitochondrial membrane potential ( $\Delta\psi(m)$ ) in pancreatic RINm5F cells: prevention by Bcl-2. *Mol Cell Endocrinol*. 2002; 190:75-82.
19. Newsholme P, Haber EP, Hirabara SM, Rebelato EL, Procopio J, Morgan D, et al. Diabetes associated cell stress and dysfunction: role of mitochondrial and non-mitochondrial ROS production and activity. *J Physiol*. 2007; 583:9-24.
20. Hetz CA. ER stress signaling and the BCL-2 family of proteins: from adaptation to irreversible cellular damage. *Antioxid Redox Signal*. 2007; 9:2345-55.
21. Igoillo-Esteve M, Marselli L, Cunha DA, Ladrière L, Ortis F, Grieco FA, et al. Palmitate induces a pro-inflammatory response in human pancreatic islets that

- mimics CCL2 expression by beta cells in type 2 diabetes. *Diabetologia*. 2010; 53:1395-405.
22. Choi HJ, Hwang S, Lee SH, Lee YR, Shin J, Park KS, et al. Genome-wide identification of palmitate-regulated immediate early genes and target genes in pancreatic beta-cells reveals a central role of NF-kappaB. *Molecular biology reports*. 2012; 39:6781-9.
  23. Lu T, Jackson MW, Wang B, Yang M, Chance MR, Miyagi M, et al. Regulation of NF-kappaB by NSD1/FBXL11-dependent reversible lysine methylation of p65. *Proc Natl Acad Sci U S A*. 2010; 107:46-51.
  24. Mokhtari D, Li T, Lu T, Welsh N. Effects of Imatinib Mesylate (Gleevec) on human islet NF-kappaB activation and chemokine production in vitro. *PLoS one*. 2011; 6:e24831.
  25. Buteau J, El-Assaad W, Rhodes CJ, Rosenberg L, Joly E, Prentki M. Glucagon-like peptide-1 prevents beta cell glucolipotoxicity. *Diabetologia* 2004; 47:806-15.
  26. Santangelo C, Matarrese P, Masella R, Di Carlo MC, Di Lillo A, Scazzocchio B, et al. Hepatocyte growth factor protects rat RINm5F cell line against free fatty acid-induced apoptosis by counteracting oxidative stress. *J Mol Endocrinol*. 2007; 38:147-58.
  27. Koshkin V, Dai FF, Robson-Doucette CA, Chan CB, Wheeler MB. Limited mitochondrial permeabilization is an early manifestation of palmitate-induced lipotoxicity in pancreatic beta-cells. *J Biol Chem*. 2008; 283:7936-48.
  28. Oprescu AI, Bikopoulos G, Naassan A, Allister EM, Tang C, Park E, et al. Free fatty acid-induced reduction in glucose-stimulated insulin secretion: evidence for a role of oxidative stress in vitro and in vivo. *Diabetes*. 2007; 56:2927-37.
  29. Maestre I, Jordan J, Calvo S, Reig JA, Ceña V, Soria B, et al. Mitochondrial dysfunction is involved in apoptosis induced by serum withdrawal and fatty acids in the beta-cell line INS-1. *Endocrinology*. 2003; 144:335-45.
  30. Michalska M, Wolf G, Walther R, Newsholme P. Effects of pharmacological inhibition of NADPH oxidase or iNOS on pro-inflammatory cytokine, palmitic acid or H<sub>2</sub>O<sub>2</sub>-induced mouse islet or clonal pancreatic beta-cell dysfunction. *Biosci Rep* 2010; 30:445-53.
  31. Cai Y, Martens GA, Hinke SA, Heimberg H, Pipeleers D, Van de Castele M. Increased oxygen radical formation and mitochondrial dysfunction mediate beta cell apoptosis under conditions of AMP-activated protein kinase stimulation. *Free Radic Biol Med*. 2007; 42:64-78.
  32. Costantini P, Chernyak BV, Petronilli V, Bernardi P. Modulation of the mitochondrial permeability transition pore by pyridine nucleotides and dithiol oxidation at two separate sites. *J Biol Chem*. 1996; 271:6746-51.
  33. Kultz D, Madhany S, Burg MB. Hyperosmolality causes growth arrest of murine kidney cells. Induction of GADD45 and GADD153 by osmosensing via stress-activated protein kinase 2. *J Biol Chem*. 1998; 273:13645-51.
  34. Brenner B, Koppenhoefer U, Weinstock C, Linderkamp O, Lang F, Gulbins E. Fas- or ceramide-induced apoptosis is mediated by a Rac1-regulated activation of Jun N-terminal kinase/p38 kinases and GADD153. *J Biol Chem*. 1997; 272:22173-81
  35. Schapansky J, Olson K, Van Der Ploeg R, Glazner G. NF-kappaB activated by ER calcium release inhibits Abeta-mediated expression of CHOP protein:

- enhancement by AD-linked mutant presenilin 1. *Exp Neurol.* 2007; 208:169-76.
36. Santin I, Moore F, Colli ML, Gurzov EN, Marselli L, Marchetti P, et al. PTPN2, a candidate gene for type 1 diabetes, modulates pancreatic beta-cell apoptosis via regulation of the BH3-only protein Bim. *Diabetes.* 2011; 60:3279-88.
37. Koo MS, Kwo YG, Park JH, Choi WJ, Billiar TR, Kim YM. Signaling and function of caspase and c-jun N-terminal kinase in cisplatin-induced apoptosis. *Molecules and cells.* 2002; 13:194-201.

## Figure legends

**Fig.1.** Expression of Bcl-2 in neo (B45) and bcl-2 (BMG) transfected RINm5F cell clones and effects of palmitate and FCCP on B45 and BMG transfected cell viability. (A) Expression of Bcl-2 in BMG and B45 cell clones. (B) Effects of palmitate and FCCP on B45 and BMG transfected cell viability. RIN cell clones were incubated with 0.5 mM palmitate (0.5% BSA or 1% BSA + 1% FBS) or 1 µg/ml FCCP for 8 h. Results are means ± S.E.M. for five separate experiments. \* denotes P<0.05 using paired Student's t-test when comparing vs corresponding control. (C) One representative immunoblot showing Bcl-2 expression during the 8-h-incubation with 0.5 mM palmitate (0.5% BSA). (D) One representative immunoblot showing cleaved caspase 3 levels from five experiments. (E) Mean optical density measurements of the immunoblots of cleaved caspase 3. The results are expressed as percentages of the control (B45 cells; Time zero) and shown as means ± S.E.M for five separate experiments. \* denotes P<0.05 using paired Student's t-test.

**Fig.2.** Effects of palmitate on GADD 153 (CHOP) expression in B45 and BMG cells. RIN cell clones were incubated with 0.5 mM palmitate (0.5% BSA) for 8 h. (A) Mean optical density measurements of the immunoblots of CHOP. Protein values were normalized to amidoblack staining of total protein. The results are expressed as percentages of the control (B45 cells; Time zero) and shown as means ± S.E.M. \* denotes P<0.05 using paired Student's t-test when comparing protein induction in BMG cells with corresponding B45 cells. (B) One representative immunoblot of CHOP expression from five experiments is shown.

**Fig.3.** Effects of palmitate on ratio of BimEL upper to lower band intensity in B45 and BMG cells. Cell clones were incubated with cytokines (IL-1β 20 ng/ml and IFN-γ 20 ng/ml) for 1 h or 0.5 mM palmitate (0.5% BSA) for 2-8 h. After the treatments, the cells were lysed and analyzed by immunoblotting with anti-Bim antibody. BimEL is the longest splice form (23 kDa) of Bim. Two bands of BimEL were observed and the upper band is probably the phosphorylated form of BimEL as verified by phosphatase treatment. (A) Ratio of BimEL upper (phosphorylated) to lower (non-phosphorylated) bands. The results are expressed as percentages of the control (B45 cells; Time zero) and shown as means ± S.E.M. \* denotes P<0.05 and # denotes P<0.01 when comparing BMG cells with corresponding B45 cells. (B) One representative

immunoblot of BimEL phosphorylation from 4 experiments is shown. (C) One representative immunoblot of phosphatase treatment,

**Fig.4.** (A) Effects of palmitate on JC-1 fluorescence in B45 and BMG cells. BMG and B45 cells were cultured for 1-8 h with 0.5 mM palmitate, 0.5% BSA. Data are expressed as the 585/530 nm fluorescence ratio normalized to the control of each time point. Results are means  $\pm$  S.E.M for three separate experiments. # denotes  $P < 0.01$  using paired Student's t-test when comparing uncoupling rate of BMG cells with corresponding B45 cells. The basal JC-1 fluorescence ratio at time zero is B45:  $100 \pm 5\%$  and BMG:  $86 \pm 22\%$ . (B) Effects of palmitate on DCF fluorescence in B45 and BMG cells. BMG and B45 cells were cultured for 1, 2 or 3 h with 0.5 mM palmitate (0.5% BSA). Data are expressed as 530 nm fluorescence normalized to the control of each time point. Results are means  $\pm$  S.E.M for three separate experiments. The basal signal of DCF fluorescence at time zero is B45:  $100 \pm 8\%$  and BMG:  $97 \pm 8\%$ .

**Fig.5.** Effects of palmitate on Phospho-IRE1 $\alpha$  and Phospho-eIF2 $\alpha$  expression in B45 and BMG cells. RIN cell clones were incubated with 0.5 mM palmitate (0.5% BSA) for 6 or 8 h. Western blots were quantified and protein values were normalized to amidoblack staining of total protein. Mean optical density measurements of the immunoblots for Phospho-IRE1 $\alpha$  (A) and Phospho-eIF2 $\alpha$  (C). The results are expressed as percentages of the control (B45 cells; Time zero) and shown as means  $\pm$  S.E.M. (B) One representative immunoblot of Phospho-IRE1 $\alpha$  expression from five experiments is shown. (D) One representative immunoblot of Phospho-eIF2 $\alpha$  expression from six experiments is shown.

**Fig.6.** Effects of palmitate on Lys221-p65 methylation in B45 and BMG cells. Cell clones were incubated with cytokines (IL-1 $\beta$  20 ng/ml and IFN- $\gamma$  20 ng/ml) for 1h or 0.5 mM palmitate (0.5% BSA) for 2-8 h. After the treatments, the cells were lysed and analyzed by immunoblotting with anti-Methyl-Lys221-p65 antibody [23]. Immunoblots were quantified and protein values were normalized to amidoblack staining of total protein. (A) Cytokines and palmitate induced methylation of p65. The results are expressed as percentages of the control (B45 cells; Time zero) and shown as means  $\pm$  S.E.M. \* denotes  $P < 0.05$  and # denotes  $P < 0.01$  using paired Student's t-test when comparing BMG cells with corresponding B45 cells. (B) One representative immunoblot of Lys221-p65 methylation from 7 experiments is shown.