

Social Attention: Developmental Foundations and Relevance for Autism Spectrum Disorder

Terje Falck-Ytter, Johan Lundin Kleberg, Ana Maria Portugal, and Emilia Thorup

ABSTRACT

The use of the term “social attention” (SA) in the cognitive neuroscience and developmental psychopathology literature has increased exponentially in recent years, in part motivated by the aim to understand the early development of autism spectrum disorder (ASD). Unfortunately, theoretical discussions around the term have lagged behind its various uses. Here, we evaluate SA through a review of key candidate SA phenotypes emerging early in life, from newborn gaze cueing and preference for face-like configurations to later emerging skills such as joint attention. We argue that most of the considered SA phenotypes are unlikely to represent unique socioattentional processes and instead have to be understood in the broader context of bottom-up and emerging top-down (domain-general) attention. Some types of SA behaviors (e.g., initiation of joint attention) are linked to the early development of ASD, but this may reflect differences in social motivation rather than attention per se. Several SA candidates are not linked to ASD early in life, including the ones that may represent uniquely socioattentional processes (e.g., orienting to faces, predicting others’ manual action goals). Although SA may be a useful superordinate category under which one can organize certain research questions, the widespread use of the term without proper definition is problematic. Characterizing gaze patterns and visual attention in social contexts in infants at elevated likelihood of ASD may facilitate early detection, but conceptual clarity regarding the underlying processes at play is needed to sharpen research questions and identify potential targets for early intervention.

<https://doi.org/10.1016/j.biopsych.2022.09.035>

Looking at Hugues Merle’s painting *Contes enfantines* (Figure 1), few would disagree that social attention (SA) occurs in the scene. The younger children look at the older girl’s face with a magnetic intensity, completely absorbed by the words coming from her mouth, her gaze, and her facial expressions. The term “social attention” is typically used to describe phenomena linked to selective attention to eyes and faces, nonverbal social communication, and joint attention. The term is also commonly used when describing differences in looking behaviors in people with autism spectrum disorder (ASD) (1), and atypical SA has been suggested to represent a potential biomarker for the condition (2). A common assumption is that atypicalities in SA very early in life may be central for understanding early developmental trajectories in children with ASD (3). While looking in social contexts could potentially be facilitated by automatized SA processes in typical development, it may draw on different processes in ASD. This potential fundamental link to ASD and the general use of eye-tracking technology in (social) cognitive developmental neuroscience may explain why the use of the term “social attention” appears to be increasing exponentially in the literature (4).

Although theoretical discussions of the term “social attention” are scarce (4), 2 distinct views regarding the nature of SA can be identified in the existing literature:

Hypothesis A. SA is a domain-specific attentional process, distinct from other attentional processes. This view is implicated by research suggesting that it builds on highly

specialized brain circuits (4,5), i.e., as a social parallel to the domain-general attentional networks that have been identified over the last few decades (6,7). A more extreme version of this hypothesis is that SA is both domain specific and unitary (across many ostensibly different SA phenotypes) (4), a possibility we will return to in the Discussion.

Hypothesis B. Attention allocation to social objects, events, and scenes can be explained by general attentional processes (Figure 2). These involve 2 interlinked types: exogenous (stimulus driven, reflexive) and endogenous (goal-directed, voluntary) orienting (8). Furthermore, attentional mechanisms that are spatially unselective (e.g., general alerting) can be differentiated from visuospatial orienting (6). In infancy, many looking behaviors reflect a blend of exogenous and emerging endogenous processes (9).

These 2 views represent the extreme ends of a spectrum of possibilities, including that SA becomes distinct from other attentional processes over time, driven by experience and interaction with social stimuli (4,10,11).

To improve clarity around these contrasting views on SA, this review presents and discusses specific phenotypes that have been termed SA in the past (hereafter referred to as candidate SA phenotypes) while briefly considering available evidence supporting either hypothesis A or hypothesis B described above for the phenotype in question. The candidates are selected based on their prevalence in the ASD and



Figure 1. *Contes enfantines* by Hugues Merle (1823–1881).

developmental cognitive neuroscience literature. Because SA is thought to be tightly linked to ASD, which develops early in childhood, our focus is on developmental data from the first years of life. We also summarize the link, or lack thereof, to ASD early in life for each candidate SA phenotype. Whether a single SA candidate is affected in ASD is not in itself informative for evaluating hypothesis A versus hypothesis B.

However, looking at the emerging pattern across several SA candidates could be informative for understanding the nature of SA (e.g., its unity) and ASD. If ASD is fundamentally linked to SA, we expect that the candidate SA phenotypes with the most support for hypothesis A will be particularly affected in infants who develop ASD symptoms.

Studies of SA have been criticized for their reliance on 2-dimensional stimuli shown on a computer screen (12–14). In this review, we also illustrate how researchers are increasingly using modern technology to assess infants in more ecologically valid settings to try to circumvent this limitation (15).

SA: THE CANDIDATES

Orienting to Faces and Preference for Social Information

The term social attention is often used when describing how individuals distribute their gaze to social versus nonsocial stimuli. Given our focus on early development in ASD, we have selected looking to faces, eyes, and mouths, but there are other examples of stimuli that belong to this category, including preferential looking to biological motion [for which the link to ASD below age 2 years is largely unexplored; see (3,17,18)] and the actions of others (19).

Candidate 1: Basic Orienting to Face-like Patterns.

There is a robust body of evidence demonstrating a fast capture of attention by face-like patterns (Figure 3A) at birth and throughout the life span [e.g., (20–24)]. Such face orienting is reflexive and exogenously driven (22,23) and is thought to have a canalizing role in cortical specialization and perceptual narrowing (9,25–27).

It has been argued that newborn face orienting is the product of a domain-general process based on low-level stimulus characteristics [e.g., (25,28–32)], yet the prevailing view is that it represents a domain-specific social bias [e.g.,




Figure 2. How social is social attention (SA)? The use of the term in the developmental and autism spectrum disorder literature often implies a specialized process for SA in the brain (central dashed box). According to an alternative view, behavioral instances of SA (rightmost box) can be explained by domain-general attentional processes (bottom, top). The figure provides a simplified overview of the main factors behind SA behaviors. It is not intended to provide a full overview of all possible links (e.g., bidirectional).




Figure 3. The many faces of social attention. **(A)** Human newborns prefer to look at stimuli with patterns that resemble the basic configuration of high-contrast areas of a face (left). Whether newborns with later autism spectrum disorder (ASD) display similar attentional patterns is not yet known. **(B)** Responding to and initiating joint attention, assessed in a live eye-tracking paradigm. [Reproduced with permission from Nyström *et al.* (49).] While children with ASD showed atypicalities in initiating joint attention (IJA), their tendency to follow gaze was like typical infant performance (responding to joint attention [RJA]). **(C)** Predictive eye movements used in action observation. When observing a model moving objects to a container, 6-month-old infants reactively track the moving object/hand. In contrast, 12-month-old infants and adults, who are able to perform such actions themselves, use goal-directed predictive eye movements, which is believed to tax their own motor system and representations of similar actions (113). The colored lines show the direction of eye movements when leaving the object area (right) and entering the goal area (left), respectively, illustrating the striking similarity between 12-month-old infants and adults, who do not follow the moving hand/object, but instead move gaze toward the goal area (bucket). Most data indicate that this basic predictive ability is intact in children with ASD (119). Data are replotted based on Falck-Ytter *et al.* (111). **(D)** Attending to complex and dynamic social scenes involving a nonverbal video-recorded interaction between 2 children. In the video, the girl refuses to give the object to the boy despite his request. The top row shows the full stimuli, and the bottom row shows how typically developing 6-year-old children tend to attend to this scene. The final frame (**D3**) illustrates how virtually all typically developing children quickly moved their gaze to the face of the girl after the boy's (neglected) request. The tendency to do so was significantly weaker in ASD (area under the curve > 0.85). [Reproduced and adapted with permission from Falck-Ytter *et al.* (143).] This result indicates that analyses that take the timing of gaze shifts into account may provide insights that are not captured by aggregated looking time measures.

(26,33)] hypothesized to be generated by a subcortical processing route that includes the superior colliculus, pulvinar, and amygdala complex (26). The existence of a specialized and very early-emerging face orienting mechanism is supported by one report of preferential head turns to face-like stimuli in fetuses during the third trimester [(34); but see (35)] and by a recent finding that newborn brain responses to face-like patterns recruit an adult-like face-specific cortical circuit (36).

Available evidence suggests that this basic face bias is no different in infants with a later ASD diagnosis (37) or in autistic adult individuals (22), although one study found that it was decreased in preschoolers with ASD (18). In the infant study, Elsabbagh *et al.* (37) measured 6- to 10-month-old infants' tendency to orient to faces among nonface distractors presented on a computer screen. All groups, including infants who later met the criteria for an ASD diagnosis, tended to move their gaze to faces in about 50% of trials, high above the level

expected by random gaze allocation (20%). Notably, no study has examined whether face orienting is typical in newborns with later ASD.

Candidate 2: Preferential Looking to Naturalistic Faces in More Complex Visual Displays. From around 4 to 6 months of age, infants' sustained face preference also reflects top-down control (37–40). Thus, older infants' face preference and consequential emerging face expertise are the products of both unique SA orienting biases and multiple other attentional and executive abilities (26,27,29,41). Face preference (when in competition with nonsocial objects) already has a substantial contribution from genetic factors at 5 months [heritability of 0.46, (42)], implying that infants select social versus nonsocial input partly due to their genetic predispositions (43).

Looking at nonfacial body parts and nonsocial objects and events represents adaptive behaviors that are expected to dominate during certain age periods (18,44). This complexity may explain the mixed findings with regard to face preference in ASD. Some studies have shown that infants and toddlers later diagnosed with ASD look at faces longer than infants and toddlers in control groups do (37,45,46), with the investigators hypothesizing that this indicates slower maturation of attention control in ASD. Other studies based on screen-based eye tracking and video recordings from homes (47,48) have suggested shorter looking time to faces during the first 2 years of life in infants with later ASD, while a recent live eye-tracking study found no clear group differences in terms of face preference (49).

Candidate 3: Preference for Specific Facial Parts (Eyes and Mouth). A bias to attend to faces with direct as compared to averted gaze is seen shortly after birth (21), and preferential looking to the eyes over other facial features is seen throughout the first half of the first year (50–52). Neurophysiological responses from infants elicited during the earliest time stages of visual attention that are sensitive to the presence of whole faces are equally, or even more strongly, elicited by images of eyes alone or among nonsocial objects (53,54). While these results demonstrate that eyes are prioritized during visual processing, it is not clear whether this effect is driven by unique attentional mechanisms underlying attention to eyes rather than by orienting to faces per se (i.e., candidate 1).

Toward the end of their first year when infants are in an intense phase of language learning, they tend to increase their looking to the mouth (vs. eyes), which provides rich audio-visual speech information (50,51,55,56). During the second year of life when infants have gained some native language experience and narrowed their perceptual expertise, their preference for eyes increases again. Individual differences in mouth (vs. eyes) preference are substantial, and recent twin data suggest that this variability has a moderate-to-high contribution from genetic influences at 5 months (52) and at 21 months (57) of age.

Regarding infants later diagnosed with ASD, results are mixed, probably reflecting age differences as well as other study-specific factors (47,58). A small effect of decreased looking to the mouth has been found in a meta-analysis of attention in children with diagnosed ASD (59). Eye- versus mouth-looking has not been evaluated in an ecologically valid context in infants at elevated risk for ASD [but see ref (60) for a study of school-aged children].

Phenotypes Linked to Joint Attention

Joint attention refers to the triadic sharing of attention between 2 individuals toward a common object or event (61,62). Here we review 3 SA candidates of increasing complexity that are linked to this concept.

Candidate 4: Gaze Cueing. Gaze cueing refers to the tendency to orient faster to peripheral stimuli appearing in locations previously cued by other people's eyes. Gaze cueing is present across the life span (including in newborns) (63–66) and is typically seen as an exogenous form of attention cueing subserved by activity in the anterior superior temporal sulcus (67). Gaze cueing has been suggested to indicate the presence of a specific sensitivity to spatial information conveyed by eyes (63). This view was challenged by the finding that arrows produce a similar cueing effect (68). However, later studies have indicated that attention cued by gaze is more strongly reflexive (and less amenable to top-down processes) than attention cued by arrows (69–71), thus supporting the hypothesis that eye gaze has a special status compared with nonbiological cues and is potentially supported by a distinct neural network (72).

To our knowledge, gaze cueing in infants who later receive an ASD diagnosis has not yet been investigated. In children with a confirmed ASD diagnosis, results are mixed (73–79). Both Senju *et al.* (80) and Chawarska *et al.* (81) found typical gaze cueing in autistic children but reported more subtle differences between these children and typically developing children in terms of their responses to gaze cues versus nonbiological cues (e.g., arrows). This may reflect dissociable underlying processes (82), a view supported by a functional magnetic resonance imaging study (83).

Candidate 5: Gaze Following. Gaze following refers to the tendency to respond to changes in others' gaze by looking in the same direction. In contrast to gaze cueing, gaze following is not exogenously triggered by changes in peripheral stimulation. Gaze following emerges around 3 to 4 months of age (84), coinciding with a period of intense development of cortical control over eye movements (85). Gaze following then gradually becomes faster (86) and more accurate (49) during infancy and early childhood. Gaze following plays an important role in language development and learning (87–89). In adulthood, gaze following is associated with activity in several areas of the medial prefrontal cortex as well as the right posterior temporal sulcus (90–92). Gaze following in infancy is associated with partly similar brain activation patterns (93,94).

The ability to follow gaze reflects several processes, such as attention to eyes and faces and the ability to disengage attention. It has been argued that gaze following in very young infants may rely on the perceived motion of the head or pupils [e.g., (66,95)] and that nonbiological moving objects may evoke a similar response (96). One recent study indicated that young infants' gaze following is affected by both social cues (direction of looking) and nonsocial cues (general movement), but that gaze following comes to rely more on social cues with increasing age (97).

Live observational studies of gaze following early in life in ASD tend to find group differences (98–101), but these findings are not replicated in eye tracking studies using prerecorded

stimuli (102–105). This suggests that screen-based methods could fail to capture some important element of the interaction (106). However, using a live eye-tracking approach (i.e., real interaction with another person) (Figure 3B), Nyström *et al.* (49) recently found that infants with later ASD were equally likely to follow gaze as infants without subsequent ASD. Thus, impaired gaze following per se does not stand out as a clear early marker of ASD.

Candidate 6: Initiation of Joint Attention. Infants' gaze is not just an information gathering device. Because eye movements are visible to others, the use of gaze is also a key component of their nonverbal communication, e.g., with parents (43,49,107). Toward the end of the first year, infants begin to use gaze behaviors that can initiate joint attention episodes with others (49). Specifically, around 8 to 9 months of age, infants start looking back and forth between an object that has caught their attention and another person, a behavior referred to as alternating gaze (88). At 10 to 13 months, more overt behaviors, such as pointing, start to emerge (88,108). Initiation of joint attention (at least in adulthood) specifically recruits reward-related brain areas linked to social motivation, such as the ventral striatum, suggesting involvement of top-down attentional control (92). The same brain areas are not typically activated during more reactive gaze following.

Children with ASD use gestures such as pointing and showing less often than typically developing children [e.g., (98)], and infants later diagnosed with ASD have been shown to engage less in alternating gaze than typically developing infants as early as 8 to 10 months of age (49,109). Compared with gaze following, impairment in initiation of joint attention in children with ASD appears to be both more pronounced and long-lasting (110).

Attention During Observation of Other People's Manual Actions

Although gaze performance during action observation can be seen in part as a special case of preferential looking to social information (similar to preference for faces; see above), evidence suggests that manual actions give rise to a special type of eye movements that deserve a separate discussion.

Candidate 7: Predictive, Goal-Directed Eye Movements. Real-time, predictive eye movements to goals during action observation emerge toward the end of the first year in infancy (111,112) (Figure 3C) and, at least in adults, depend on a somatotopic recruitment of the observer's motor system (113). It is believed that observers recruit their own action plans during action observation to predict other people's action goals (114). According to one interpretation of the literature, this action mirroring is a highly automatic process ("direct matching") not requiring top-down control (115,116). If correct, these eye movements may reflect low-level SA processes that are triggered specifically by hand-object interactions. According to an alternative re-enactment view, motor recruitment happens after the observer has understood an agent's intention (115). However, the latter view cannot explain why predictive eye movements are typically not observed when infants

look at self-propelled objects moving toward target objects, even though they do attribute goals to such events (116). Although both predictive abilities in general and perception-action mirroring have been proposed to be atypical in ASD (117,118), the available data from young children and infants at elevated risk do not indicate substantial group differences, if any (119–121). It appears as if the distinction between live versus screen-based eye tracking is not crucial for this phenotype because it is observable in both contexts in infancy and adulthood (111,112,114).

DISCUSSION

In the literature on basic attentional processes, careful experimentation over many years indicates that the underlying neural systems are anatomically separate from other information processing systems and that multiple distinct attentional systems exist (6). How SA fits into this picture has not been altogether clear. Our review indicates that there is no unequivocal support for the involvement of unique socioattentional processes (see hypothesis A) (Table 1) for any of the SA candidates. For example, there is no consensus in the field that joint attention behaviors such as gaze following and alternating gaze (Figure 3B), very often referred to as reflecting SA in the literature, involve distinct social attention processes in this sense. In fact, during joint attention and when looking at complex social scenes more generally (e.g., Figure 3D), top-down influences (reflecting past experiences, social motivation, mentalizing, and other forms of social cognition) are likely to play a major role in moment-to-moment gaze allocation. This may explain why visual saliency models (122) tend to fail to predict attention in such contexts (123). However, for some SA behaviors there is partial support for domain specificity and uniqueness at a mechanistic level. Prediction of other people's action goals (Figure 3B) appears to reflect a unique [at both behavioral and brain levels (113)] process linked to perceptual selection that requires social input to be triggered (111,114).

Table 1. Overview of Reviewed Social Attention Phenotypes

Candidate Number	Candidate	Support for Hypothesis A vs. B	Atypical in Infants With Later ASD?
1	Basic orienting to face-like patterns	Leaning A	No (but not yet studied in newborns)
2	Preferential looking to naturalistic faces in more complex visual displays	Leaning B	Mixed findings
3	Preference for specific facial parts (eyes and mouth)	Leaning B	Mixed findings (age-dependent)
4	Gaze cueing	Inconclusive	Not yet studied in infancy
5	Gaze following	Inconclusive	Mixed findings (methods-dependent)
6	Initiation of joint attention	Leaning B	Yes
7	Predictive, goal-directed eye movements	Leaning A	No

ASD, autism spectrum disorder.

Similarly, newborn orienting to face-like configurations (Figure 3A) is considered by many [but not all; see (25,28–32)] to constitute a uniquely social attentional process operating early in life (26,33).

How fundamental are SA differences in the development of ASD? Currently, there is no evidence available to suggest that SA behaviors that may be considered strong candidates in a mechanistic sense, such as action prediction (116) or orienting to faces (37), are atypical in infants with emerging ASD. In fact, the relatively few studies that have reported evidence supporting atypical SA behaviors (other than initiation of joint attention; candidate 6) during the first year of life in infants later diagnosed with ASD tend to have modest effect sizes or small sample sizes (47,58,124). However, it is worth noting that some SA behaviors, such as gaze cueing, orienting to biological motion, and orienting to faces, have not yet been probed in relation to ASD during the first few months of life. The fact that initiation of joint attention in infancy shows a relatively strong relation to later ASD could be consistent with the social motivation theory of ASD, which states that interaction and communication impairments in ASD stem from a reduced intrinsic reward value of social interaction for individuals with ASD as compared with typically developing individuals (125). Although it is questionable whether atypical gaze behaviors during social interaction in ASD reflect differences in domain-specific attentional functions per se, they may still be important because they could alter child-parent interaction dynamics and developmental trajectories.

We have focused our review mainly on the question of whether specific SA behaviors reflect unique socioattentional processes, i.e., processes that are distinct from general attention and from nonattentional processes such as social motivation and social cognition (Figure 2). An arguably more extreme interpretation of SA is that it is also unitary (4). The wide range of behaviors referred to as SA [ranging from basic biases, present at or even before birth (34), to highly dynamic gaze behaviors (Figure 3)], their different developmental trajectories, and their differential links to ASD speak against strong claims of unity (but notably, a differential link to ASD for early vs. later emerging SA behaviors does not per se exclude the possibility that SA is unitary because it could be linked to the timing of gene expression in children with the condition). It is likely that correlations across different SA phenotypes would emerge because they are similarly modulated by the social context in the same individual, e.g., via social motivation or general attention.

Despite the reservations mentioned previously regarding the SA category, one could argue that it is a useful umbrella term under which one can organize a set of related research questions. For example, some types of early SA phenotypes are associated with later social developmental outcomes (42,49) and thus are important to study irrespective of the exact biological/psychological underpinnings. However, more conceptual and methodological clarity is needed for progress at both a basic and an applied level. Studying looking at social objects and events with an eye tracker does not necessarily mean that what has been measured is best described as SA.

It is likely that differences in infants' gaze behaviors in social settings partly reflect genetic differences in the population (42,43,52,57), meaning that selection of visual input to some

extent is rooted in genetics (active gene-environment correlation). Furthermore, because the eyes are visible to others, these differences could in turn affect caregiver responses toward their offspring (evocative gene-environment interplay).

Lately, interest in SA has spread to many other neurodevelopmental and child psychiatric disorders partly phenotypically overlapping with ASD, including social anxiety disorder (126,127), attention-deficit/hyperactivity disorder (128), and even anorexia nervosa (129). Behavioral inhibition is a temperamental precursor of social anxiety that can be detected in infancy (130). Longitudinal studies of SA behaviors in infants with behavioral inhibition could be informative about the role of emotional reactivity and regulation in relation to specific SA behaviors. Similarly, studies of different SA behaviors may compare developmental trajectories in infants with (later) autism versus attention-deficit/hyperactivity disorder (16) to understand the role of more general attentional atypicalities. Furthermore, studying SA behaviors in disorders with a known genetic cause can inform us about the underlying biological mechanisms and thus potentially identify stratification markers for ASD. Infants with fragile X syndrome look at other's faces less often than typically developing infants do, mirroring the adult literature from the condition (131) and also seemingly consistent with the high prevalence of ASD in the condition (132). In contrast, although between 70% and 90% of children with Smith-Magenis syndrome have symptoms consistent with an ASD diagnosis (133,134), preliminary evidence suggests that SA is relatively spared in infants with Smith-Magenis syndrome (135) and even enhanced at school age (136). Infants with Williams syndrome (whose personality later in life is often described as hypersocial) have atypically high levels of attention to faces (137).

Ecologically valid studies of SA phenotypes have mostly been conducted in older individuals, but distinct processing during passive social observation versus reciprocal interaction could emerge early (13) (but see candidate 7). As such, and as exemplified by the recent live joint attention eye-tracking study of Nyström *et al.* (49), more work needs to be done to characterize and validate SA behaviors (and potential atypicalities in ASD) in real-life contexts early in life (107). New analytic approaches that take into account the dynamic and bidirectional nature of everyday social stimuli and contexts may reveal more distinct underlying processes than aggregated looking time measures based on passive viewing (Figure 3C, D). Most studies aiming to measure SA have focused on visual attention. This has also been the focus in this review, but it is important that future research expands to include nonvisual modalities. Experiments specifically addressing attention to social auditory cues in ASD early in life are few [but see (138,139)]. To comprehensively address the questions of SA domain specificity and unity from a developmental perspective, it would be important to collect multiple SA candidate measures together with carefully assessed contrast phenotypes longitudinally over time. Furthermore, studies that compare different types of attentional functions (e.g., orienting and disengagement of attention) during processing of social and nonsocial stimuli, beyond examining gaze allocation, are likely to provide more definitive answers that advance the field. It is important to keep in mind that although a phenotype may recruit a domain-specific process at the group level, individual

differences in this phenotype (or case control differences for ASD) may reflect other processes influencing the phenotype in question. In terms of intervention research for ASD, a better understanding of the nature of SA atypicalities is needed to develop better support and intervention programs (140,141). These will necessarily differ depending on whether atypicalities are reflected in social understanding, in social motivation, or in lower-level attention (Figure 2). Relatedly, because gaze differences in ASD may be adaptive responses to other, more fundamental information processing or motivational biases (142), it is not self-evident that it is meaningful, or even ethical, to target such behaviors in intervention trials.

ACKNOWLEDGMENTS AND DISCLOSURES

The work leading to these results was supported by Riksbankens Jubileumsfond (Grant No. P18-0743:1 [to ET]), Knut and Alice Wallenberg Foundation (to TFY), and Innovative Medicines Initiative 2 Joint Undertaking (Grant No. 777394 [to TFY]). This Joint Undertaking receives support from the European Union's Horizon 2020 research and innovation program and EFPIA and AUTISM SPEAKS, Autistica, SFARI.

The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

We thank Kim Astor, Christine Fawcett, and Gustaf Gredebäck for valuable comments on an earlier version of this manuscript.

The authors report no biomedical financial interests or potential conflicts of interest.

ARTICLE INFORMATION

From the Development and Neurodiversity Lab, Department of Psychology, Uppsala University, Uppsala, Sweden (TF-Y, AMP, ET); Center of Neurodevelopmental Disorders, Centre for Psychiatry Research, Department of Women's and Children's Health, Karolinska Institutet & Stockholm Health Care Services, Region, Stockholm, Stockholm, Sweden (TF-Y, AMP); Swedish Collegium for Advanced Study, Uppsala, Sweden (TF-Y); Rare Diseases Research Group, Department of Molecular Medicine and Surgery, Karolinska Institutet, Stockholm, Sweden (JLK); Center for Psychiatry Research, Department of Clinical Neuroscience, Karolinska Institutet & Stockholm Health Care Services, Region Stockholm, Stockholm, Sweden (JLK); Department of Psychology, Stockholm University, Stockholm, Sweden (JLK); and Department of Psychology, Lund University, Lund, Sweden (ET).

Address correspondence to Terje Falck-Ytter, Ph.D., at terje.falck-ytter@psyk.uu.se, or Emilia Thorup, Ph.D., at emilia.thorup@psyk.uu.se.

Received Dec 20, 2021; revised Aug 27, 2022; accepted Sep 27, 2022.

REFERENCES

- Frazier TW, Strauss M, Klingemier EW, Zetzer EE, Hardan AY, Eng C, Youngstrom EA (2017): A meta-analysis of gaze differences to social and nonsocial information between individuals with and without autism. *J Am Acad Child Adolesc Psychiatry* 56:546–555.
- Murias M, Major S, Davlantis K, Franz L, Harris A, Rardin B, *et al.* (2018): Validation of eye-tracking measures of social attention as a potential biomarker for autism clinical trials. *Autism Res* 11:166–174.
- Klin A, Lin DJ, Gorrindo P, Ramsay G, Jones W (2009): Two-year-olds with autism orient to non-social contingencies rather than biological motion. *Nature* 459:257–261.
- Salley B, Colombo J (2016): Conceptualizing social attention in developmental research. *Soc Dev* 25:687–703.
- Nummenmaa L, Calder AJ (2009): Neural mechanisms of social attention. *Trends Cogn Sci* 13:135–143.
- Petersen SE, Posner MI (2012): The attention system of the human brain: 20 years after. *Annu Rev Neurosci* 35:73–89.
- Posner MI (1992): Attention as a cognitive and neural system. *Curr Dir Psychol Sci* 1:11–14.
- Corbetta M, Shulman GL (2002): Control of goal-directed and stimulus-driven attention in the brain. *Nat Rev Neurosci* 3:201–215.
- Amso D, Scerif G (2015): The attentive brain: Insights from developmental cognitive neuroscience. *Nat Rev Neurosci* 16:606–619.
- Posner MI, Rothbart MK, Sheese BE, Voelker P (2012): Control networks and neuromodulators of early development. *Dev Psychol* 48:827–835.
- Mundy P, Bullen J (2022): The bidirectional social-cognitive mechanisms of the social-attention symptoms of autism. *Front Psychiatry* 12:752274.
- Schilbach L, Timmermans B, Reddy V, Costall A, Bente G, Schlicht T (2013): A second-person neuroscience in interaction. *Behav Brain Sci* 36:441–462.
- Redcay E, Schilbach L (2019): Using second-person neuroscience to elucidate the mechanisms of social interaction. *Nat Rev Neurosci* 20:495–505.
- Pfeiffer UJ, Vogeley K, Schilbach L (2013): From gaze cueing to dual eye-tracking: Novel approaches to investigate the neural correlates of gaze in social interaction. *Neurosci Biobehav Rev* 37:2516–2528.
- Valtakari NV, Hooge ITC, Viktorsson C, Nyström P, Falck-Ytter T, Hessel RS (2021): Eye tracking in human interaction: Possibilities and limitations. *Behav Res Methods* 53:1592–1608.
- Braithwaite EK, Gui A, Jones EJM (2020): Social attention: What is it, how can we measure it, and what can it tell us about autism and ADHD? *Prog Brain Res* 254:271–303.
- Mason L, Shic F, Falck-Ytter T, Chakrabarti B, Charman T, Loth E, *et al.* (2021): Preference for biological motion is reduced in ASD: Implications for clinical trials and the search for biomarkers. *Mol Autism* 12:74.
- Falck-Ytter T, Nyström P, Gredebäck G, Gliga T, Bölte S (2018): Reduced orienting to audiovisual synchrony in infancy predicts autism diagnosis at 3 years of age. *J Child Psychol Psychiatry* 59:872–880.
- Shic F, Bradshaw J, Klin A, Scassellati B, Chawarska K (2011): Limited activity monitoring in toddlers with autism spectrum disorder. *Brain Res* 1380:246–254.
- Morton J, Johnson MH (1991): CONSPEC and CONLERN: A two-process theory of infant face recognition. *Psychol Rev* 98:164–181.
- Farroni T, Johnson MH, Menon E, Zulian L, Faraguna D, Csibra G (2005): Newborns' preference for face-relevant stimuli: Effects of contrast polarity. *Proc Natl Acad Sci U S A* 102:17245–17250.
- Shah P, Gaule A, Bird G, Cook R (2013): Robust orienting to proto-facial stimuli in autism. *Curr Biol* 23:R1087–R1088.
- Hirai M, Muramatsu Y, Nakamura M (2020): Developmental changes in orienting towards faces: A behavioral and eye-tracking study. *Int J Behav Dev* 44:157–165.
- Johnson M, Morton J (1991): *Biology and Cognitive Development: The case of Face Recognition*. Oxford: Blackwell Publishing.
- Reynolds GD, Roth KC (2018): The development of attentional biases for faces in infancy: A developmental systems perspective. *Front Psychol* 9:222.
- Johnson MH, Senju A, Tomalski P (2015): The two-process theory of face processing: Modifications based on two decades of data from infants and adults. *Neurosci Biobehav Rev* 50:169–179.
- Simion F, Giordano ED (2015): Face perception and processing in early infancy: Inborn predispositions and developmental changes. *Front Psychol* 6:969.
- Nelson CA (2001): The development and neural bases of face recognition. *Inf Child Develop* 10:3–18.
- Simion F, Macchi Cassia V, Turati C, Valenza E (2001): The origins of face perception: Specific versus non-specific mechanisms. *Inf Child Develop* 10:59–65.
- Wilkinson N, Paikan A, Gredebäck G, Rea F, Metta G (2014): Staring us in the face? An embodied theory of innate face preference. *Dev Sci* 17:809–825.
- Guillon Q, Rogé B, Afzali MH, Baduel S, Kruck J, Hadjikhani N (2016): Intact perception but abnormal orientation towards face-like objects in young children with ASD. *Sci Rep* 6:22119.

32. Turati C (2004): Why faces are not special to newborns: An alternative account of the face preference. *Curr Dir Psychol Sci* 13:5–8.
33. Farroni T, Menon E, Johnson MH (2006): Factors influencing newborns' preference for faces with eye contact. *J Exp Child Psychol* 95:298–308.
34. Reid VM, Dunn K, Young RJ, Amu J, Donovan T, Reissland N (2017): The human fetus preferentially engages with face-like visual stimuli [published correction appears in *Curr Biol* 2017;27:2052] [published correction appears in *Curr Biol* 2018;28:824]. *Curr Biol* 27:1825–e1823.
35. Scheel AM, Ritchie SJ, Brown NJ, Jacques SL (2018): Methodological problems in a study of fetal visual perception. *Curr Biol* 28:R594–R596.
36. Buiatti M, Di Giorgio E, Piazza M, Polloni C, Menna G, Taddei F, *et al.* (2019): Cortical route for facelike pattern processing in human newborns. *Proc Natl Acad Sci U S A* 116:4625–4630.
37. Elsabbagh M, Gliga T, Pickles A, Hudry K, Charman T, Johnson MH, *et al.* (2013): The development of face orienting mechanisms in infants at-risk for autism. *Behav Brain Res* 251:147–154.
38. Di Giorgio E, Turati C, Altoè G, Simion F (2012): Face detection in complex visual displays: An eye-tracking study with 3- and 6-month-old infants and adults. *J Exp Child Psychol* 113:66–77.
39. Gliga T, Elsabbagh M, Andravizou A, Johnson M (2009): Faces attract infants' attention in complex displays. *Infancy* 14:550–562.
40. Frank MC, Vul E, Johnson SP (2009): Development of infants' attention to faces during the first year. *Cognition* 110:160–170.
41. Johnson MH, Kamiloff-Smith A (2004): Perspectives on infant development. *Theor Infant Dev* 121–141.
42. Portugal AM, Viktorsson C, Taylor M, Mason L, Tammimies K, Ronald A, *et al.* (2022): Infants' looking preferences for social versus nonsocial objects reflect genetic variation and are linked to later language development. *OSF Preprints*.
43. Kennedy DP, D'Onofrio BM, Quinn PD, Bölte S, Lichtenstein P, Falck-Ytter T (2017): Genetic influence on eye movements to complex scenes at short timescales. *Curr Biol* 27:3554–3560.e3.
44. Bertenthal BI, Boyer TW (2015): The development of social attention in human infants. In: Puce A, Bertenthal B, editors. *The Many Faces of Social Attention*. Berlin: Springer, 21–65.
45. Webb SJ, Jones EJ, Merkle K, Namkung J, Toth K, Greenson J, *et al.* (2010): Toddlers with elevated autism symptoms show slowed habituation to faces. *Child Neuropsychol* 16:255–278.
46. Hendry A, Jones E, Bedford R, Gliga T, Charman T, Johnson MH, *et al.* (2018): Developmental change in look durations predicts later effortful control in toddlers at familial risk for ASD. *J Neurodev Disord* 10:3.
47. Chawarska K, Macari S, Shic F (2013): Decreased spontaneous attention to social scenes in 6-month-old infants later diagnosed with autism spectrum disorders. *Biol Psychiatry* 74:195–203.
48. Saint-Georges C, Cassel RS, Cohen D, Chetouani M, Laznik M, Maestro S, *et al.* (2010): What studies of family home movies can teach us about autistic infants: A literature review. *Res Autism Spec Disord* 4:355–366.
49. Nyström P, Thorup E, Bölte S, Falck-Ytter T (2019): Joint attention in infancy and the emergence of autism. *Biol Psychiatry* 86:631–638.
50. Oakes LM, Ellis AE (2013): An eye-tracking investigation of developmental changes in infants' exploration of upright and inverted human faces. *Infancy* 18:134–148.
51. Lewkowicz DJ, Hansen-Tift AM (2012): Infants deploy selective attention to the mouth of a talking face when learning speech. *Proc Natl Acad Sci U S A* 109:1431–1436.
52. Viktorsson C, Portugal AM, Li D, Rudling M, Siqueiros Sanchez M, Tammimies K, *et al.* (2023): Preferential looking to eyes versus mouth in early infancy: Heritability and link to concurrent and later development. *J Child Psychol Psychiatry* 64:311–319.
53. Itier RJ, Batty M (2009): Neural bases of eye and gaze processing: The core of social cognition. *Neurosci Biobehav Rev* 33:843–863.
54. Hoehl S (2015): How do neural responses to eyes contribute to face-sensitive ERP components in young infants? A rapid repetition study. *Brain Cogn* 95:1–6.
55. Tenenbaum EJ, Shah RJ, Sobel DM, Malle BF, Morgan JL (2013): Increased focus on the mouth among infants in the first year of life: A longitudinal eye-tracking study. *Infancy* 18:534–553.
56. Hillairet de Boisferon A, Tift AH, Minar NJ, Lewkowicz DJ (2017): Selective attention to a talker's mouth in infancy: Role of audiovisual temporal synchrony and linguistic experience. *Dev Sci* 20:e12381.
57. Constantino JN, Kennon-McGill S, Weichselbaum C, Marrus N, Haider A, Glowinski AL, *et al.* (2017): Infant viewing of social scenes is under genetic control and is atypical in autism. *Nature* 547:340–344.
58. Jones W, Klin A (2013): Attention to eyes is present but in decline in 2–6-month-old infants later diagnosed with autism. *Nature* 504:427–431.
59. Chita-Tegmark M (2016): Social attention in ASD: A review and meta-analysis of eye-tracking studies. *Res Dev Disabil* 48:79–93.
60. Falck-Ytter T, Carlström C, Johansson M (2015): Eye contact modulates cognitive processing differently in children with autism. *Child Dev* 86:37–47.
61. Bruner JS (1974): From communication to language—A psychological perspective. *Cognition* 3:255–287.
62. Scaife M, Bruner JS (1975): The capacity for joint visual attention in the infant. *Nature* 253:265–266.
63. Friesen CK, Kingstone A (1998): The eyes have it! Reflexive orienting is triggered by nonpredictive gaze. *Psychon Bull Rev* 5:490–495.
64. Ristic J, Friesen CK, Kingstone A (2002): Are eyes special? It depends on how you look at it. *Psychon Bull Rev* 9:507–513.
65. Farroni T, Johnson MH, Brockbank M, Simion F (2000): Infants' use of gaze direction to cue attention: The importance of perceived motion. *Vis Cogn* 7:705–718.
66. Farroni T, Massaccesi S, Pividori D, Johnson MH (2004): Gaze following in newborns. *Infancy* 5:39–60.
67. Calder AJ, Beaver JD, Winston JS, Dolan RJ, Jenkins R, Eger E, *et al.* (2007): Separate coding of different gaze directions in the superior temporal sulcus and inferior parietal lobule. *Curr Biol* 17:20–25.
68. Tipples J (2002): Eye gaze is not unique: Automatic orienting in response to uninformative arrows. *Psychon Bull Rev* 9:314–318.
69. Friesen CK, Ristic J, Kingstone A (2004): Attentional effects of counterpredictive gaze and arrow cues. *J Exp Psychol Hum Percept Perform* 30:319–329.
70. Langdon R, Smith P (2005): Spatial cueing by social versus nonsocial directional signals. *Vis Cogn* 12:1497–1527.
71. Ristic J, Wright A, Kingstone A (2007): Attentional control and reflexive orienting to gaze and arrow cues. *Psychon Bull Rev* 14:964–969.
72. Hietanen JK, Leppänen JM, Peltola MJ, Linna-Aho K, Ruuhiala HJ (2008): Seeing direct and averted gaze activates the approach-avoidance motivational brain systems. *Neuropsychologia* 46:2423–2430.
73. Kylläinen A, Hietanen JK (2004): Attention orienting by another's gaze direction in children with autism. *J Child Psychol Psychiatry* 45:435–444.
74. Rombough A, Iarocci G (2013): Orienting in response to gaze and the social use of gaze among children with autism spectrum disorder. *J Autism Dev Disord* 43:1584–1596.
75. Gillespie-Lynch K, Elias R, Escudero P, Hutman T, Johnson SP (2013): Atypical gaze following in autism: A comparison of three potential mechanisms. *J Autism Dev Disord* 43:2779–2792.
76. Goldberg MC, Mostow AJ, Vecera SP, Larson JCG, Mostofsky SH, Mahone EM, *et al.* (2008): Evidence for impairments in using static line drawings of eye gaze cues to orient visual-spatial attention in children with high functioning autism. *J Autism Dev Disord* 38:1405–1413.
77. Pruett JR, LaMacchia A, Hoertel S, Squire E, McVey K, Todd RD, *et al.* (2011): Social and non-social cueing of visuospatial attention in autism and typical development. *J Autism Dev Disord* 41:715–731.
78. Swettenham J, Condie S, Campbell R, Milne E, Coleman M (2003): Does the perception of moving eyes trigger reflexive visual orienting in autism? *Philos Trans R Soc Lond B Biol Sci* 358:325–334.

79. Johnson MH, Griffin R, Csibra G, Halit H, Farroni T, De Haan M, *et al.* (2005): The emergence of the social brain network: Evidence from typical and atypical development. *Dev Psychopathol* 17:599–619.
80. Senju A, Tojo Y, Dairoku H, Hasegawa T (2004): Reflexive orienting in response to eye gaze and an arrow in children with and without autism. *J Child Psychol Psychiatry* 45:445–458.
81. Chawarska K, Klin A, Volkmar F (2003): Automatic attention cueing through eye movement in 2-year-old children with autism. *Child Dev* 74:1108–1122.
82. Nation K, Penny S (2008): Sensitivity to eye gaze in autism: Is it normal? Is it automatic? Is it social? *Dev Psychopathol* 20:79–97.
83. Greene DJ, Colich N, Iacoboni M, Zaidel E, Bookheimer SY, Dapretto M (2011): Atypical neural networks for social orienting in autism spectrum disorders. *Neuroimage* 56:354–362.
84. D'Entremont B, Hains SMJ, Muir DW (1997): A demonstration of gaze following in 3- to 6-month-olds. *Infant Behav Dev* 20:569–572.
85. Klin A, Shultz S, Jones W (2015): Social visual engagement in infants and toddlers with autism: Early developmental transitions and a model of pathogenesis. *Neurosci Biobehav Rev* 50:189–203.
86. Gredebäck G, Fikke L, Melinder A (2010): The development of joint visual attention: A longitudinal study of gaze following during interactions with mothers and strangers. *Dev Sci* 13:839–848.
87. Baldwin DA (1991): Infants contribution to the achievement of joint reference. *Child Dev* 62:875–890.
88. Beuker KT, Rommelse NN, Donders R, Buitelaar JK (2013): Development of early communication skills in the first two years of life. *Infant Behav Dev* 36:71–83.
89. Vaughan Van Hecke AV, Mundy P, Block JJ, Delgado CE, Parlade MV, Pomares YB, *et al.* (2012): Infant responding to joint attention, executive processes, and self-regulation in preschool children. *Infant Behav Dev* 35:303–311.
90. Caruana N, Brock J, Woolgar A (2015): A frontotemporoparietal network common to initiating and responding to joint attention bids. *Neuroimage* 108:34–46.
91. Redcay E, Kleiner M, Saxe R (2012): Look at this: The neural correlates of initiating and responding to bids for joint attention. *Front Hum Neurosci* 6:169.
92. Schilbach L, Wilms M, Eickhoff SB, Romanzetti S, Tepest R, Bente G, *et al.* (2010): Minds made for sharing: Initiating joint attention recruits reward-related neurocircuitry. *J Cogn Neurosci* 22:2702–2715.
93. Elison JT, Wolff JJ, Heimer DC, Paterson SJ, Gu H, Hazlett HC, *et al.* (2013): Frontolimbic neural circuitry at 6 months predicts individual differences in joint attention at 9 months. *Dev Sci* 16:186–197.
94. Grossmann T, Johnson MH (2010): Selective prefrontal cortex responses to joint attention in early infancy. *Biol Lett* 6:540–543.
95. Deák GO (2015): When and where do infants follow gaze? Joint IEEE international conference on development and learning and epigenetic robotics (ICDL-EpiRob). IEEE Publications, 182–187.
96. Deligianni F, Senju A, Gergely G, Csibra G (2011): Automated gaze-contingent objects elicit orientation following in 8-month-old infants. *Dev Psychol* 47:1499–1503.
97. Astor K, Thiele M, Gredebäck G (2021): Gaze following emergence relies on both perceptual cues and social awareness. *Cogn Dev* 60:101121.
98. Chiang CH, Soong WT, Lin TL, Rogers SJ (2008): Nonverbal communication skills in young children with autism. *J Autism Dev Disord* 38:1898–1906.
99. Dawson G, Toth K, Abbott R, Osterling J, Munson J, Estes A, *et al.* (2004): Early social attention impairments in autism: Social orienting, joint attention, and attention to distress. *Dev Psychol* 40:271–283.
100. Landa RJ, Holman KC, Garrett-Mayer E (2007): Social and communication development in toddlers with early and later diagnosis of autism spectrum disorders. *Arch Gen Psychiatry* 64:853–864.
101. Rozga A, Hutman T, Young GS, Rogers SJ, Ozonoff S, Dapretto M, *et al.* (2011): Behavioral profiles of affected and unaffected siblings of children with autism: Contribution of measures of mother–infant interaction and nonverbal communication. *J Autism Dev Disord* 41:287–301.
102. Akechi H, Senju A, Kikuchi Y, Tojo Y, Osanai H, Hasegawa T (2011): Do children with ASD use referential gaze to learn the name of an object? An eye-tracking study. *Res Autism Spec Disord* 5:1230–1242.
103. Falck-Ytter T, Thorup E, Bölte S (2015): Brief report: Lack of processing bias for the objects other people attend to in 3-year-olds with autism. *J Autism Dev Disord* 45:1897–1904.
104. Bedford R, Elsabbagh M, Gliga T, Pickles A, Senju A, Charman T, *et al.* (2012): Precursors to social and communication difficulties in infants at-risk for autism: Gaze following and attentional engagement. *J Autism Dev Disord* 42:2208–2218.
105. Parsons JP, Bedford R, Jones EJJ, Charman T, Johnson MH, Gliga T (2019): Gaze following and attention to objects in infants at familial risk for ASD. *Front Psychol* 10:1799.
106. Stallworthy IC, Lasch C, Berry D, Wolff JJ, Pruett JR Jr, Marrus N, *et al.* (2022): Variability in responding to joint attention cues in the first year is associated with autism outcome. *J Am Acad Child Adolesc Psychiatry* 61:413–422.
107. Nasiopoulos E, Risko EF, Kingstone A (2015): Social attention, social presence, and the dual function of gaze. In: Puce A, Bertenthal B, editors. *The Many Faces of Social Attention*. Berlin: Springer, 129–155.
108. Carpenter M, Nagell K, Tomasello M (1998): Social cognition, joint attention, and communicative competence from 9 to 15 months of age. *Monogr Soc Res Child Dev* 63:i-vi, 1–143.
109. Ibañez LV, Grantz CJ, Messinger DS (2013): The development of referential communication and autism symptomatology in high-risk infants. *Infancy* 18:1–18.
110. Gotham K, Risi S, Pickles A, Lord C (2007): The autism diagnostic observation schedule: Revised algorithms for improved diagnostic validity. *J Autism Dev Disord* 37:613–627.
111. Falck-Ytter T, Gredebäck G, von Hofsten C (2006): Infants predict other people's action goals. *Nat Neurosci* 9:878–879.
112. Rosander K, von Hofsten C (2011): Predictive gaze shifts elicited during observed and performed actions in 10-month-old infants and adults. *Neuropsychologia* 49:2911–2917.
113. Elsner C, D'Ausilio A, Gredebäck G, Falck-Ytter T, Fadiga L (2013): The motor cortex is causally related to predictive eye movements during action observation. *Neuropsychologia* 51:488–492.
114. Flanagan JR, Johansson RS (2003): Action plans used in action observation. *Nature* 424:769–771.
115. Csibra G (2007): Action mirroring and action interpretation: An alternative account. In: Haggard P, Rosetti Y, Kawato M, editors. *Sensorimotor Foundations of Higher Cognition Attention and Performance XXII*. Oxford: Oxford University Press, 435–459.
116. Gredebäck G, Falck-Ytter T (2015): Eye movements during action observation. *Perspect Psychol Sci* 10:591–598.
117. Iacoboni M, Dapretto M (2006): The mirror neuron system and the consequences of its dysfunction. *Nat Rev Neurosci* 7:942–951.
118. Van de Cruys S, Evers K, Van der Hallen R, Van Eylen L, Boets B, de Wit L, *et al.* (2014): Precise minds in uncertain worlds: Predictive coding in autism. *Psychol Rev* 121:649–675.
119. Falck-Ytter T (2010): Young children with autism spectrum disorder use predictive eye movements in action observation. *Biol Lett* 6:375–378.
120. Braukmann R, Ward E, Hessels RS, Bekkering H, Buitelaar JK, Hunnius S (2018): Action prediction in 10-month-old infants at high and low familial risk for autism spectrum disorder. *Res Autism Spec Disord* 49:34–46.
121. Krogh-Jespersen S, Kaldy Z, Valadez AG, Carter AS, Woodward AL (2018): Goal prediction in 2-year-old children with and without autism spectrum disorder: An eye-tracking study. *Autism Res* 11:870–882.
122. Itti L, Koch C (2001): Computational modelling of visual attention. *Nat Rev Neurosci* 2:194–203.
123. Birmingham E, Bischof WF, Kingstone A (2009): Saliency does not account for fixations to eyes within social scenes. *Vision Res* 49:2992–3000.
124. Macari S, Milgramm A, Reed J, Shic F, Powell KK, Macris D, *et al.* (2021): Context-specific dyadic attention vulnerabilities during the first year in infants later developing autism spectrum disorder. *J Am Acad Child Adolesc Psychiatry* 60:166–175.
125. Chevallier C, Kohls G, Troiani V, Brodtkin ES, Schultz RT (2012): The social motivation theory of autism. *Trends Cogn Sci* 16:231–239.

126. Chen J, van den Bos E, Westenberg PM (2020): A systematic review of visual avoidance of faces in socially anxious individuals: Influence of severity, type of social situation, and development. *J Anxiety Disord* 70:102193.
127. Kleberg JL, Högström J, Sundström K, Frick A, Serlachius E (2021): Delayed gaze shifts away from others' eyes in children and adolescents with social anxiety disorder. *J Affect Disord* 278:280–287.
128. Ioannou C, Seemani D, Stefanou ME, Riedel A, Tebart van Elst L, Smyrnis N, *et al.* (2020): Comorbidity matters: Social visual attention in a comparative study of autism spectrum disorder, attention-deficit/hyperactivity disorder and their comorbidity. *Front Psychiatry* 11:545567.
129. Kerr-Gaffney J, Jones E, Mason L, Hayward H, Murphy D, Loth E, Tchanturia K (2022): Social attention in anorexia nervosa and autism spectrum disorder: Role of social motivation. *Autism* 26:1641–1655.
130. Tang A, Crawford H, Morales S, Degnan KA, Pine DS, Fox NA (2020): Infant behavioral inhibition predicts personality and social outcomes three decades later. *Proc Natl Acad Sci U S A* 117:9800–9807.
131. Black CJ, Hogan AL, Smith KD, Roberts JE (2021): Early behavioral and physiological markers of social anxiety in infants with fragile X syndrome. *J Neurodev Disord* 13:11.
132. McDuffie A, Thurman AJ, Hagerman RJ, Abbeduto L (2015): Symptoms of autism in males with fragile X syndrome: A comparison to nonsyndromic ASD using current ADI-R scores. *J Autism Dev Disord* 45:1925–1937.
133. Laje G, Morse R, Richter W, Ball J, Pao M, Smith AC (2010): Autism spectrum features in Smith–Magenis syndrome. *Am J Med Genet C Semin Med Genet* 154C:456–462.
134. Nag HE, Nordgren A, Anderlid BM, Nærland T (2018): Reversed gender ratio of autism spectrum disorder in Smith–Magenis syndrome. *Mol Autism* 9:1.
135. Wolters PL, Gropman AL, Martin SC, Smith MR, Hildenbrand HL, Brewer CC, *et al.* (2009): Neurodevelopment of children under 3 years of age with Smith–Magenis syndrome. *Pediatr Neurol* 41:250–258.
136. Wilde L, Silva D, Oliver C (2013): The nature of social preference and interactions in Smith–Magenis syndrome. *Res Dev Disabil* 34:4355–4365.
137. Mervis CB, Morris CA, Klein-Tasman BP, Bertrand J, Kwitny S, Appelbaum LG, *et al.* (2003): Attentional characteristics of infants and toddlers with Williams syndrome during triadic interactions. *Dev Neuropsychol* 23:243–268.
138. Čeponienė R, Lepistö T, Shestakova A, Vanhala R, Alku P, Nääätänen R, *et al.* (2003): Speech–sound-selective auditory impairment in children with autism: They can perceive but do not attend. *Proc Natl Acad Sci U S A* 100:5567–5572.
139. Rudling M, Nystrom P, Bolte S, Falck-Ytter T (2022): Larger pupil dilation to nonsocial sounds in infants with subsequent autism diagnosis. *J Child Psychol Psychiatry* 63:793–801.
140. Green J, Charman T, Pickles A, Wan MW, Elsabbagh M, Slonims V, *et al.* (2015): Parent-mediated intervention versus no intervention for infants at high risk of autism: A parallel, single-blind, randomised trial. *Lancet Psychiatry* 2:133–140.
141. Walsh P, Elsabbagh M, Bolton P, Singh I (2011): In search of biomarkers for autism: Scientific, social and ethical challenges. *Nat Rev Neurosci* 12:603–612.
142. Johnson MH, Charman T, Pickles A, Jones EJJ (2021): Annual Research Review: Anterior Modifiers in the Emergence of Neurodevelopmental Disorders (AMEND)—A systems neuroscience approach to common developmental disorders. *J Child Psychol Psychiatry* 62:610–630.
143. Falck-Ytter T, von Hofsten C, Gillberg C, Fernell E (2013): Visualization and analysis of eye movement data from children with typical and atypical development. *J Autism Dev Disord* 43:2249–2258.