



PAPER • OPEN ACCESS

Tunable phonon-driven magnon–magnon entanglement at room temperature

To cite this article: Yuefei Liu *et al* 2023 *New J. Phys.* **25** 113032

View the [article online](#) for updates and enhancements.

You may also like

- [Magnon-magnon entanglement generation between two remote interaction-free optomagnonic systems via optical Bell-state measurement](#)
S Golkar, E Ghasemian, M Setodeh Kheirabady et al.
- [Modulation-induced long-range magnon bound states in one-dimensional optical lattices](#)
Wenjie Liu, Yongguan Ke, Bo Zhu et al.
- [Enhanced the Bi-\(Tri-\)partite entanglement between three magnons in a cavity magnomechanics system](#)
Abdelkader Hidki, Ya-long Ren, Abderrahim Lakhfif et al.



PAPER

OPEN ACCESS

RECEIVED

5 September 2023

REVISED

29 October 2023

ACCEPTED FOR PUBLICATION

9 November 2023

PUBLISHED

20 November 2023

Original Content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.



Tunable phonon-driven magnon–magnon entanglement at room temperature

Yuefei Liu^{1,*}, Anders Bergman², Andrey Bagrov³, Anna Delin^{1,4,7}, Danny Thonig^{2,5}, Manuel Pereiro² ,
Olle Eriksson^{2,8}, Simon Streib², Erik Sjöqvist^{2,*} and Vahid Azimi-Mousolou^{6,2,*} 

¹ Department of Applied Physics, School of Engineering Sciences, KTH Royal Institute of Technology, AlbaNova University Center, SE-10691 Stockholm, Sweden

² Department of Physics and Astronomy, Uppsala University, Box 516, SE-751 20 Uppsala, Sweden

³ Institute for Molecules and Materials, Radboud University, Heyendaalseweg 135, 6525AJ Nijmegen, The Netherlands

⁴ Swedish e-Science Research Center (SeRC), KTH Royal Institute of Technology, SE-10044 Stockholm, Sweden

⁵ School of Science and Technology, Örebro University, SE-701 82 Örebro, Sweden

⁶ Department of Applied Mathematics and Computer Science, Faculty of Mathematics and Statistics, University of Isfahan, Isfahan 81746-73441, Iran

⁷ Wallenberg Initiative Materials Science for Sustainability (WISE), KTH Royal Institute of Technology, SE-10044 Stockholm, Sweden

⁸ Wallenberg Initiative Materials Science for Sustainability (WISE), Uppsala University, 75121 Uppsala, Sweden

* Authors to whom any correspondence should be addressed.

E-mail: yuefei@kth.se, erik.sjoqvist@physics.uu.se and v.azimi@sci.ui.ac.ir

Keywords: magnon, phonon, steady-state, quantum entanglement, room temperature

Abstract

We report the existence of entangled steady-states in bipartite quantum magnonic systems at elevated temperatures. We consider dissipative dynamics of two magnon modes in a bipartite antiferromagnet, subjected to interaction with a phonon mode and an external rotating magnetic field. To quantify the bipartite magnon–magnon entanglement, we use entanglement negativity and compute its dependence on temperature and magnetic field. We provide evidence that the coupling between magnon and phonon modes is necessary for the entanglement, and that, for any given phonon frequency and magnon–phonon coupling rate, there are always ranges of the magnetic field amplitudes and frequencies for which magnon–magnon entanglement persists at room temperature.

1. Introduction

Entanglement is a central concept of quantum physics that has developed into a key resource in quantum technology [1]. It plays a role in quantum information processing, opens a way towards intercept-resilient quantum communications, and enables increased sensitivity in quantum metrology. If it were possible to overcome the thermal noise and prepare robust entangled quantum states at ambient conditions, the corresponding implications for the future quantum devices would be significant [2–9].

Magnons exhibit quantum properties over a wide range of temperatures and offer a possibility to access quantum phenomena at room temperature [10, 11]. Compatibility of magnonics with hybrid quantum platforms [11–15] makes magnons potentially useful for quantum data processing. Most recently, particular attention has been paid to antiferromagnetic (AFM) quantum magnonics due to its large equilibrium squeezing and entanglement [11, 16–26].

Here, we report evidence for equilibrium phonon-driven magnon–magnon entanglement in bipartite AFM materials. We compute entanglement negativity [27] of a pair of magnon modes coupled to a lattice vibration, in the presence of an off-resonant external rotating magnetic field. Our aim is to study the stability of bipartite entanglement between magnons to thermal noise. We search for the regime where the magnonic degrees of freedom remain entangled with each other rather than with the environment, and hence can be used as coherent channels in quantum information protocols. We obtain a steady-state entanglement diagram in the temperature and magnetic field plane, which shows the possibility of having magnon–magnon entanglement at room temperature. We analyze the magnon entanglement stability against

dissipation caused by magnon–phonon coupling and observe that the two-mode magnon system indeed permits high-temperature entanglement at any value of the magnetic field provided that the magnetic field frequency is adjusted to the proper range. In turn, the existence of crystalline and synthetic AFM materials, such as NiO [28], 2D Ising systems like MnPSe₃ [29, 30], YIG-based synthetic AFMs [31], and perovskite manganites [32, 33], particularly SrTcO₃ [34], CaTcO₃ [35], BiFeO₃ [36] that have high Neel temperatures, provide a space for further study of high-temperature entanglement in quantum magnonics.

2. Physical system

To pursue our analysis, we assume a pair of magnon modes in a bipartite AFM material subjected to a vibrational mode in an off-resonant rotating magnetic field as shown in figure 1. The system Hamiltonian reads

$$\hat{H} = \hat{H}_m + \hat{H}_{m-m} + \hat{H}_z + \hat{H}_{ph} + \hat{H}_{m-ph} \quad (1)$$

with each term defined as follows. \hat{H}_m and \hat{H}_{m-m} correspond to single- and two-mode magnon Hamiltonian, respectively, at a given \mathbf{k} -vector. To find the explicit form of these terms, we consider an AFM spin Hamiltonian, $\hat{H}_s = \sum_{\langle i,j \rangle} \hat{\mathbf{S}}_i \mathbb{I}_{ij} \hat{\mathbf{S}}_j$ with nearest neighbor interactions, where $\hat{\mathbf{S}}_i$ is the spin vector operator at lattice site i , and \mathbb{I}_{ij} is the bilinear interaction tensor matrix between sites i and j , which involves all different types of interactions, e.g. Heisenberg interaction, Dzyaloshinskii–Moriya interaction, anisotropy [23–26]. By using the linearized Holstein–Primakoff transformation

$$\begin{aligned} \text{Sublattice A : } & \begin{cases} \hat{S}_i^z = S - \hat{a}_i^\dagger \hat{a}_i, \\ \hat{S}_i^+ = \hat{S}_i^x + i\hat{S}_i^y \approx \sqrt{2S} \hat{a}_i, \end{cases} \\ \text{Sublattice B : } & \begin{cases} \hat{S}_j^z = \hat{b}_j^\dagger \hat{b}_j - S, \\ \hat{S}_j^+ = \hat{S}_j^x + i\hat{S}_j^y \approx \sqrt{2S} \hat{b}_j^\dagger, \end{cases} \end{aligned}$$

and Fourier transformation [11], one obtains $\hat{H}_s = \hat{H}_m + \hat{H}_{m-m}$ with

$$\begin{aligned} \hat{H}_m &= \hbar \omega_{a_k} \hat{a}_k^\dagger \hat{a}_k + \hbar \omega_{b_{-k}} \hat{b}_{-k}^\dagger \hat{b}_{-k} \\ \hat{H}_{m-m} &= \hbar g_{a_k b_{-k}}^* \hat{a}_k^\dagger \hat{b}_{-k}^\dagger + \hbar g_{a_k b_{-k}} \hat{a}_k \hat{b}_{-k}, \end{aligned} \quad (2)$$

for a given momentum \mathbf{k} -vector. Here, the bosonic annihilation operators \hat{a}_k and \hat{b}_{-k} represent two counter-propagating magnon modes that describe identifiable magnon modes associated with each sublattice of the AFM.

We further assume that the prepared magnon system is subjected to an off-resonant rotating magnetic field with frequency $\omega_0 < \omega_{a_k}, \omega_{b_{-k}}$. An in-plane rotating magnetic field

$$\mathbf{B} = B_0 (\cos(\phi_0 + \omega_0 t), \sin(\phi_0 + \omega_0 t), 0) \quad (3)$$

interacts with a spin system through the Zeeman term $\hat{H}_z = \sum_i \mu \mathbf{B} \cdot \hat{\mathbf{S}}_i$. By using the same transformations as for \hat{H}_s , we obtain

$$\begin{aligned} \hat{H}_z &= \hbar \left[\frac{\mu B_0 \Gamma_{\mathbf{k}}}{\hbar} \right] \sqrt{\frac{SN}{2}} \left(e^{-i(\phi_0 + \zeta_{\mathbf{k}} + \omega_0 t)} \hat{a}_{\mathbf{k}} + e^{i(\phi_0 + \zeta_{\mathbf{k}} + \omega_0 t)} \hat{a}_{\mathbf{k}}^\dagger \right) \\ &\quad + \hbar \left[\frac{\mu B_0 \Gamma_{\mathbf{k}}}{\hbar} \right] \sqrt{\frac{SN}{2}} \left(e^{i(\phi_0 + \zeta_{\mathbf{k}} + \omega_0 t)} \hat{b}_{-\mathbf{k}} + e^{-i(\phi_0 + \zeta_{\mathbf{k}} + \omega_0 t)} \hat{b}_{-\mathbf{k}}^\dagger \right), \end{aligned} \quad (4)$$

for a given \mathbf{k} -vector. B_0 is the strength of the magnetic field, N is the number of spins in the system, S is the total spin per site, and $\Gamma_{\mathbf{k}} e^{-i\zeta_{\mathbf{k}}} = \frac{2}{N} \sum_{i \in A(B)} e^{-i\mathbf{k} \cdot \mathbf{r}_i}$ with \mathbf{r}_i being the position vector of site i in the corresponding sublattice. $\Gamma_{\mathbf{k}}$ and $\zeta_{\mathbf{k}}$ come about by Fourier transforming the Hamiltonian from real space to \mathbf{k} -space. It is assumed that the values of $\Gamma_{\mathbf{k}}$ and $\zeta_{\mathbf{k}}$ for sublattice A are identical to those for sublattice B .

For the magnon–phonon interaction, we consider magnetoelastic coupling, where magnetic degrees of freedom interact with elastic displacements of atoms from their equilibrium positions. Assuming single phonon process [37] with translational and rotational symmetries, the quantized phonon and magnetoelastic spin-phonon coupling terms of the Hamiltonian read [28]




Figure 1. Schematic illustration of bipartite collective spin waves in an antiferromagnetic material subjected to a vibrational phonon mode in an off-resonant rotating magnetic field. The rotating magnetic field is illustrated in the lower left corner while the coupling between magnons in sublattice A and B to the phonons are illustrated in the lower right part of the figure. The upper figure depicts schematically the spins interacting with the lattice.

$$\begin{aligned}\hat{H}'_{\text{ph}} &= \hbar \sum_{\mathbf{k}'} \omega_{\mathbf{k}'} \hat{c}_{\mathbf{k}'}^\dagger \hat{c}_{\mathbf{k}'}, \\ \hat{H}'_{\text{m-ph}} &= \hbar \sum_{\mathbf{k}'} \sum_i \sum_{\delta} \sum_{\alpha, \beta} G_{\mathbf{k}'\delta}^{\alpha\beta} \mathbf{S}_i^\alpha \mathbf{S}_{i+\delta}^\beta \left(\hat{c}_{\mathbf{k}'}^\dagger + \hat{c}_{-\mathbf{k}'} \right) e^{i\mathbf{k}' \cdot \mathbf{r}_i},\end{aligned}\quad (5)$$

where i is summed over all magnetic lattice sites and δ denotes a vector pointing from lattice site i to a neighboring site. $G_{\mathbf{k}'\delta}^{\alpha\beta}$ is the coupling strength between the spins in spatial directions $\alpha, \beta \in \{x, y, z\}$ and the elastic displacement mode at a given \mathbf{k}' . The operators $\hat{c}_{\mathbf{k}'}^\dagger$ ($\hat{c}_{\mathbf{k}'}$) represent annihilation (creation) of an elastic phonon mode with frequency $\omega_{\mathbf{k}'}$, and crystal momentum \mathbf{k}' . Taking into account the magnetoelastic coupling only in the transverse direction, the interacting spin-phonon coupling term of the Hamiltonian reduces to

$$\hat{H}'_{\text{m-ph}} = \hbar \sum_{\mathbf{k}'} \sum_{i, \delta} G_{\mathbf{k}'\delta}^{zz} \mathbf{S}_i^z \mathbf{S}_{i+\delta}^z \left(\hat{c}_{\mathbf{k}'}^\dagger + \hat{c}_{-\mathbf{k}'} \right) e^{i\mathbf{k}' \cdot \mathbf{r}_i}. \quad (6)$$

By applying the same transformations to the spin degrees of freedom as for the other terms in the Hamiltonian above, the resulting magnetoelastic Hamiltonian takes the form

$$\hat{H}'_{\text{m-ph}} = \hbar \sum_{\mathbf{k}, \mathbf{k}'} \left[g_{ac}^{(\mathbf{k}, \mathbf{k}')} \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}+\mathbf{k}'} + g_{bc}^{(-\mathbf{k}, -\mathbf{k}')} \hat{b}_{-\mathbf{k}}^\dagger \hat{b}_{-\mathbf{k}-\mathbf{k}'} \right] \left(\frac{\hat{c}_{\mathbf{k}'}^\dagger + \hat{c}_{-\mathbf{k}'}}{\sqrt{2}} \right), \quad (7)$$

where a term corresponding to a shift of each phonon mode is neglected. This type of magnon–phonon interaction naturally describes the scattering of magnons with emission or absorption of phonons based on momentum conservation [15, 38–41]. However, if a given pair of magnon modes $\hat{a}_{\mathbf{k}}$ and $\hat{b}_{-\mathbf{k}}$ are initially prepared in an excited state having large occupation number, the subsequent evolution will be dominated by interactions involving $\hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}}$ and $\hat{b}_{-\mathbf{k}}^\dagger \hat{b}_{-\mathbf{k}}$, thereby forcing the phonon they couple with to have zero momentum, i.e. $\mathbf{k}' = 0$. In the light of the dominant term, the effective magnon–phonon interaction Hamiltonian takes the form

$$\hat{H}_{\text{m-ph}} = \hbar \left[g_{ac}^{(\mathbf{k}, 0)} \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}} + g_{bc}^{(-\mathbf{k}, 0)} \hat{b}_{-\mathbf{k}}^\dagger \hat{b}_{-\mathbf{k}} \right] \left(\frac{\hat{c}_0^\dagger + \hat{c}_0}{\sqrt{2}} \right). \quad (8)$$

This implies that the phonon energy in the model system can be written as

$$\hat{H}_{\text{ph}} = \hbar \omega_{c_0} \hat{c}_0^\dagger \hat{c}_0, \quad (9)$$

which physically corresponds to zone-center optical phonon modes. Note that phonon contributions from non-zero \mathbf{k}' are included implicitly in the dynamics of the system as dissipation and noise terms in the

quantum Langevin equations (QLEs) below. It is worth noting that the type of bosonic Hamiltonian expressed in equation (8) is commonly used in optomechanical systems [1, 42].

By collecting the above terms for a system with magnons of a specific value of \mathbf{k} being prepared, the relevant Hamiltonian in the rotating frame, given by applying $\hat{U}(t) = e^{i\omega_0 t[\hat{b}_{-\mathbf{k}}^\dagger \hat{b}_{-\mathbf{k}} - \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}}]}$, takes the following form

$$\begin{aligned}\hat{H} &= i\hbar \frac{d\hat{U}}{dt} \hat{U}^\dagger + \hat{U} \hat{H} \hat{U}^\dagger \\ &= \hbar(\omega_{a_{\mathbf{k}}} + \omega_0) \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}} + \hbar(\omega_{b_{-\mathbf{k}}} - \omega_0) \hat{b}_{-\mathbf{k}}^\dagger \hat{b}_{-\mathbf{k}} + \hbar g_{a_{\mathbf{k}}b_{-\mathbf{k}}}^* \hat{a}_{\mathbf{k}}^\dagger \hat{b}_{-\mathbf{k}}^\dagger + \hbar g_{a_{\mathbf{k}}b_{-\mathbf{k}}} \hat{a}_{\mathbf{k}} \hat{b}_{-\mathbf{k}} \\ &\quad + \hbar\Omega \left(e^{-i(\phi_0 + \zeta_{\mathbf{k}})} \hat{a}_{\mathbf{k}} + e^{i(\phi_0 + \zeta_{\mathbf{k}})} \hat{a}_{\mathbf{k}}^\dagger \right) + \hbar\Omega \left(e^{i(\phi_0 + \zeta_{\mathbf{k}})} \hat{b}_{-\mathbf{k}} + e^{-i(\phi_0 + \zeta_{\mathbf{k}})} \hat{b}_{-\mathbf{k}}^\dagger \right) \\ &\quad + \hbar\omega_{c_0} \hat{c}_0^\dagger \hat{c}_0 + \hbar \left[g_{a_{\mathbf{k}}c_0} \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}} + g_{b_{-\mathbf{k}}c_0} \hat{b}_{-\mathbf{k}}^\dagger \hat{b}_{-\mathbf{k}} \right] \left(\frac{\hat{c}_0^\dagger + \hat{c}_0}{\sqrt{2}} \right)\end{aligned}\quad (10)$$

with $\Omega = \left[\frac{\mu B_0 \Gamma_{\mathbf{k}}}{\hbar} \right] \sqrt{\frac{SN}{2}}$. Note that the parameters Ω and $\zeta_{\mathbf{k}}$ can be tuned by the strength B_0 and azimuthal angle ϕ_0 of the off-resonant rotating magnetic field. Thus, without loss of generality, we assume $\phi_0 + \zeta_{\mathbf{k}} \equiv 0$.

To summarize, the model Hamiltonian takes the form

$$\begin{aligned}\hat{H} &= \hbar\omega_a \hat{a}^\dagger \hat{a} + \hbar\omega_b \hat{b}^\dagger \hat{b} + \hbar\omega_c \hat{c}^\dagger \hat{c} + \hbar\Omega (\hat{a} + \hat{a}^\dagger) + \hbar\Omega (\hat{b} + \hat{b}^\dagger) \\ &\quad + \hbar g_{ab}^* \hat{a}^\dagger \hat{b}^\dagger + \hbar g_{ab} \hat{a} \hat{b} + \hbar \left[g_{ac} \hat{a}^\dagger \hat{a} + g_{bc} \hat{b}^\dagger \hat{b} \right] \left(\frac{\hat{c} + \hat{c}^\dagger}{\sqrt{2}} \right),\end{aligned}\quad (11)$$

where we drop the subscripts $\mathbf{k}, -\mathbf{k}, 0$ for simplicity. We assume that the magnon modes in the AFM have the same frequency ω that are split by an off-resonant external rotating magnetic field with frequency ω_0 . Thus, we have the detuned frequencies $\omega_a = \omega + \omega_0$ and $\omega_b = \omega - \omega_0$ of the two magnon modes. The magnon–magnon coupling with strength g_{ab} is known to give rise to magnon squeezing and entanglement in each magnon eigenstate in AFMs [23–26].

Physical systems that are relevant for this investigation should have zone center (optical) phonons that couple to magnons with finite crystal momentum (up to the zone boundary). Typically this happens when the frequencies are of similar magnitude. There are many reports in the literature on such materials [28, 30, 38]. Thus, we pay a particular attention to the frequency ratio $\omega_c/\omega = 1$, although, as figures 4 and 5 below show, our results are valid for a wider range of ω_c/ω as well.

3. Dynamics of the system

We consider the system under dissipation caused by thermal fluctuations, uncontrolled coupling to other modes (e.g. phonons with non-zero momentum \mathbf{k}'), and Brownian motion. Such dissipative dynamics can be described by nonlinear QLEs [43].

To specify the dynamics of the system, it is handy to introduce dimensionless quadratures

$$\begin{aligned}\hat{X} &= (\hat{a} + \hat{a}^\dagger) / \sqrt{2}, & \hat{Y} &= i(\hat{a}^\dagger - \hat{a}) / \sqrt{2}, \\ \hat{x} &= (\hat{b} + \hat{b}^\dagger) / \sqrt{2}, & \hat{y} &= i(\hat{b}^\dagger - \hat{b}) / \sqrt{2},\end{aligned}\quad (12)$$

for the magnon modes \hat{a}, \hat{b} . The dissipative dynamics for magnon modes are described by

$$\dot{\hat{O}} = \frac{i}{\hbar} [\hat{H}, \hat{O}] - \kappa_o \hat{O} + \sqrt{2\kappa_o} \hat{O}^{in}, \quad (13)$$

where κ_o is the dissipation rate, $\hat{O}^{in} = (\hat{o}^{in} + \hat{o}^{in\dagger})/\sqrt{2}$ or $\hat{O}^{in} = i(\hat{o}^{in\dagger} - \hat{o}^{in})/\sqrt{2}$ with $\hat{o} = \hat{a}, \hat{b}$, is the input noise operator associated with the magnon modes $\hat{O} = \hat{X}, \hat{x}$ or $\hat{O} = \hat{Y}, \hat{y}$, respectively. The input noise operators are characterized by the correlation functions

$$\begin{aligned}\langle \hat{o}^{in\dagger}(t) \hat{o}^{in}(t') \rangle &= 2\kappa_o N(\omega_o) \delta(t - t'), \\ \langle \hat{o}^{in}(t) \hat{o}^{in\dagger}(t') \rangle &= 2\kappa_o [N(\omega_o) + 1] \delta(t - t'),\end{aligned}\quad (14)$$

with the equilibrium thermal mean magnon occupation numbers $N(\omega_o) = [\exp(\hbar\omega_o/k_B T) - 1]^{-1}$.

In a similar manner we may define

$$\hat{q} = (\hat{c} + \hat{c}^\dagger) \sqrt{2}, \quad \hat{p} = i(\hat{c}^\dagger - \hat{c}) / \sqrt{2}, \quad (15)$$

to be the dimensionless quadratures associated with the mechanical phonon mode. A Markovian description of quantum Brownian motion for a mechanical mode with large quality factor $Q = \omega_c / \gamma_c \gg 1$ is set by [12, 43, 44]

$$\dot{\hat{q}} = \frac{i}{\hbar} [\hat{H}, \hat{q}], \quad \dot{\hat{p}} = \frac{i}{\hbar} [\hat{H}, \hat{p}] - \gamma_c \hat{p} + \hat{\xi}, \quad (16)$$

where the mechanical dissipation rate γ_c is associated mainly with phonons of non-zero momentum \mathbf{k}' , which are not considered in the Hamiltonian \hat{H} . The input noise operator $\hat{\xi}$ is given by the correlation function

$$\langle \hat{\xi}(t) \hat{\xi}(t') + \hat{\xi}(t') \hat{\xi}(t) \rangle \simeq 2\gamma_c [2N_c(\omega_c) + 1] \delta(t - t'), \quad (17)$$

where $N_c(\omega_c) = [\exp(\hbar\omega_c/k_B T) - 1]^{-1}$ is the equilibrium thermal mean phonon occupation number.

Therefore, the corresponding nonlinear QLEs, in terms of dimensionless quadratures, take the form

$$\begin{aligned} \dot{\hat{X}} &= -\kappa_a \hat{X} + \omega_a \hat{Y} - \text{Im}(g_{ab}) \hat{x} - \text{Re}(g_{ab}) \hat{y} + g_{ac} \hat{Y} \hat{q} + \sqrt{2} \Omega + \hat{X}^{in}, \\ \dot{\hat{Y}} &= -\omega_a \hat{X} - \kappa_a \hat{Y} - \text{Re}(g_{ab}) \hat{x} + \text{Im}(g_{ab}) \hat{y} - g_{ac} \hat{X} \hat{q} + \hat{Y}^{in\dagger}, \\ \dot{\hat{x}} &= -\text{Im}(g_{ab}) \hat{X} - \text{Re}(g_{ab}) \hat{Y} - \kappa_b \hat{x} + \omega_b \hat{y} + g_{bc} \hat{Y} \hat{q} + \sqrt{2} \Omega + \hat{x}^{in}, \\ \dot{\hat{y}} &= -\text{Re}(g_{ab}) \hat{X} + \text{Im}(g_{ab}) \hat{Y} - \omega_b \hat{x} - \kappa_b \hat{y} - g_{bc} \hat{X} \hat{q} + \hat{y}^{in\dagger}, \\ \dot{\hat{q}} &= \omega_c \hat{p}, \\ \dot{\hat{p}} &= -\frac{g_{ac}}{2} (\hat{X}^2 + \hat{Y}^2 - 1) - \frac{g_{bc}}{2} (\hat{x}^2 + \hat{y}^2 - 1) - \omega_c \hat{q} - \gamma_c \hat{p} + \hat{\xi}. \end{aligned} \quad (18)$$

4. Results and discussion

We now show that it is possible to maintain magnon–magnon entanglement in an equilibrium setting. We focus on the steady-state regime, where $\frac{d}{dt} \langle \hat{O} \rangle = 0$ for $\hat{O} = \hat{X}, \hat{Y}, \hat{x}, \hat{y}, \hat{p}, \hat{q}$. Any operator can then be written as a steady-state expectation value plus an additional quantum fluctuation term, i.e. $\hat{O} = \langle \hat{O} \rangle + \delta \hat{O}(t)$ [45]. Imposing the steady-state conditions on equation (18) and retaining fluctuations up to linear order, we obtain a set of linearized QLEs

$$\dot{u}(t) = Au(t) + n(t), \quad (19)$$

where $u(t) = (\delta \hat{X}, \delta \hat{Y}, \delta \hat{x}, \delta \hat{y}, \delta \hat{q}, \delta \hat{p})^T$ and $n(t) = (\hat{X}^{in}, \hat{Y}^{in}, \hat{x}^{in}, \hat{y}^{in}, 0, \hat{\xi})^T$. Information about steady-state expectation values and coupling constants is encoded into the drift matrix:

$$A = \begin{bmatrix} -\kappa_a & \Delta_{ac} & -\text{Im}(g_{ab}) & -\text{Re}(g_{ab}) & M_Y & 0 \\ -\Delta_{ac} & -\kappa_a & -\text{Re}(g_{ab}) & \text{Im}(g_{ab}) & -M_X & 0 \\ -\text{Im}(g_{ab}) & -\text{Re}(g_{ab}) & -\kappa_b & \Delta_{bc} & M_Y & 0 \\ -\text{Re}(g_{ab}) & \text{Im}(g_{ab}) & -\Delta_{bc} & -\kappa_b & -M_X & 0 \\ 0 & 0 & 0 & 0 & 0 & \omega_p \\ -M_X & -M_Y & -M_X & -M_Y & -\omega_p & -\gamma_c \end{bmatrix}. \quad (20)$$

Here, $\Delta_{ac} = \omega_a + g_{ac} \langle \hat{q} \rangle$ and $\Delta_{bc} = \omega_b + g_{bc} \langle \hat{q} \rangle$ are the effective magnon frequency detunings induced by the magnon–phonon interaction. The effective magnon–phonon couplings are given by $M_X = g_{ac} \langle \hat{X} \rangle$, $M_Y = g_{ac} \langle \hat{Y} \rangle$, and $M_x = g_{bc} \langle \hat{x} \rangle$, $M_y = g_{bc} \langle \hat{y} \rangle$. Assuming low dissipation of the magnon modes compared to their frequencies in typical AFM structures [46, 47], i.e. $\kappa_a, \kappa_b \ll \Delta_{ac}, \Delta_{bc}$, we obtain the steady-state solutions as




Figure 2. Phonon-driven magnon-magnon entanglement E_N as a function of temperature and the frequency Ω . Note that Ω is associated with the strength of the off-resonant rotating magnetic field, B_0 (see discussion below equation (10)). The color bar displays a qualitative illustration of the entanglement, where the white area indicates the separable zone while the colored zones show non-zero magnon-magnon entanglement E_N in the $\Omega - T$ plane. Double peaks are associated with the degeneracy of two polarized magnons in AFM systems, which is broken in the presence of non-zero magnetic field frequency ω_0 . The gap between the peaks is proportional to the frequency difference of the two magnons. The ratio ω_0/ω with respect to the magnon frequency at the degeneracy point, i.e. ω , allows one to control the magnon-magnon entanglement region in the $\Omega - T$ parameter space. The lower the ratio ω_0/ω , the weaker the external magnetic field required to achieve high entanglement at a given temperature (see also figure 3). Here, the remaining parameters of the Hamiltonian in equation (11) satisfy $\omega_c/\omega = 1$ and $g_{ac}/g_{ab} = g_{bc}/g_{ab} = 0.001$.

$$\begin{aligned}
 \langle X \rangle &= \frac{\sqrt{2}\Omega \text{Im}(g_{ab})}{\Delta_{ac}\Delta_{bc} - |g_{ab}|^2}, \quad \langle Y \rangle = \frac{-\sqrt{2}\Omega [\Delta_{bc} + \text{Re}(g_{ab})]}{\Delta_{ac}\Delta_{bc} - |g_{ab}|^2}, \\
 \langle x \rangle &= \frac{\sqrt{2}\Omega \text{Im}(g_{ab})}{\Delta_{ac}\Delta_{bc} - |g_{ab}|^2}, \quad \langle y \rangle = \frac{-\sqrt{2}\Omega [\Delta_{ac} + \text{Re}(g_{ab})]}{\Delta_{ac}\Delta_{bc} - |g_{ab}|^2}, \\
 \frac{\omega_p}{\Omega^2} \langle q \rangle + \frac{g_{ac}(\Delta_{bc} + g_{ab})^2 + g_{bc}(\Delta_{ac} + g_{ab})^2}{(\Delta_{ac}\Delta_{bc} - |g_{ab}|^2)^2} &= 0.
 \end{aligned} \tag{21}$$

Once the frequencies and exchange parameters are set as parameters of the model Hamiltonian in equation (11), the explicit form of the drift matrix A mainly depends on the steady-state value of $\langle \hat{q} \rangle$, which is given as a root of the polynomial of degree five specified by the last algebraic expression in equation (21). As a frequency shift in magnon modes, only the real-valued roots of $\langle \hat{q} \rangle$ are physically meaningful. For a given set of parameter values, the polynomial has only one real-valued root.

The linear form of QLEs in equation (19) ensures that the dynamics of quantum fluctuations is Gaussian and can be completely characterized by the corresponding covariance correlation matrix (CCM) [48]. By using the fact that different components of the input noise vector $n(t)$ are uncorrelated, the steady-state CCM denoted by V can be obtained through the Lyapunov equation [12, 44]


$$A \cdot V + V \cdot A^T = -D, \tag{22}$$

where $D = \text{diag}[\kappa_a(2N_a + 1), \kappa_a(2N_a + 1), \kappa_b(2N_b + 1), \kappa_b(2N_b + 1), 0, \gamma_c(2N_p + 1)]$ is the diffusion matrix. Solving equation (22) for V , one can evaluate the bipartite magnon-magnon entanglement by computing the logarithmic negativity of two-mode Gaussian states defined as [12, 27, 48, 49]

$$E_N \equiv \max[0, -\ln(2\tilde{\nu}_-)], \tag{23}$$

where $\tilde{\nu}_-$ is the minimum symplectic eigenvalue of a matrix, V_4 , being the 4×4 reduced CCM obtained by projecting V onto the two magnon modes. This eigenvalue can be computed as $\tilde{\nu}_- = \min \text{eig}[i\Upsilon \tilde{V}_4]$, for the symplectic matrix $\Upsilon = \oplus_{j=1}^2 i\sigma_y$ (with σ_y being the Pauli- y matrix) and the partial transpose $\tilde{V}_4 = P_{1|2} V_4 P_{1|2}$ with $P_{1|2} = \text{diag}(1, -1, 1, 1)$.

The magnon-magnon logarithmic negativity in the steady-state limit as a function of temperature T and external magnetic field strength Ω is shown in figure 2 for $\omega_c/\omega = 1$. Note that in the following numerical analysis, γ_c is taken to be 10^2 Hz, and $\kappa_{a,b} \approx 1$ GHz. As can be seen, our analysis reveals magnon-magnon entanglement at room temperature under the phonon mode effect. The high-temperature result for magnon-magnon entanglement is reproduced across a wide range of Hamiltonian parameters through



complementary plots in figures 3–5. For any given rotation frequency ω_0 of the magnetic field, there always exists a certain range of the field amplitude Ω , where magnon–magnon entanglement can be realized at high temperature. Moreover, the entangled region including the high-temperature entanglement zone in the $\Omega - T$ parameter space can be efficiently placed around practically feasible parameter values by controlling the frequency ratio ω_0/ω . In other words, magnon–magnon entanglement can be realized for any positive value of Ω at a given temperature T , provided the frequency ratio ω_0/ω is properly adjusted. The results depicted in figure 2 were qualitatively replicated across a range of magnon–phonon coupling ratio $g_{ac}/g_{ab} = g_{bc}/g_{ab} \in [10^{-5}, 10^{-3}]$. This shows that, regardless of the strength of the magnon–phonon interaction in the reasonable range relative to the coupling strength between the two magnon modes, a similar outcome as presented in figure 2 can be achieved. Thus, the strength of magnon–phonon interaction, as treated here, does not have a detrimental effect on the phase transition of the magnon–magnon entanglement. It is instead so that, in the steady-state limit, the phonon degrees of freedom have a key role in establishing magnon–magnon entanglement, in the sense that in the absence of phonons, the two magnon modes are completely disentangled. We demonstrate this by noting that the Hamiltonian in equation (11) reduces to


$$\hat{H}_{ab} = \hbar\omega_a\hat{a}^\dagger\hat{a} + \hbar\omega_b\hat{b}^\dagger\hat{b} + \hbar\Omega(\hat{a} + \hat{a}^\dagger) + \hbar\Omega(\hat{b} + \hat{b}^\dagger) + \hbar g_{ab}^*\hat{a}^\dagger\hat{b}^\dagger + \hbar g_{ab}\hat{a}\hat{b}, \quad (24)$$

when the phonon mode is absent. Through the similar method as shown above, we obtain the corresponding drift matrix

$$A_{ab} = \begin{bmatrix} -\kappa_a & \omega_a & -\text{Im}(g_{ab}) & -\text{Re}(g_{ab}) \\ -\omega_a & -\kappa_a & -\text{Re}(g_{ab}) & \text{Im}(g_{ab}) \\ -\text{Im}(g_{ab}) & -\text{Re}(g_{ab}) & -\kappa_b & \omega_b \\ -\text{Re}(g_{ab}) & \text{Im}(g_{ab}) & -\omega_b & -\kappa_b \end{bmatrix} \quad (25)$$

and numerically confirm that $E_N^{ab} = 0$, for any choice of physical parameters of the spin Hamiltonian. We employ the drift matrix from equation (25) to solve equation (22) for the CCM, i.e. V . Subsequently, we determine the minimum eigenvalue ν_- for evaluating for evaluating the logarithmic negativity. The procedure is repeated for a wide range of distinct parameter values within the drift matrix described in equation (25). In all cases, $\tilde{\nu}_-$ consistently exceeds $\frac{1}{2}$ at any temperature, indicating a complete absence of entanglement. This indicates that the nontrivial entanglement diagram obtained in figure 2 is dissipation- and phonon-driven.

To elaborate further on this point, in figure 3 we show how some cuts of the entangled region in figure 2 at three different temperatures move along the parameter domain of Ω upon varying the control parameter ω_0/ω . The plots indicate that for a given value of Ω , there is a narrow interval for ω_0/ω where magnon–magnon entanglement is non-zero and that this interval depends on temperature. The appearance



of two lines in these plots at higher values of ω_0/ω and temperatures corresponds to the two peaks in figure 2. These peaks result from the detuning frequencies of the two magnon modes, where $\omega_a = \omega + \omega_0$ and $\omega_b = \omega - \omega_0$. As ω_0 approaches zero, causing the two magnon modes to become degenerate, the two peaks converge and coincide at $\omega_0 = 0$, as indicated in figure 3.

Figures 2 and 3 are plotted for fixed value of the frequency ratio $\omega_c/\omega = 1$ and fixed ratios g_{ac}/g_{ab} and g_{bc}/g_{ab} , and shows segments of entanglement in a parameter space of magnetic field strength and frequency ratio ω_0/ω . In a search for a real material that can serve as an optimal host for robustly entangled magnon modes, it is instructive to also understand the dependence of entanglement on these material parameters. This is shown in figures 4 and 5. One can see that, in a wide range of couplings and phonon frequencies, it is possible to have non-zero entanglement at rather large (and even ambient) temperatures if frequency and amplitude of the rotating magnetic field are tuned to proper values. Figure 4 demonstrates that for any given magnon–phonon coupling $g_{ac} = g_{bc}$, there always exists a narrow window of phonon frequencies, where the same magnon–magnon entanglement as shown in the figure 2 occurs.

We further examine the effect of asymmetric magnon–phonon couplings on the magnon-magnon entanglement region. For given ω_c/ω , figure 5 illustrates a wide range of different coupling strengths $g_{ac} \neq g_{bc}$, which allows the same entanglement phase diagram as in figure 2. Figure 5 shows that, even up to as high temperatures as $T = 300$ K, for different values of coupling rates varying in a wide range, there is a value of phonon frequency ω_c/ω that guarantees non-zero magnon–magnon entanglement. Besides, it implies that a broad family of antiferromagnetic materials are likely to have a similar magnon–magnon entanglement diagram, as shown in figure 2.

There are numerous suitable classes of AFM compounds that can maintain the steady-state entanglement of magnon modes within the AFM structure, even up to room temperature. Among them, NiO is a concrete example of a crystalline AFM [50–53] with high Néel temperature of $T_N^{\text{NiO}} = 523$ K and large magnon lifetime. Notably, it has been shown that the zone center phonon frequency matches the magnon frequency at finite \mathbf{k} frequencies ~ 11.3 THz and ~ 17.3 THz [28]. The perovskite manganites also form a promising class of materials in this regard, with many compounds of varying chemical composition that have been synthesized (see, e.g. [31, 32]). For these systems, upon chemical modulation, both the magnetic and lattice properties can be tuned. This has particular relevance for finding compositions that allow for finer adjustments of the magnon and phonon frequencies relevant for the results of the present investigation.

We end our discussion with a remark that the dissipative magnon entanglement studied here is of the mixed-state type and is achieved in the asymptotic steady-state limit. This is different from the entanglement observed in the pure energy eigenstates of ideal closed magnon systems studied previously in [23].




Figure 5. The range of asymmetric magnon-phonon couplings, which allows for the same magnon-magnon entanglement pattern in figure 2, for a given phonon frequency. The contour plots correspond to the non-zero magnon-magnon entanglement segment at $\Omega = 0.1932$ THz, $\omega_0/\omega = 0.07$ and temperature $T = 300$ K ($T = 100$ K for inset) for $\omega_c/\omega = 0.75$ (yellow), $\omega_c/\omega = 1$ (pink) and $\omega_c/\omega = 1.25$ (blue). The uncolored regions depict zero-entanglement (separable). Similar patterns occur for any other point in the entangling diagram of figure 2 and any other value of ω_c/ω . The higher the temperature the narrower the range of asymmetric magnon-phonon couplings for a given ω_c/ω .

5. Conclusion

We have examined bipartite magnon-magnon entanglement in a general setting of an AFM material subjected to a dissipative elastic displacement (phonon) mode, thermal noise, and an external rotating magnetic field. We have provided evidence that it is possible to observe phonon-driven entanglement between the magnon modes of the two sublattices even at room temperature. The high-temperature magnon-magnon entanglement for a given material and magnon-phonon coupling, can be ensured by tuning the external magnetic field frequency and amplitude in a physically reasonable range of scales. The presence of a non-zero magnon-phonon coupling is necessary for maintaining entanglement between magnons in antiferromagnets in a dissipative steady-state regime, while the strength of the coupling does not significantly affect the existence of this entanglement. It would be interesting to examine whether AFMs can be used as a resource for different quantum information applications, for instance quantum state transfer [54], and to study the potential role of magnon-magnon entanglement in such processes.

Data availability statement

All data that support the findings of this study are included within the article (and any supplementary files).

Acknowledgments

The authors acknowledge financial support from the Knut and Alice Wallenberg Foundation through Grants No. 2018.0060 and No. 2021.0246. A Bergman and O E acknowledges eSENCE. A D acknowledges financial support from the Swedish Research Council (VR) through Grants No. 2019-05304 and 2016-05980. O E acknowledges support by the Swedish Research Council (VR), the Foundation for Strategic Research (SSF), the European Research Council (854843-FASTCORR) and STandUP. D T acknowledges support from the Swedish Research Council (VR) with Grant No. 2019-03666. A D and O E acknowledge support from the Wallenberg Initiative Materials Science for Sustainability (WISE) funded by the Knut and Alice Wallenberg Foundation (KAW). Y L acknowledges financial support from the KTH-CSC scholarship agreement (No. 201907090094). E S acknowledges financial support from the Swedish Research Council (VR) through Grant No. 2017-03832.

ORCID iDs

Manuel Pereiro <https://orcid.org/0000-0003-2977-255X>

Vahid Azimi-Mousolou <https://orcid.org/0000-0002-7838-0053>

References

- [1] Browne D, Bose S, Mintert F and Kim M S 2017 From quantum optics to quantum technologies *Prog. Quantum Electron.* **54** 2–18
- [2] Galve F, Pachón L A and Zueco D 2010 Bringing entanglement to the high temperature limit *Phys. Rev. Lett.* **105** 180501
- [3] Dolde F, Jakobi I, Naydenov B, Zhao N, Pezzagna S, Trautmann C, Meijer J, Neumann P, Jelezko F and Wrachtrup J 2013 Room-temperature entanglement between single defect spins in diamond *Nat. Phys.* **9** 139
- [4] Wang F et al 2017 Room-temperature storage of quantum entanglement using decoherence-free subspace in a solid-state spin system *Phys. Rev. B* **96** 134314
- [5] Lin Q, He B and Xiao M 2020 Entangling two macroscopic mechanical resonators at high temperature *Phys. Rev. Appl.* **13** 034030
- [6] Zhong H-S et al 2020 Quantum computational advantage using photons *Science* **370** 1460
- [7] Arrazola J M et al 2021 Quantum circuits with many photons on a programmable nanophotonic chip *Nature* **591** 54
- [8] Gulka M, Wirtitsch D, Ivády V, Vodnik J, Hruba J, Magchiels G, Bourgeois E, Gali A, Trupke M and Nesladek M 2021 Room-temperature control and electrical readout of individual nitrogen-vacancy nuclear spins *Nat. Commun.* **12** 4421
- [9] Bello F D, Kongsuwan N and Hess O 2022 Near-field generation and control of ultrafast, multipartite entanglement for quantum nanoplasmonic networks *Nano Lett.* **22** 2801
- [10] Barman A et al 2021 The 2021 magnonics roadmap *J. Phys.: Condens. Matter* **33** 413001
- [11] Yuan H Y, Cao Y, Kamra A, Duine R A and Yan P 2022 Quantum magnonics: when magnon spintronics meets quantum information science *Phys. Rep.* **965** 1
- [12] Li J, Zhu S-Y and Agarwal G S 2018 Magnon-photon-phonon entanglement in cavity magnomechanics *Phys. Rev. Lett.* **121** 203601
- [13] Toklikishvili Z, Chotorlishvili L, Khomeriki R, Jandieri V V and Berakdar J J 2023 Electrically controlled entanglement of cavity photons with electromagnons *Phys. Rev. B* **107** 115126
- [14] Tabuchi Y, Ishino S, Noguchi A, Ishikawa O, Yamazaki R, Usami K and Nakamura Y 2016 Quantum magnonics: the magnon meets the superconducting qubit *C. R. Physique* **17** 729
- [15] Lachance-Quirion D, Tabuchi Y, Glöppe A, Usami K and Nakamura Y 2019 Hybrid quantum systems based on magnonics *Appl. Phys. Express* **12** 070101
- [16] Bossini D et al 2019 Laser-driven quantum magnonics and terahertz dynamics of the order parameter in antiferromagnets *Phys. Rev. B* **100** 024428
- [17] Kamra A, Thingstad E, Rastelli G, Duine R A, Brataas A, Belzig W and Sudbø A 2019 Antiferromagnetic magnons as highly squeezed Fock states underlying quantum correlations *Phys. Rev. B* **100** 174407
- [18] Yuan H Y, Zheng S, Ficek Z, He Q Y and Yung M-H 2020 Enhancement of magnon-magnon entanglement inside a cavity *Phys. Rev. B* **101** 014419
- [19] Wührer D, Rohling N and Belzig W 2022 Theory of quantum entanglement and structure of the two-mode squeezed antiferromagnetic magnon vacuum *Phys. Rev. B* **105** 054406
- [20] Kamra A, Belzig W and Brataas A 2020 Magnon-squeezing as a niche of quantum magnonics *Appl. Phys. Lett.* **117** 090501
- [21] Yuan H Y, Kamra A, Hartmann D M F and Duine R A 2021 Electrically switchable entanglement channel in van der Waals magnets *Phys. Rev. Appl.* **16** 024047
- [22] Rezende S M, Azevedo A and Rodríguez-Suárez R L 2019 Introduction to antiferromagnetic magnons *J. Appl. Phys.* **126** 151101
- [23] Azimi-Mousolou V, Bagrov A, Bergman A, Delin A, Eriksson O, Liu Y, Pereiro M, Thonig D and Sjöqvist E 2020 Hierarchy of magnon mode entanglement in antiferromagnets *Phys. Rev. B* **102** 224418
- [24] Azimi-Mousolou V, Liu Y, Bergman A, Delin A, Eriksson O, Pereiro M, Thonig D and Sjöqvist E 2021 Magnon-magnon entanglement and its quantification via a microwave cavity *Phys. Rev. B* **104** 224302
- [25] Azimi-Mousolou V, Bergman A, Delin A, Eriksson O, Pereiro M, Thonig D and Sjöqvist E 2023 Transmon probe for quantum characteristics of magnons in antiferromagnets *Phys. Rev. B* **108** 094430
- [26] Shiranzaei M, Fransson J and Azimi-Mousolou V 2023 Temperature-anisotropy conjugate magnon squeezing in antiferromagnets *Phys. Rev. B* **108** 144302
- [27] Vidal G and Werner R F 2002 Computable measure of entanglement *Phys. Rev. A* **65** 032314
- [28] Simensen H T, Troncoso R E, Kamra A and Brataas A 2019 Magnon-polarons in cubic collinear antiferromagnets *Phys. Rev. B* **99** 064421
- [29] Mai T T, Garrity K F, McCreary A, Argo J, Simpson J R, Doan-Nguyen V, Valdés Aguilar R and Hight Walker A R 2021 Magnon-phonon hybridization in 2D antiferromagnet MnPSe₃ *Sci. Adv.* **9** eabj3106
- [30] Liu S et al 2021 Direct observation of magnon-phonon strong coupling in two-dimensional antiferromagnet at high magnetic fields *Phys. Rev. Lett.* **127** 097401
- [31] Dai C and Ma F 2021 Strong magnon-magnon coupling in synthetic antiferromagnets *Appl. Phys. Lett.* **118** 112405
- [32] Rini E G, Rao M N, Chaplot S L, Gaur N K and Singh R K 2007 Phonon dynamics of lanthanum manganite LaMnO₃ using an interatomic shell model potential *Phys. Rev. B* **75** 214301
- [33] Ulbrich H, Krüger F, Nugroho A A, Lamago D, Sidis Y and Braden M 2011 Spin-wave excitations in the ferromagnetic metallic and in the charge-, orbital- and spin-ordered states in Nd_{1-x}Sr_xMnO₃ with $x \approx 0.5$ *Phys. Rev. B* **84** 094453
- [34] Rodríguez E E, Poineau F, Llobet A, Kennedy B J, Avdeev M, Thorogood G J, Carter M L, Seshadri R, Singh D J and Cheetham A K 2011 High temperature magnetic ordering in the 4d perovskite SrTcO₃ *Phys. Rev. Lett.* **106** 067201
- [35] Avdeev M, Thorogood G J, Carter M L, Kennedy B J, Ting J, Singh D J and Wallwork K S 2011 Antiferromagnetism in a technetium oxide. Structure of CaTcO₃ *J. Am. Chem. Soc.* **133** 1654
- [36] Palkar V R, John J and Pinto R 2002 Observation of saturated polarization and dielectric anomaly in magnetoelectric BiFeO₃ thin films *Appl. Phys. Lett.* **80** 1628–30
- [37] Sinha K P and Upadhyaya U N 1962 Phonon-magnon interaction in magnetic crystals *Phys. Rev.* **127** 432
- [38] Streib S, Vidal-Silva N, Shen K and Bauer G E W 2019 Magnon-phonon interactions in magnetic insulators *Phys. Rev. B* **99** 184442
- [39] Zhang X, Zou C-L, Jiang L and Tang H X 2016 Cavity magnomechanics *Sci. Adv.* **2** e1501286
- [40] Woods L M 2001 Magnon-phonon effects in ferromagnetic manganites *Phys. Rev. B* **65** 014409
- [41] Rückriegel A, Kopietz P, Bozhko D A, Serga A A and Hillebrands B 2014 Magnetoelastic modes and lifetime of magnons in thin yttrium iron garnet films *Phys. Rev. B* **89** 184413
- [42] Sanavio C, Peano V and Xuereb A 2020 Nonreciprocal topological phononics in optomechanical arrays *Phys. Rev. B* **101** 085108
- [43] Giovannetti V and Vitali D 2001 Phase-noise measurement in a cavity with a movable mirror undergoing quantum Brownian motion *Phys. Rev. A* **63** 023812

- [44] Vitali D, Gigan S, Ferreira A, Böhm H R, Tombesi P, Guerreiro A, Vedral V, Zeilinger A and Aspelmeyer M 2007 Optomechanical entanglement between a movable mirror and a cavity fields *Phys. Rev. Lett.* **98** 030405
- [45] Mancini S and Tombesi P 1994 Quantum noise reduction by radiation pressure *Phys. Rev. A* **49** 4055
- [46] Kruglyak V V, Demokritov S O and Grundler D 2010 Magnonics *J. Phys. D: Appl. Phys.* **43** 260301
- [47] Lenk B, Ulrichs H, Garbs F and Münzenberg M 2011 The building blocks of magnonics *Phys. Rep.* **507** 107
- [48] Adesso G, Serafini A and Illuminati F 2004 Extremal entanglement and mixedness in continuous variable systems *Phys. Rev. A* **70** 022318
- [49] Plenio M B 2005 The logarithmic negativity: a full entanglement monotone that is not convex *Phys. Rev. Lett.* **95** 090503
- [50] Roth W L 1958 Magnetic structures of MnO, FeO, CoO and NiO *Phys. Rev.* **101** 1333
- [51] Moriyama T, Hayashi K, Yamada K, Shima M, Ohya Y and Ono T 2019 Intrinsic and extrinsic antiferromagnetic damping in NiO *Phys. Rev. Mater.* **3** 051402
- [52] Tomlinson J R, Domash L, Hay R G and Montgomery C W 1955 The high temperature heat content of nickel oxide *J. Am. Chem. Soc.* **77** 909
- [53] Momida H and Oguchi T 2003 First-principles studies of antiferromagnetic MnO and NiO surfaces *J. Phys. Soc. Japan* **72** 588
- [54] Cirac J I, Zoller P, Kimble H J and Mabuchi H 1997 Quantum state transfer and entanglement distribution among distant nodes in a quantum network *Phys. Rev. Lett.* **78** 3221