

LETTER

Association of *MGMT* and *BIN1* genes with Alzheimer's disease risk across sex and *APOE* ϵ 4 status

Chung et al. reported a novel association of the Alzheimer's disease (AD) risk with genetic variants in the *MGMT* gene in women.¹ The genome-wide significant signals were found in women lacking the apolipoprotein E ϵ 4 allele (*APOE* ϵ 4-) from 30 studies of the Alzheimer's Disease Genetics Consortium (ADGC) (3399 AD cases and 6905 controls), and in a Hutterite cohort (31 members of a consanguineous kindred with different *APOE* ϵ 4 statuses, including 5 AD cases who were all women). The effect sizes reported were large: odds ratio [OR] = 1.44 [1.26–1.64], $P = 4.95 \times 10^{-8}$ in ADGC for rs12775171, and OR = 2.02 [1.80–2.26], $P = 1.9 \times 10^{-14}$ in the Hutterites for rs2803456 and rs12256016. The association found in the ADGC was consistent across studies and not significant in the three other subsets defined by sex and *APOE* ϵ 4 status (women *APOE* ϵ 4+, men *APOE* ϵ 4-, and men *APOE* ϵ 4+) for which effect sizes were not reported.

We aimed at replicating the association of *MGMT* with AD risk in the meta-analysis of 6 case-control studies from the European Alzheimer & Dementia Biobank (EADB) consortium: EADB-core,² EADI (European Alzheimer's Disease Initiative),^{3,4} GERAD (Genetic and Environmental Risk in AD),⁵ DemGene,⁶ GR@ACE-DEGESCO,⁷ and Bonn.² We considered a total of 33,677 AD cases and 48,158 controls, all of European ancestry, including 10,354 AD cases and 19,910 controls who were female and *APOE* ϵ 4- (Figure 1, Tables S1, and S2 in supporting information). The samples were genotyped with different chips and then imputed using the TOPMed reference panel² (supporting infor-

mation). In each study, we tested the association of *MGMT* variants with AD in the four subsets defined by sex and *APOE* ϵ 4 status. Analyses were adjusted on principal components, and results were combined across studies in a fixed effect meta-analysis with an inverse-variance weighted approach (supporting information).

None of the *MGMT* variants identified by Chung et al. were found to be associated with AD risk ($P < 0.05$) in the different subsets (Figures S1–S6 in supporting information). The effect of rs12775171 was larger in *APOE* ϵ 4- women (OR = 1.06 [0.98–1.14], $P = 0.17$) than in the other subsets (OR = 1.03, 1.00, and 0.99 in *APOE* ϵ 4- men, *APOE* ϵ 4+ women, and *APOE* ϵ 4+ men, respectively), but those differences were not significant ($P = 0.69, 0.37, \text{ and } 0.39$ for the comparison of the OR in *APOE* ϵ 4- women with the one in *APOE* ϵ 4- men, *APOE* ϵ 4+ women, and *APOE* ϵ 4+ men, respectively, Figure 1). Of note, our study in *APOE* ϵ 4- women had more than 99% power to detect the association with rs12775171 as described by Chung et al., at the nominal significance level of 0.05 (supporting information).

The authors also identified in ADGC *APOE* ϵ 4- women a genome-wide significant association with AD for a known AD gene, *BIN1* (rs11680911, OR = 1.21 [1.13–1.29], $P = 2.22 \times 10^{-8}$). We sought to assess whether this association differed across the four sex-*APOE* ϵ 4 subsets. We detected a genome-wide significant association ($P < 5 \times 10^{-8}$) with AD risk for rs11680911 in *APOE* ϵ 4- women (OR = 1.12 [1.07–1.16], $P = 2.21 \times 10^{-8}$) and in *APOE* ϵ 4- men (OR = 1.16 [1.10–1.21], $P = 1.75 \times 10^{-9}$), but not in the 2 other




FIGURE 1 Results of rs12775171 association with Alzheimer's disease (AD) risk in apolipoprotein E (*APOE*) ϵ 4- women and the other sex-*APOE* ϵ 4 subsets compared with the effect reported in the Alzheimer's Disease Genetics Consortium (ADGC) *APOE* ϵ 4- women from Chung et al. 2022. The effect allele is G with a frequency of 0.06 in all models. The black square whose size is proportional to the sample size represents the odds ratio (OR) and the blue line the confidence interval (CI). Interaction P are p-values of the heterogeneity test between the different group pairs (1 degree of freedom test) using the *APOE* ϵ 4- women group as a reference for each test (supporting information). EA, Effect allele; P, p-value.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Alzheimer's & Dementia* published by Wiley Periodicals LLC on behalf of Alzheimer's Association.

subsets (Figures S7–S8 in supporting information). However, the effects in all the subsets were similar (OR = 1.12, 1.16, 1.14, and 1.13 in APOEε4- women, APOEε4- men, APOEε4+ women, and APOEε4+ men, respectively), and effects were not significantly different between the subsets (Figures S7–S8 and Table S3 in supporting information).

We performed several sensitivity analyses in the EADB studies (supporting information, Tables S1–S4 and Figures S1–S12), but none of them identified a significant association of MGMT with AD risk in APOEε4- women or differences of association between subsets in BIN1.

In conclusion, we did not find a significant, nor suggestive association of the MGMT variants identified by Chung et al. with AD risk, in any of the subsets defined by sex and APOEε4 status, where our sample size was up to three times larger than in the original publication. Additionally, we did not identify a significant effect difference of BIN1 rs11680911 variant across sex and APOEε4 status subsets.

ACKNOWLEDGMENTS

We thank the many study participants, researchers, and staff for collecting and contributing to the data of the EADB consortium.

CONFLICT OF INTEREST STATEMENT

Martin Ingelsson is a paid consultant to BioArtic.

FUNDING INFORMATION

This study was supported by a grant from the Fondation pour la Recherche sur Alzheimer, convention 2022-A-01, the JPco-fuND-2 “Multinational research projects on Personalized Medicine for Neurodegenerative Diseases” PREADAPT project (ANR-19-JPW2-0004), and the JPco-fuND EADB grant. Ole Andreassen was supported by the Research Council of Norway (RCN grants 223273, 283799, 324252, 344121). Alfredo Ramirez was supported by the German Federal Ministry of Education and Research (BMBF: 01ED1619A). Agustin Ruiz was supported by GRIFOLS-GR@ACE DEGESCO, LA CAIXA-GR@ACE DEGESCO and ISCIII-Ministry of Health Spain. Rebecca Sims was supported by the Medical Research Council UK. Julie Williams was supported by UKDRI-IPSC Platform to Model Alzheimer's Disease Risk (IPMAR).

Julie Le Borgne¹

EADB, GR@ACE, Degesco, EADI, GERAD, DemGene

Philippe Amouyel¹

Ole Andreassen²

Ruth Frikke-Schmidt^{3,4}

Mikko Hiltunen⁵

Martin Ingelsson^{6,7,8}

Alfredo Ramirez^{9,10,11,12,13}

Giacomina Rossi¹⁴

Agustin Ruiz^{15,16}

Pascual Sanchez-Juan^{16,17}

Rebecca Sims¹⁸

Kristel Slegers^{19,20}

Magda Tsolaki^{21,22}

Sven J. van der Lee^{23,24,25}

Julie Williams^{18,26}

Jean-Charles Lambert¹

Céline Bellenguez¹

¹Univ. Lille, Inserm, CHU Lille, Institut Pasteur Lille, LabEx DISTALZ - U1167 - RID-AGE - Facteurs de risque et déterminants moléculaires des maladies liées au vieillissement, Lille, France

²NORMENT Centre, University of Oslo, Oslo, Norway

³Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark

⁴Department of Clinical Biochemistry, Rigshospitalet, Copenhagen, Denmark

⁵Institute of Biomedicine, University of Eastern Finland, Kuopio, Finland

⁶Department of Public Health and Caring Sciences/Geriatrics, Uppsala University, Uppsala, Sweden

⁷Krembil Brain Institute, University Health Network, Toronto, Ontario, Canada

⁸Tanz Centre for Research in Neurodegenerative Diseases, Departments of Medicine and Laboratory Medicine & Pathobiology, University of Toronto, Toronto, Ontario, Canada

⁹Department of Neurodegenerative Diseases and Geriatric Psychiatry, University Hospital Bonn, Bonn, Germany

¹⁰Division of Neurogenetics and Molecular psychiatry, Department of Psychiatry and Psychotherapy, University of Cologne, Medical Faculty, Cologne, Germany

¹¹German Center for Neurodegenerative Diseases (DZNE Bonn), Bonn, Germany

¹²Glenn Biggs Institute for Alzheimer's & Neurodegenerative Diseases, University of Texas Health Sciences Center, San Antonio, Texas, USA

¹³Cluster of Excellence on Cellular Stress responses in Aging-Associated Diseases (CECAD), University of Cologne, Cologne, Germany

¹⁴Fondazione IRCCS, Istituto Neurologico Carlo Besta, Milan, Italy

¹⁵Research Center and Memory Clinic Fundació ACE, Institut Català de Neurociències Aplicades, Universitat Internacional de Catalunya, Barcelona, Spain

¹⁶CIBERNED, Network Center for Biomedical research in Neurodegenerative Diseases, National Institute of Health Carlos III, Madrid, Spain

¹⁷Alzheimer's Centre Reina Sofia-CIEN Foundation, Madrid, Spain

¹⁸Division of Psychological Medicine and Clinical Neuroscience, School of Medicine, Cardiff University, Cardiff, UK

¹⁹Complex Genetics of Alzheimer's Disease Group, VIB Center for Molecular Neurology, VIB, Antwerp, Belgium

²⁰Department of Biomedical Sciences, University of Antwerp, Antwerp, Belgium

²¹First Department of Neurology, Medical School, Aristotle University of Thessaloniki, Thessaloniki, Greece

²²Alzheimer Hellas, Thessaloniki, Greece

²³Genomics of Neurodegenerative Diseases and Aging, Human Genetics, Vrije Universiteit Amsterdam, Amsterdam UMC location VUmc, Amsterdam, The Netherlands

²⁴Alzheimer Center Amsterdam, Neurology, Vrije Universiteit Amsterdam, Amsterdam UMC location VUmc, Amsterdam, The Netherlands

²⁵Amsterdam Neuroscience, Neurodegeneration, Amsterdam, The Netherlands

²⁶UKDRI@Cardiff, School of Medicine, Cardiff University, Cardiff, UK

Correspondence

Céline Bellenguez, Inserm UMR-1167, Institut Pasteur de Lille, 1 rue du Professeur Calmette, BP 245 - 59019 Lille, cedex, France.

Email: celine.bellenguez@pasteur-lille.fr

REFERENCES

1. Chung J, Das A, Sun X, et al. Genome-wide association and multi-omics studies identify MGMT as a novel risk gene for Alzheimer's disease among women. *Alzheimer's Dement*. 2023; 19:896-908. doi:[10.1002/alz.12719](https://doi.org/10.1002/alz.12719)
2. Bellenguez C, Küçükali F, Jansen IE, et al. New insights into the genetic etiology of Alzheimer's disease and related dementias. *Nat Genet*. 2022;54(4):412-436.
3. 3C Study Group. Vascular factors and risk of dementia: design of the Three-City Study and baseline characteristics of the study population. *Neuroepidemiology*. 2003;22:316-325.
4. Lambert JC, Heath S, Even G, et al. Genome-wide association study identifies variants at CLU and CR1 associated with Alzheimer's disease. *Nat Genet*. 2009;41:1094-1099.
5. Harold D, Abraham R, Hollingworth P, et al. Genome-wide association study identifies variants at CLU and PICALM associated with Alzheimer's disease. *Nat Genet*. 2009;41:1088-1093.
6. Jansen IE, Savage JE, Watanabe K, et al. Genome-wide meta-analysis identifies new loci and functional pathways influencing Alzheimer's disease risk. *Nat Genet*. 2019;51(3):405s4-413.
7. Moreno-Grau S, de Rojas I, Hernández I, et al. Genomeswide association analysis of dementia and its clinical endophenotypes reveal novel loci associated with Alzheimer's disease and three causality networks: the GR@ACE project. *Alzheimer's Dement*. 2009;15:1333-1347.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.