

## REDUCED THEORY OF SYMMETRIC AND ANTISYMMETRIC EXCHANGE INTERACTIONS IN NANOWIRES

GIOVANNI DI FRATTA<sup>1</sup>, FILIPP N. RYBAKOV<sup>2</sup> AND  
VALERIY SLASTIKOV<sup>3,\*</sup>

**Abstract.** We investigate the behavior of minimizers of perturbed Dirichlet energies supported on a wire generated by a regular simple curve  $\gamma$  and defined in the space of  $\mathbb{S}^2$ -valued functions. The perturbation  $K$  is represented by a matrix-valued function defined on  $\mathbb{S}^2$  with values in  $\mathbb{R}^{3 \times 3}$ . Under natural regularity conditions on  $K$ , we show that the family of perturbed Dirichlet energies converges, in the sense of  $\Gamma$ -convergence, to a simplified energy functional on  $\gamma$ . The reduced energy unveils how part of the antisymmetric exchange interactions contribute to an anisotropic term whose specific shape depends on the curvature of  $\gamma$ . We also discuss the significant implications of our results for studies of ferromagnetic nanowires when Dzyaloshinskii–Moriya interaction (DMI) is present.

**Mathematics Subject Classification.** 49S05, 35C20, 35Q51, 82D40.

Received July 22, 2024. Accepted November 30, 2024.

### 1. INTRODUCTION AND MOTIVATION

The motivation for our study comes from investigating the behavior of ferromagnetic nanowires. Microscopically ferromagnetic nanowires such as nanotubes, nanowhiskers, or nanostrips can exhibit unique physical properties. These include anisotropic magnetoresistance in helimagnets [1, 2], proximity exchange fields in ferromagnet–superconductor heterostructures [3, 4], fast domain wall motion [5], impact of wire twisting onto magnetic textures [6], and many others [7, 8]. Techniques for fabricating magnetic nanowires make available a wide variety of cross-sections and shapes, from almost perfectly straight to braided [6, 9]. Of significant interest for research and applications are wireframes and networks made of magnetic nanowires [10–13].

The theoretical description of the statics and dynamics of magnetization in nanomagnets is a challenging task even from the numerical point of view due to the nonlinear, nonconvex, and nonlocal nature of the problem, see, *e.g.*, the monograph [14], the review articles [15, 16] and the references therein. Brute-force numerical simulations are typically performed within the framework of 3D micromagnetic model [13, 17–19], which are challenging and often time-consuming.

---

*Keywords and phrases:* Dirichlet energy, harmonic maps,  $\Gamma$ -convergence, micromagnetics, magnetic nanowires, Dzyaloshinskii–Moriya interaction, domain walls.

<sup>1</sup> Dipartimento di Matematica e Applicazioni “R. Caccioppoli”, Università degli Studi di Napoli “Federico II”, Via Cintia, Complesso Monte S. Angelo, 80126 Napoli, Italy.

<sup>2</sup> Department of Physics and Astronomy, Uppsala University, Box-516, Uppsala SE-751 20, Sweden.

<sup>3</sup> School of Mathematics, University of Bristol, University Walk, Bristol BS8 1UG, UK.

\* Corresponding author: [valeriy.slastikov@bristol.ac.uk](mailto:valeriy.slastikov@bristol.ac.uk)

While an essential goal of quantitative simulations is description, the basic explanation of magnetic phenomena typically requires having simplified yet accurate models at hand. From the mathematical point of view, such models can be derived by dimension reduction techniques of the calculus of variations as  $\Gamma$ -limits of the micromagnetic energy functional under suitable asymptotic relations among the material and geometric scaling parameters. Dimensional reduction arguments in micromagnetics are a subject with a long history as they date back to the seminal paper [20], where the authors show that in *planar* thin films, the nonlocal effects of the stray field operator reduce to an easy-plane anisotropy term. Later, various static reduced theories for planar thin-film micromagnetics were established under different scaling regimes [21–27]. Further asymptotic regimes, *e.g.*, in the presence of strong perpendicular anisotropy, have been the subject of [28]. The induced thin-film dynamics have been analyzed in [29–32].

The latest developments in nanotechnology have made it possible to create nanostructures, with thicknesses as small as a few atomic layers and lateral sizes down to tens of nanometers. These structures often demonstrate the impact of interfacial effects, such as the *Dzyaloshinskii-Moriya interaction* (DMI) [33, 34], leading to the formation of magnetic skyrmions [35]. Mathematical studies of magnetic domain walls in the regime of planar thin films, taking into account also DMI, have been done in [36–39].

In recent years, interest has grown in ferromagnetic systems with a *curved* shape due to their capability of hosting magnetic skyrmions even in materials where the DMI can be neglected. The evidence of these states sheds light on the role of the geometry in magnetism: chiral spin-textures can be stabilized by curvature effects only, in contrast to the planar case where DMI is required (we refer the reader to the recent monograph [8]). From the mathematical perspective, dimension reduction results for curved thin films have been studied in [21, 27, 40, 41] and, more recently, in [42] where the curved thin film limit of chiral Dirichlet energies is derived.

The effective magnetic behavior of ferromagnetic nanowires has been investigated in [43] where both long-range dipolar interactions and symmetric exchange interactions are considered, also for curved wire. The main aim of this paper is to complement the analysis in [43] by taking into account general antisymmetric exchange interactions. As in [36] and [42], the derived limiting model reveals new physics: part of the antisymmetric exchange interactions contributes to an increase in the shape anisotropy originating from the magnetostatic self-energy. Several examples of applications to the micromagnetic theory complete our analytical derivation.

## 2. CONTRIBUTIONS OF THE PRESENT WORK

Given a regular simple curve<sup>1</sup>  $\gamma : I \mapsto \mathbb{R}^3$  and a smooth domain  $Q \subseteq \mathbb{R}^2$ , for every  $\varepsilon > 0$ , we denote by  $\Omega_\varepsilon \subseteq \mathbb{R}^3$  the  $\varepsilon$ -tube along  $\gamma$  with cross-sectional shape  $Q$  and thickness  $\varepsilon$ , defined as the image of the cylindrical region  $I \times Q \subseteq \mathbb{R} \times \mathbb{R}^2$  through the parameterization (see Fig. 1)

$$\varphi_\varepsilon : (s, z) \in I \times Q \mapsto \gamma(s) + \varepsilon z_1 \mathbf{n}(s) + \varepsilon z_2 \mathbf{b}(s) \in \Omega_\varepsilon. \quad (2.1)$$

Here, we set  $z := (z_1, z_2)$  and denoted by  $(\mathbf{t}(s), \mathbf{n}(s), \mathbf{b}(s))$  the FRENET–SERRET frame consisting, respectively, of the tangent, normal, and binormal unit vectors to the curve at  $\gamma(s)$  – if  $\gamma$  is a straight line, we complete  $\mathbf{t}$  to an orthonormal basis of  $\mathbb{R}^3$ . In what follows, we assume that  $\gamma$  is parameterized by arc length. Also, we assume that the interval  $I$  is compact so that according to the tubular neighborhood theorem, there exists a  $\delta > 0$  such that for every  $0 < \varepsilon < \delta$ , the map  $\varphi_\varepsilon$  defined by (2.1) is a diffeomorphism of  $I \times Q$  onto the  $\varepsilon$ -tube along  $\gamma$ .

The main aim of this paper is to investigate the curved thin-wire limit ( $\varepsilon \rightarrow 0$ ) of the perturbed Dirichlet energy ( $0 < \varepsilon < \delta$ )

$$\mathcal{G}_\varepsilon(\mathbf{v}) := \frac{1}{2\varepsilon^2|Q|} \int_{\Omega_\varepsilon} |D\mathbf{v}(x) + K(\mathbf{v}(x))|^2 dx, \quad (2.2)$$

<sup>1</sup>By a regular simple curve, we mean the image of a  $C^2$ -map  $\gamma : I \mapsto \mathbb{R}^3$ ,  $I \subseteq \mathbb{R}$  a compact interval, such that  $\partial_s \gamma(s) \neq 0$  for every  $s \in I$ , and with no self-intersections, *i.e.*, such that the only possible loss of injectivity in  $\gamma$  arises at the endpoints of  $I$ , case in which the curve closes into a loop.




FIGURE 1. Sketches of suitable wires. The wires can be straight or curved and can have a nonsmooth boundary or a multiply connected cross-section (not every case is shown in the picture). The cross-section is assumed to be invariant along the wire.

defined on  $H^1$  Sobolev maps  $\mathbf{v} : \Omega_\varepsilon \rightarrow \mathbb{S}^2$ , where for every  $\varepsilon > 0$  the domain  $\Omega_\varepsilon \subseteq \mathbb{R}^3$  is an  $\varepsilon$ -tube of  $\mathbb{R}^3$ , and  $\mathbb{S}^2$  is the two-sphere in  $\mathbb{R}^3$ . In writing (2.2), we have made the common and convenient abuse of denoting by  $|\cdot|$  both the Euclidean norm on  $\mathbb{R}^{3 \times 3}$  and the Lebesgue measure when applied to sets, as in the expression  $|Q|$ .

In physical terms,  $\mathcal{G}_\varepsilon$  represents a free energy functional that captures the cost of deformations in the magnetization configuration  $\mathbf{v}$  under both exchange interactions and additional effects from antisymmetric exchange interactions, such as the Dzyaloshinskii–Moriya interaction, encapsulated by the operator  $K$  (see Sect. 3).

The perturbation  $K : \sigma \in \mathbb{S}^2 \mapsto K(\sigma) \in \mathbb{R}^{3 \times 3}$  is represented by a matrix-valued function defined on  $\mathbb{S}^2$  and with values in  $\mathbb{R}^{3 \times 3}$  and is assumed to be Lipschitz continuous, *i.e.*, that there exists  $c_K > 0$  such that

$$|K(\sigma_1) - K(\sigma_2)|_{n \times m} \leq c_K |\sigma_1 - \sigma_2|_m \quad \forall \sigma_1, \sigma_2 \in \mathbb{S}^2. \quad (2.3)$$

Since  $\mathbb{S}^2$  is compact, the continuity condition (2.3) implies that the image of  $K$  is bounded (in fact, compact). In what follows, to simplify notation, we will assume that a Lipschitz constant  $c_K$  is chosen big enough so that there holds

$$|K(\sigma)| \leq c_K \quad \forall \sigma \in \mathbb{S}^2. \quad (2.4)$$

For any  $0 < \varepsilon < \delta$ , the existence of at least a minimizer for  $\mathcal{G}_\varepsilon$  in  $H^1(\Omega_\varepsilon, \mathbb{S}^2)$  is a simple application of the direct method of the calculus of variations. We are interested in the asymptotic behavior of the family of minimizers of  $(\mathcal{G}_\varepsilon)_\varepsilon$  as  $\varepsilon \rightarrow 0$ . Our main result (*cf.* Thm. 2.1 below) shows that the family  $(\mathcal{G}_\varepsilon)_\varepsilon$  converges, in the sense of  $\Gamma$ -convergence to an energy functional defined on the curve  $\gamma$ , which strongly depends on  $K$  and the target space  $\mathbb{S}^2$ , and has remarkable physical implications for relevant systems, for instance, to magnetic materials: in the curved thin-wire regime, generic exchange interactions (symmetric and antisymmetric) can manifest themselves under an additional anisotropy term whose specific shape depends both on the shape of the wire and the curvature of the target surface.

To properly state our result, we must finish setting up the stage. For every  $0 < \varepsilon < \delta$  we introduce the following functional defined on  $H^1(\mathcal{C}, \mathbb{S}^2)$ , which can be thought of as the pull-back of  $\mathcal{G}_\varepsilon$  on the cylindrical domain  $\mathcal{C} := I \times Q$ :

$$\begin{aligned} \mathcal{E}_\mathcal{C}^\varepsilon(\mathbf{v}_\varepsilon) := & \frac{1}{2|Q|} \int_{I \times Q} |\partial_s \mathbf{v}_\varepsilon + \tau(z \wedge \nabla_z \mathbf{v}_\varepsilon) + \alpha_\varepsilon K(\mathbf{v}_\varepsilon) \mathbf{t}|^2 \frac{1}{\alpha_\varepsilon} dz ds \\ & + \frac{1}{2|Q|} \int_{I \times Q} (|\varepsilon^{-1} \partial_1 \mathbf{v}_\varepsilon + \alpha_\varepsilon K(\mathbf{v}_\varepsilon) \mathbf{n}|^2 + |\varepsilon^{-1} \partial_2 \mathbf{v}_\varepsilon + \alpha_\varepsilon K(\mathbf{v}_\varepsilon) \mathbf{b}|^2) \frac{1}{\alpha_\varepsilon} dz ds \end{aligned} \quad (2.5)$$

where we set

$$\partial_1 := \partial_{z_1}, \partial_2 := \partial_{z_2}, \quad z \wedge \nabla_z \mathbf{v}_\varepsilon := z_2 \partial_1 \mathbf{v}_\varepsilon - z_1 \partial_2 \mathbf{v}_\varepsilon, \quad \alpha_\varepsilon(s, z) := (1 - \varepsilon \varpi(s) z_1), \quad (2.6)$$

with  $\varpi(s)$  being the curvature of the space curve  $\gamma$  at  $\gamma(s)$ .

Our main result is stated in the following statement.

**Theorem 2.1.** *For any  $0 < \varepsilon < \delta$ , the minimization problem for  $\mathcal{G}_\varepsilon$  in  $H^1(\Omega_\varepsilon, \mathbb{S}^2)$  is equivalent to the minimization in  $H^1(\mathcal{C}, \mathbb{S}^2)$  of the functional  $\mathcal{E}_\mathcal{C}^\varepsilon$  defined by (2.5) in the sense that a configuration  $\mathbf{v}_\varepsilon \in H^1(\Omega_\varepsilon, \mathbb{S}^2)$  minimizes  $\mathcal{G}_\varepsilon$  if and only if  $\mathbf{v}_\varepsilon \circ \varphi_\varepsilon \in H^1(\mathcal{C}, \mathbb{S}^2)$  minimizes  $\mathcal{E}_\mathcal{C}^\varepsilon$ .*

*The family  $(\mathcal{E}_\mathcal{C}^\varepsilon)_\varepsilon$  is equicoercive in the weak topology of  $H^1(\mathcal{C}, \mathbb{S}^2)$  and the  $\Gamma$ -limit  $\mathcal{E}_\mathcal{C} := \Gamma\text{-}\lim_{\varepsilon \rightarrow 0} \mathcal{E}_\mathcal{C}^\varepsilon$  is defined for every  $\mathbf{v} \in H^1(\mathcal{C}, \mathbb{S}^2)$  by*

$$\mathcal{E}_\mathcal{C}(\mathbf{v}) = \begin{cases} \frac{1}{2} \int_I |\partial_s \mathbf{v}(s) + K(\mathbf{v}(s)) \mathbf{t}(s)|^2 ds \\ \quad + \frac{1}{2} \int_I |K^\top(\mathbf{v}(s)) \mathbf{v}(s)|^2 - (K^\top(\mathbf{v}(s)) \mathbf{v}(s) \cdot \mathbf{t}(s))^2 ds & \text{if } \nabla_z \mathbf{v} = 0, \\ +\infty & \text{otherwise.} \end{cases} \quad (2.7)$$

Moreover,

$$\min_{H^1(\Omega_\varepsilon, \mathbb{S}^2)} \mathcal{G}_\varepsilon = \min_{H^1(\mathcal{C}, \mathbb{S}^2)} \mathcal{E}_\mathcal{C}^\varepsilon = \min_{H^1(\mathcal{C}, \mathbb{S}^2)} \mathcal{E}_\mathcal{C} + o(1) \quad (2.8)$$

and if  $(\mathbf{v}_\varepsilon)_\varepsilon$  is a minimizing family for  $(\mathcal{E}_\mathcal{C}^\varepsilon)_\varepsilon$  then  $(\mathbf{v}_\varepsilon)_\varepsilon$  converges, strongly in  $H^1(\mathcal{C}, \mathbb{S}^2)$ , to a minimum point of  $\mathcal{E}_\mathcal{C}$ .

**Remark 2.2.** In terms of normal and binormal vectors to the curve, the expression in (2.7) is equivalent to

$$\frac{1}{2} \int_I |\partial_s \mathbf{v}(s) + K(\mathbf{v}(s)) \mathbf{t}(s)|^2 ds + \frac{1}{2} \int_I (K^\top(\mathbf{v}(s)) \mathbf{v}(s) \cdot \mathbf{n}(s))^2 + (K^\top(\mathbf{v}(s)) \mathbf{v}(s) \cdot \mathbf{b}(s))^2 ds.$$

*Proof.* (EQUIVALENCE) First, we show that for any  $0 < \varepsilon < \delta$ , the minimization problem for  $\mathcal{G}_\varepsilon$  in  $H^1(\Omega_\varepsilon, \mathbb{S}^2)$  is equivalent to the minimization in  $H^1(\mathcal{C}, \mathbb{S}^2)$  of the functional  $\mathcal{E}_\mathcal{C}^\varepsilon$  defined by (2.5). To proceed, let  $F := (\mathbf{t} | \mathbf{n} | \mathbf{b})$  denote the matrix whose columns are the components of the Frenet frame. Using the Frenet–Serret formulas, with  $\varpi$  and  $\tau$  denoting the curvature and the torsion of  $\gamma$ , respectively:

$$\dot{\mathbf{n}}(s) = -\varpi \mathbf{t} + \tau \mathbf{b}, \quad \dot{\mathbf{t}} = \varpi \mathbf{n}, \quad \dot{\mathbf{b}} = -\tau \mathbf{n},$$

we obtain that for every  $0 < \varepsilon < \delta$ , it holds that

$$D\varphi_\varepsilon = \left( \partial_s \varphi_\varepsilon \mid \partial_1 \varphi_\varepsilon \mid \partial_2 \varphi_\varepsilon \right) = F \cdot \begin{pmatrix} 1 - \varepsilon \varpi z_1 & 0 & 0 \\ -\varepsilon \tau z_2 & \varepsilon & 0 \\ \varepsilon \tau z_1 & 0 & \varepsilon \end{pmatrix}.$$

The determinant of  $D\varphi_\varepsilon$  is then  $\det D\varphi_\varepsilon = \varepsilon^2 \alpha_\varepsilon$ , with  $\alpha_\varepsilon(s, z_1) := 1 - \varepsilon \varpi(s) z_1$ . Therefore,  $D\varphi_\varepsilon$  remains invertible for small  $\varepsilon$ . The inverse of  $D\varphi_\varepsilon$  is given by

$$(D\varphi_\varepsilon)^{-1} = \frac{1}{\alpha_\varepsilon} \Phi_\varepsilon F^\top, \quad \Phi_\varepsilon := \begin{pmatrix} 1 & 0 & 0 \\ \tau z_2 & \varepsilon^{-1} & 0 \\ -\tau z_1 & 0 & \varepsilon^{-1} \end{pmatrix}.$$

The previous considerations allow showing the equivalence between the minimization problem for  $\mathcal{G}_\varepsilon$  and  $\mathcal{E}_\mathcal{C}^\varepsilon$ . Indeed with  $\tilde{\mathbf{v}}_\varepsilon := \mathbf{v}_\varepsilon \circ \varphi_\varepsilon$ , we have

$$\begin{aligned} \mathcal{G}_\varepsilon(\mathbf{v}_\varepsilon) &= \frac{1}{2\varepsilon^2|Q|} \int_{\Omega_\varepsilon} |D\mathbf{v}_\varepsilon(x) + K(\mathbf{v}_\varepsilon(x))|^2 dx \\ &= \frac{1}{2} \int_{I \times Q} |D\tilde{\mathbf{v}}_\varepsilon(D\varphi_\varepsilon)^{-1} + K(\tilde{\mathbf{v}}_\varepsilon)|^2 \alpha_\varepsilon dz ds \\ &= \frac{1}{2|Q|} \int_{I \times Q} |(\partial_s \tilde{\mathbf{v}}_\varepsilon + \tau(z_2 \partial_1 \tilde{\mathbf{v}}_\varepsilon - z_1 \partial_2 \tilde{\mathbf{v}}_\varepsilon)) + \alpha_\varepsilon K(\tilde{\mathbf{v}}_\varepsilon) \mathbf{t}|^2 \frac{1}{\alpha_\varepsilon} dz ds \\ &\quad + \frac{1}{2|Q|} \int_{I \times Q} (|\varepsilon^{-1} \partial_1 \tilde{\mathbf{v}}_\varepsilon + \alpha_\varepsilon K(\tilde{\mathbf{v}}_\varepsilon) \mathbf{n}|^2 + |\varepsilon^{-1} \partial_2 \tilde{\mathbf{v}}_\varepsilon + \alpha_\varepsilon K(\tilde{\mathbf{v}}_\varepsilon) \mathbf{b}|^2) \frac{1}{\alpha_\varepsilon} dz ds \\ &= \mathcal{E}_\mathcal{C}^\varepsilon(\tilde{\mathbf{v}}_\varepsilon), \end{aligned}$$

from which the equivalence follows.

(COMPACTNESS/EQUICOERCIVENESS) We want to show that the family  $(\mathcal{E}_\mathcal{C}^\varepsilon)_\varepsilon$  is equicoercive in the weak topology of  $H^1(\mathcal{C}, \mathbb{S}^2)$ , *i.e.*, the existence of a weakly compact set  $J(\mathcal{C}, \mathbb{S}^2) \subseteq H^1(\mathcal{C}, \mathbb{S}^2)$  such that for every  $0 < \varepsilon < \delta$  there holds

$$\min_{H^1(\mathcal{C}, \mathbb{S}^2)} \mathcal{E}_\mathcal{C}^\varepsilon = \min_{J(\mathcal{C}, \mathbb{S}^2)} \mathcal{E}_\mathcal{C}^\varepsilon. \quad (2.9)$$

Ensuring the equicoercivity of  $(\mathcal{E}_\mathcal{C}^\varepsilon)_\varepsilon$  will guarantee that the fundamental theorem of  $\Gamma$ -convergence hypotheses are met. Consequently, the  $\Gamma$ -limit  $\mathcal{E}_\mathcal{C}$  will satisfy relation (2.8) on the variational convergence of minimum problems.

To show (2.9), we first observe that for any constant in space configuration  $\xi \in H^1(\mathcal{C}, \mathbb{S}^2)$  we have

$$\min_{\mathbf{v} \in H^1(\mathcal{C}, \mathbb{S}^2)} \mathcal{E}_\mathcal{C}^\varepsilon(\mathbf{v}) \leq \mathcal{E}_\mathcal{C}^\varepsilon(\xi) = \frac{1}{2|Q|} \int_{I \times Q} \frac{|K(\xi)|^2}{\sqrt{\alpha_\varepsilon(s, z)}} dz ds. \quad (2.10)$$

Given that for  $\varepsilon$  sufficiently small  $\alpha_\varepsilon(s, z)$  is uniformly bounded away from zero, there exists  $c_\varpi > 0$  such that (*cf.* the expression of  $\alpha_\varepsilon$  in (2.6))

$$c_\varpi^{-1} < \alpha_\varepsilon(s, z) < c_\varpi \quad \text{for every } (s, z) \in I \times Q. \quad (2.11)$$

Also, as stated in (2.4),  $|K(\sigma)| \leq c_K$ . Therefore, for  $\delta$  sufficiently small, the minimizers of  $(\mathcal{E}_\mathcal{C}^\varepsilon)_{0 < \varepsilon < \delta}$  are all contained in the  $\varepsilon$ -independent set

$$J(\mathcal{C}, \mathbb{S}^2) := \bigcup_{0 < \varepsilon < \delta} \{\mathbf{v} \in H^1(\mathcal{C}, \mathbb{S}^2) : \mathcal{E}_\mathcal{C}^\varepsilon(\mathbf{v}) \leq c\}, \quad (2.12)$$

where  $c > 0$  is a positive constant that depends only on  $c_\varpi, c_K$  and on the geometry of  $\gamma$ . After that, we use that for every  $a, b \in \mathbb{R}$  and every  $\eta > 0$  there holds  $|a + b|^2 \geq (1 + \eta)^{-1}|a|^2 - \eta^{-1}|b|^2$ , to infer that

$$\begin{aligned} 2|Q|\mathcal{E}_\mathcal{C}^\varepsilon(\mathbf{v}) &\geq \frac{1}{c_\varpi(1 + \eta)} \int_{I \times Q} |\partial_s \mathbf{v} + \tau(z \wedge \nabla_z \mathbf{v})|^2 \\ &\quad + \frac{1}{\alpha_\varepsilon} dz ds + \frac{1}{\varepsilon^2(1 + \eta)} \int_{I \times Q} |\nabla_z \mathbf{v}|^2 \frac{1}{\alpha_\varepsilon} dz ds - \frac{1}{\eta} \int_{I \times Q} \frac{|K(\mathbf{v})|^2}{\sqrt{\alpha_\varepsilon}} dz ds \end{aligned}$$

$$\begin{aligned} &\geq \frac{1}{c_\varpi(1+\eta)^2} \int_{I \times Q} |\partial_s \mathbf{v}|^2 dz ds - \frac{1}{c_\varpi \eta (1+\eta)} \tau^2 \int_{I \times Q} |z \wedge \nabla_z \mathbf{v}|^2 dz ds \\ &\quad + \frac{1}{c_\varpi(1+\eta)\varepsilon^2} \int_{I \times Q} |\nabla_z \mathbf{v}|^2 \frac{1}{\alpha_\varepsilon} dz ds - \frac{c_K^2}{\eta \sqrt{c_\varpi}} |I \times Q|. \end{aligned}$$

Given that  $|z \wedge \nabla_z \mathbf{v}| \leq 2|z| \cdot |\nabla_z \mathbf{v}| \leq c_Q |\nabla_z \mathbf{v}|$  for a positive constant  $c_Q > 0$  that depends only on  $Q$ , we get that up to the constant term  $-\frac{c_K^2}{\eta \sqrt{c_\varpi}} |I \times Q|$  there holds

$$2|Q| \mathcal{E}_C^\varepsilon(\mathbf{v}) \geq \frac{1}{c_\varpi(1+\eta)^2} \int_{I \times Q} |\partial_s \mathbf{v}|^2 dz ds + \frac{1}{(1+\eta)c_\varpi} \left( \frac{1}{\varepsilon^2} - \frac{\tau^2 c_Q^2}{\eta} \right) \int_{I \times Q} |\nabla_z \mathbf{v}|^2 dz ds.$$

In particular, choosing  $\eta := 2\varepsilon^2 \tau^2 c_Q^2$ , we get the existence of a constant  $c_* > 0$  dependent only on  $\delta, c_\varpi, c_K$  and the shape of the  $\delta$ -tube generated by  $\gamma$ , such that

$$\int_{I \times Q} |\partial_s \mathbf{v}|^2 dz ds + \frac{1}{\varepsilon^2} \int_{I \times Q} |\nabla_z \mathbf{v}_\varepsilon|^2 dz ds \leq c_* (1 + \mathcal{E}_C^\varepsilon(\mathbf{v})). \quad (2.13)$$

Assuming without loss of generality that  $\delta < 1$  so that also  $\varepsilon < 1$ , from the previous relation (2.13) we can conclude that the set  $J(\mathcal{C}, \mathbb{S}^2)$  defined by (2.12) is contained in the bounded subset of  $H^1(\mathcal{C}, \mathbb{R}^3)$  given by

$$H_b^1(\mathcal{C}, \mathbb{S}^2) := \{\mathbf{v} \in H^1(\mathcal{C}, \mathbb{R}^3) : |\mathbf{v}| = 1, \|\mathbf{v}\|_{H^1(\mathcal{C}, \mathbb{R}^3)}^2 \leq c_*(1+c)\}$$

Therefore

$$\min_{\mathbf{v} \in H^1(\mathcal{C}, \mathbb{S}^2)} \mathcal{E}_C^\varepsilon(\mathbf{v}) = \min_{\mathbf{v} \in H_b^1(\mathcal{C}, \mathbb{S}^2)} \mathcal{E}_C^\varepsilon(\mathbf{v}).$$

To establish that  $H_b^1(\mathcal{C}, \mathbb{S}^2)$  is weakly compact, we need to prove that it is weakly closed. To do this, consider a sequence  $(\mathbf{v}_n)_{n \in \mathbb{N}}$  in  $H_b^1(\mathcal{C}, \mathbb{S}^2)$  such that  $\mathbf{v}_n \rightharpoonup \mathbf{v}_0$  weakly in  $H^1(\mathcal{C}, \mathbb{R}^3)$ ; by Rellich–Kondrachov theorem,  $\mathbf{v}_n \rightarrow \mathbf{v}_0$  strongly in  $L^2(\mathcal{C}, \mathbb{R}^3)$ . Thus, after extracting a subsequence that converges pointwise a.e. in  $\mathcal{C}$ , we get that the  $\mathbf{v}_0$  still takes values in  $\mathbb{S}^2$ . This completes the proof of equicoerciveness.

( $\Gamma$ -LIMINF) Let  $(\mathbf{v}_\varepsilon)$  be a family in  $H^1(\mathcal{C}, \mathbb{S}^2)$  such that

$$\sup_{0 < \varepsilon < \delta} \mathcal{E}_C^\varepsilon(\mathbf{v}_\varepsilon) < +\infty.$$

By estimate (2.13), we get that the families  $(\varepsilon^{-1} \partial_1 \mathbf{v}_\varepsilon)$ ,  $(\varepsilon^{-1} \partial_2 \mathbf{v}_\varepsilon)$  and  $(\partial_s \mathbf{v}_\varepsilon)$  are bounded in  $L^2$ . Therefore, there exist  $\mathbf{v}_0 \in H^1(\mathcal{C}, \mathbb{S}^2)$ ,  $\mathbf{d}_1, \mathbf{d}_2 \in L^2(\mathcal{C}, \mathbb{R}^3)$ , such that

$$\begin{aligned} \mathbf{v}_\varepsilon &\rightarrow \mathbf{v}_0 && \text{strongly in } L^2(\mathcal{C}, \mathbb{S}^2), \\ \partial_s \mathbf{v}_\varepsilon &\rightharpoonup \partial_s \mathbf{v}_0 && \text{weakly in } L^2(\mathcal{C}, \mathbb{R}^3), \\ \frac{1}{\varepsilon} \partial_1 \mathbf{v}_\varepsilon &\rightharpoonup \mathbf{d}_1 && \text{weakly in } L^2(\mathcal{C}, \mathbb{R}^3), \\ \frac{1}{\varepsilon} \partial_2 \mathbf{v}_\varepsilon &\rightharpoonup \mathbf{d}_2 && \text{weakly in } L^2(\mathcal{C}, \mathbb{R}^3). \end{aligned} \quad (2.14)$$

The previous relations imply that  $\mathbf{v}_0$  is 0-homogeneous along the cross-section of the wire, *i.e.*, that it depends only on the  $s$ -variable. In particular,

$$\nabla_z \mathbf{v}_\varepsilon \rightarrow 0 \quad \text{strongly in } L^2(\mathcal{C}, \mathbb{R}^3). \quad (2.15)$$

Moreover, since  $\mathbf{v}_\varepsilon$  is constrained to take values in the unit sphere  $\mathbb{S}^2$ , one has that  $\partial_1 \mathbf{v}_\varepsilon \cdot \mathbf{v}_\varepsilon = \partial_2 \mathbf{v}_\varepsilon \cdot \mathbf{v}_\varepsilon = 0$  for every  $0 < \varepsilon < \delta$ . Hence, in the limit for  $\varepsilon \rightarrow 0$ , we still get that  $\mathbf{d}_1, \mathbf{d}_2$  are pointwise orthogonal to  $\mathbf{v}_0$ , *i.e.*,

$$\mathbf{d}_1 \cdot \mathbf{v}_0 = 0, \quad \mathbf{d}_2 \cdot \mathbf{v}_0 = 0. \quad (2.16)$$

By (2.14), (2.15), and (2.16), taking into account the lower semicontinuity of the norm and that  $\alpha_\varepsilon \rightarrow 1$  when  $\varepsilon \rightarrow 0$ , we get that

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} \mathcal{E}_C^\varepsilon(\mathbf{v}_\varepsilon) &\geq \frac{1}{2} \int_I |\partial_s \mathbf{v}_0 + K(\mathbf{v}_0) \mathbf{t}|^2 ds \\ &+ \frac{1}{2|Q|} \int_{I \times Q} \left( |\mathbf{d}_1 + K(\mathbf{v}_0) \mathbf{n}|^2 + |\mathbf{d}_2 + K(\mathbf{v}_0) \mathbf{b}|^2 \right) dz ds \\ &\geq \frac{1}{2} \int_I |\partial_s \mathbf{v}_0 + K(\mathbf{v}_0) \mathbf{t}|^2 ds + \frac{1}{2} \int_I \left[ \min_{\mathbf{d}_1(s), \mathbf{d}_2(s) \in \mathbf{v}_0^\perp(s)} \mathbf{g}(\mathbf{d}_1(s), \mathbf{d}_2(s)) \right] dz ds, \end{aligned}$$

with

$$\mathbf{g}(\mathbf{d}_1, \mathbf{d}_2) := |\mathbf{d}_1 + K(\mathbf{v}_0) \mathbf{n}|^2 + |\mathbf{d}_2 + K(\mathbf{v}_0) \mathbf{b}|^2. \quad (2.17)$$

The minimal energy of  $\mathbf{g}$  is given by

$$\min_{\mathbf{d}_1(s), \mathbf{d}_2(s) \in \mathbf{v}_0^\perp(s)} \mathbf{g}(\mathbf{d}_1(s), \mathbf{d}_2(s)) = (K(\mathbf{v}_0) \mathbf{n} \cdot \mathbf{v}_0)^2 + (K(\mathbf{v}_0) \mathbf{b} \cdot \mathbf{v}_0)^2, \quad (2.18)$$

and it is achieved when  $\mathbf{d}_1$  and  $\mathbf{d}_2$  are respectively given by

$$\mathbf{d}_1(\mathbf{v}_0) := \mathbf{v}_0 \times (\mathbf{v}_0 \times K(\mathbf{v}_0) \mathbf{n}), \quad (2.19)$$

$$\mathbf{d}_2(\mathbf{v}_0) := \mathbf{v}_0 \times (\mathbf{v}_0 \times K(\mathbf{v}_0) \mathbf{b}). \quad (2.20)$$

Indeed, one can observe that

$$\begin{aligned} \mathbf{g}(\mathbf{d}_1, \mathbf{d}_2) &= |\mathbf{d}_1 - \mathbf{v}_0 \times (\mathbf{v}_0 \times K(\mathbf{v}_0) \mathbf{n}) + (K(\mathbf{v}_0) \mathbf{n} \cdot \mathbf{v}_0) \mathbf{v}_0|^2 \\ &\quad + |\mathbf{d}_2 - \mathbf{v}_0 \times (\mathbf{v}_0 \times K(\mathbf{v}_0) \mathbf{b}) + (K(\mathbf{v}_0) \mathbf{b} \cdot \mathbf{v}_0) \mathbf{v}_0|^2 \\ &= |\mathbf{d}_1 - \mathbf{v}_0 \times (\mathbf{v}_0 \times K(\mathbf{v}_0) \mathbf{n})|^2 + |\mathbf{d}_2 - \mathbf{v}_0 \times (\mathbf{v}_0 \times K(\mathbf{v}_0) \mathbf{b})|^2 \\ &\quad + (K(\mathbf{v}_0) \mathbf{n} \cdot \mathbf{v}_0)^2 + (K(\mathbf{v}_0) \mathbf{b} \cdot \mathbf{v}_0)^2. \end{aligned}$$

Summarizing, we obtained that

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} \mathcal{E}_C^\varepsilon(\mathbf{v}_\varepsilon) &\geq \frac{1}{2} \int_I |\partial_s \mathbf{v}(s) + K(\mathbf{v}(s)) \mathbf{t}(s)|^2 ds \\ &\quad + \frac{1}{2} \int_I (K^\top(\mathbf{v}(s)) \mathbf{v}(s) \cdot \mathbf{n}(s))^2 + (K^\top(\mathbf{v}(s)) \mathbf{v}(s) \cdot \mathbf{b}(s))^2 ds, \end{aligned} \quad (2.21)$$

and this is nothing but the right-hand side of (2.7).

( $\Gamma$ -LIMSUP) A direct computation shows that for any  $\mathbf{v}_0 \in H^1(\mathcal{C}, \mathbb{S}^2)$  such that  $\nabla_z \mathbf{v}_0 = 0$ , the family

$$\mathbf{v}_\varepsilon := \frac{\mathbf{v}_0 + \varepsilon \mathbf{d}(\mathbf{v}_0)}{|\mathbf{v}_0 + \varepsilon \mathbf{d}(\mathbf{v}_0)|}, \quad \mathbf{d}(\mathbf{v}_0) := z_1 \mathbf{d}_1(\mathbf{v}_0) + z_2 \mathbf{d}_2(\mathbf{v}_0), \quad (2.22)$$

with  $\mathbf{d}_1(\mathbf{v}_0)$ ,  $\mathbf{d}_2(\mathbf{v}_0)$  given by (2.19)–(2.20), is a recovery sequence, *i.e.*,

$$\mathbf{v}_\varepsilon \rightarrow \mathbf{v}_0 \text{ strongly in } H^1(\mathcal{C}, \mathbb{S}^2) \quad \text{and} \quad \mathcal{E}_\mathcal{C}^\varepsilon(\mathbf{v}_\varepsilon) \rightarrow \mathcal{E}_\mathcal{C}(\mathbf{v}_0). \quad (2.23)$$

Indeed, since  $\mathbf{d}_1(\mathbf{v}_0)$ ,  $\mathbf{d}_2(\mathbf{v}_0)$  depend only on the  $s$ -variable and  $|\mathbf{v}_0| = 1$ , we have

$$\begin{aligned} D\mathbf{v}_\varepsilon &= \frac{1}{|\mathbf{v}_0 + \varepsilon\mathbf{d}(\mathbf{v}_0)|} \left( I + \frac{\mathbf{v}_0 + \varepsilon\mathbf{d}(\mathbf{v}_0)}{|\mathbf{v}_0 + \varepsilon\mathbf{d}(\mathbf{v}_0)|} \otimes \frac{\mathbf{v}_0 + \varepsilon\mathbf{d}(\mathbf{v}_0)}{|\mathbf{v}_0 + \varepsilon\mathbf{d}(\mathbf{v}_0)|} \right) D(\mathbf{v}_0 + \varepsilon\mathbf{d}(\mathbf{v}_0)) \\ &\approx (I + \mathbf{v}_0 \otimes \mathbf{v}_0)(\partial_s|\partial_1|\partial_2)(\mathbf{v}_0 + \varepsilon\mathbf{d}(\mathbf{v}_0)) \\ &= (I + \mathbf{v}_0 \otimes \mathbf{v}_0)(\partial_s\mathbf{v}_0 + \varepsilon\partial_s\mathbf{d}(\mathbf{v}_0)|\varepsilon\mathbf{d}_1(\mathbf{v}_0)|\varepsilon\mathbf{d}_2(\mathbf{v}_0)) \end{aligned}$$

where we used the approximation symbol to denote equality up to negligible terms when  $\varepsilon \rightarrow 0$ . Given that  $0 = \partial_s\mathbf{v}_0 \cdot \mathbf{v}_0 = \mathbf{d}_1(\mathbf{v}_0) \cdot \mathbf{v}_0 = \mathbf{d}_2(\mathbf{v}_0) \cdot \mathbf{v}_0$ , we infer the following limit relations

$$\partial_s\mathbf{v}_\varepsilon \rightarrow (I + \mathbf{v}_0 \otimes \mathbf{v}_0)\partial_s\mathbf{v}_0 = \partial_s\mathbf{v}_0, \quad (2.24)$$

$$\varepsilon^{-1}\partial_1\mathbf{v}_\varepsilon \rightarrow (I + \mathbf{v}_0 \otimes \mathbf{v}_0)\mathbf{d}_1(\mathbf{v}_0) = \mathbf{d}_1(\mathbf{v}_0), \quad (2.25)$$

$$\varepsilon^{-1}\partial_2\mathbf{v}_\varepsilon \rightarrow (I + \mathbf{v}_0 \otimes \mathbf{v}_0)\mathbf{d}_2(\mathbf{v}_0) = \mathbf{d}_2(\mathbf{v}_0), \quad (2.26)$$

from which (2.23) follows.

(STRONG CONVERGENCE OF MINIMIZERS) Let  $(\mathbf{v}_\varepsilon)_\varepsilon \subset H^1(\mathcal{C}, \mathbb{S}^2)$  be a minimizing family for  $(\mathcal{E}_\mathcal{C}^\varepsilon)_\varepsilon$ . By compactness (*cf.* (2.14)–(2.15)), there exists  $\mathbf{v}_0 \in H^1(\mathcal{C}, \mathbb{S}^2)$ , depending only the  $s$ -variable, such that  $\mathbf{v}_\varepsilon \rightharpoonup \mathbf{v}_0$  weakly in  $H^1(\mathcal{C}, \mathbb{S}^2)$  and  $\partial_s\mathbf{v}_\varepsilon \rightarrow 0$  strongly in  $L^2(\mathcal{C}, \mathbb{S}^2)$ . By the  $\Gamma$ -liminf inequality, the strong convergence of  $\mathbf{v}_\varepsilon \rightarrow \mathbf{v}_0$  in  $L^2(\mathcal{C}, \mathbb{S}^2)$ , and the minimality of  $\mathbf{v}_\varepsilon$ , we get that if  $\mathbf{v}_\varepsilon^*$  is the recovery sequence built from  $\mathbf{v}_0$  as in (2.22), then

$$\mathcal{E}_\mathcal{C}(\mathbf{v}_0) \leq \liminf_{\varepsilon \rightarrow 0} \mathcal{E}_\mathcal{C}^\varepsilon(\mathbf{v}_\varepsilon) \leq \limsup_{\varepsilon \rightarrow 0} \mathcal{E}_\mathcal{C}^\varepsilon(\mathbf{v}_\varepsilon) \leq \lim_{\varepsilon \rightarrow 0} \mathcal{E}_\mathcal{C}^\varepsilon(\mathbf{v}_\varepsilon^*) = \mathcal{E}_\mathcal{C}(\mathbf{v}_0). \quad (2.27)$$

Therefore, if  $(\mathbf{v}_\varepsilon)_\varepsilon$  is a minimizing family for  $(\mathcal{E}_\mathcal{C}^\varepsilon)_\varepsilon$ , then  $\mathbf{v}_\varepsilon \rightharpoonup \mathbf{v}_0$  weakly in  $H^1(\mathcal{C}, \mathbb{S}^2)$  for some  $\mathbf{v}_0$  which depends only on the  $s$ -variable and, moreover, by (2.27),  $\lim_{\varepsilon \rightarrow 0} \mathcal{E}_\mathcal{C}^\varepsilon(\mathbf{v}_\varepsilon) = \mathcal{E}_\mathcal{C}(\mathbf{v}_0)$ . By the compactness relations (2.14)–(2.15) and the lower semicontinuity of the norm, we get that

$$\begin{aligned} \mathcal{E}_\mathcal{C}(\mathbf{v}_0) &= \lim_{\varepsilon \rightarrow 0} \mathcal{E}_\mathcal{C}^\varepsilon(\mathbf{v}_\varepsilon) \geq \limsup_{\varepsilon \rightarrow 0} \frac{1}{2|Q|} \int_{I \times Q} |\partial_s\mathbf{v}_\varepsilon + K(\mathbf{v}_\varepsilon)\mathbf{t}|^2 dz ds \\ &\quad + \frac{1}{2|Q|} \int_{I \times Q} \mathbf{g}(\mathbf{d}_1, \mathbf{d}_2) dz ds \\ &\geq \limsup_{\varepsilon \rightarrow 0} \frac{1}{2|Q|} \int_{I \times Q} |\partial_s\mathbf{v}_\varepsilon + K(\mathbf{v}_\varepsilon)\mathbf{t}|^2 dz ds \\ &\quad + \frac{1}{2} \int_I (K^\top(\mathbf{v}_0)\mathbf{v}_0 \cdot \mathbf{n})^2 + (K^\top(\mathbf{v}_0)\mathbf{v}_0 \cdot \mathbf{b})^2 ds, \end{aligned} \quad (2.28)$$

where, for the last inequality, we used the fact that the function  $\mathbf{g}$  defined by (2.17) has minimal energy  $(K(\mathbf{v}_0)\mathbf{n} \cdot \mathbf{v}_0)^2 + (K(\mathbf{v}_0)\mathbf{b} \cdot \mathbf{v}_0)^2$ . Substituting in (2.28) the equivalent expression of  $\mathcal{E}_\mathcal{C}(\mathbf{v}_0)$  given in Remark 2.2, we get that

$$\limsup_{\varepsilon \rightarrow 0} \frac{1}{2|Q|} \int_{I \times Q} |\partial_s\mathbf{v}_\varepsilon + K(\mathbf{v}_\varepsilon)\mathbf{t}|^2 dz ds \leq \frac{1}{2} \int_I |\partial_s\mathbf{v}_0 + K(\mathbf{v}_0)\mathbf{t}|^2 ds. \quad (2.29)$$



Hence, from the lower semicontinuity of the norm, it follows that

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{2|Q|} \int_{I \times Q} |\partial_s \mathbf{v}_\varepsilon + K(\mathbf{v}_\varepsilon) \mathbf{t}|^2 dz ds = \frac{1}{2} \int_I |\partial_s \mathbf{v}_0 + K(\mathbf{v}_0) \mathbf{t}|^2 ds. \quad (2.30)$$

The previous relation implies that  $\|\partial_s \mathbf{v}_\varepsilon\|_{L^2(\mathcal{C}, \mathbb{S}^2)} \rightarrow \|\partial_s \mathbf{v}_0\|_{L^2(\mathcal{C}, \mathbb{S}^2)}$ . Overall, also considered that  $\|\partial_1 \mathbf{v}_\varepsilon\|_{L^2(\mathcal{C}, \mathbb{S}^2)} \rightarrow 0$  and  $\|\partial_2 \mathbf{v}_\varepsilon\|_{L^2(\mathcal{C}, \mathbb{S}^2)} \rightarrow 0$ , we deduce the convergence of the norms  $\|\mathbf{v}_\varepsilon\|_{H^1(\mathcal{C}, \mathbb{S}^2)} \rightarrow \|\mathbf{v}_0\|_{H^1(\mathcal{C}, \mathbb{S}^2)}$  which together with the weak convergence  $\mathbf{v}_\varepsilon \rightharpoonup \mathbf{v}_0$  in  $H^1(\mathcal{C}, \mathbb{S}^2)$ , assures strong convergence in  $H^1(\mathcal{C}, \mathbb{S}^2)$ . This concludes the proof.  $\square$

### 3. APPLICATIONS TO MICROMAGNETICS

In this section, we illustrate how our analysis of curved thin wires can explain different scenarios that may occur when ferromagnetic crystals lack inversion symmetry and how the so-called Dzyaloshinskii–Moriya interaction (DMI) can distort the otherwise uniform ferromagnetic state. Before delving into specifics, we describe the physical framework and introduce relevant notation.

In the variational theory of micromagnetism (*cf.* [44–46]), the order parameter that characterizes the orientations of a rigid ferromagnetic body within a bounded Lipschitz domain  $\Omega \subseteq \mathbb{R}^3$  is the magnetization  $\mathbf{M}$ , a vector field whose magnitude  $M_s := |\mathbf{M}|$ , called the saturation magnetization, can be considered constant in  $\Omega$  at temperatures well below the Curie temperature. Hence, one can represent the magnetization as  $\mathbf{M} := M_s \mathbf{v}$ , where  $\mathbf{v} : \Omega \rightarrow \mathbb{S}^2$  is a map taking values into the unit sphere of  $\mathbb{R}^3$ . Even though the magnitude of  $\mathbf{v}$  remains constant in space, its direction is free to vary and the observable magnetization states can be expressed as local minimizers of the micromagnetic energy functional. In single-crystal ferromagnets (*cf.* [47, 48]), and upon appropriate non-dimensionalization, the relevant energy functional takes the form

$$\mathcal{E}_\Omega(\mathbf{v}) + \mathcal{W}_\Omega(\mathbf{v}) := \frac{1}{2} \int_\Omega |\nabla \mathbf{v}|^2 dx + \frac{1}{2} \int_\Omega |\nabla u|^2 dx \quad (3.1)$$

where  $\mathbf{v} \in H^1(\Omega, \mathbb{S}^2)$  represents the normalized magnetization field, and  $u$  the unique solution in  $H^1(\mathbb{R}^3)$  of the Poisson equation

$$\Delta u = \nabla \cdot (\mathbf{v} \chi_{\Omega_\varepsilon}), \quad (3.2)$$

with  $\mathbf{v} \chi_{\Omega_\varepsilon}$  denoting the extension by zero of  $\mathbf{v}$  to  $\mathbb{R}^3$ . We refer the reader to [41] for further mathematical details on this formulation. Physically, the exchange energy  $\mathcal{E}_\Omega$  reflects the effects of symmetric exchange interactions that penalize spatial variations in the orientation of the magnetization, thereby encouraging collinear alignment. The term  $\mathcal{W}_\Omega$  represents the magnetostatic self-energy, corresponding to the energy associated with the magnetic scalar potential  $u$  induced by the normalized magnetization  $\mathbf{v}$ .

In this work, we extend the analysis beyond traditional *symmetric* exchange interactions by investigating the effects of potential centrosymmetry-breaking elements within the crystal lattice structure of the ferromagnetic material (see [42, 49] for an in-depth treatment of symmetric and antisymmetric exchange interactions). Specifically, we examine the additional contributions to the energy landscape introduced by this structural asymmetry. Alongside the energy density defined in equation (3.1), we incorporate *antisymmetric* exchange interactions by accounting for terms that stem from antisymmetric mathematical structures, particularly the chirality tensor  $\nabla \times \mathbf{v}$ , a form analyzed in Lifshitz’s theory of phase transitions, often termed as Lifshitz invariants [50].

A focal point in our study is the energy contribution associated with the so-called *bulk* Dzyaloshinskii–Moriya interaction (DMI), expressed by the energy density  $\mathbf{v} \cdot (\nabla \times \mathbf{v})$ . This term represents the trace of the chirality tensor and encapsulates essential features of the material’s antisymmetric exchange interactions, potentially

leading to novel magnetic phenomena. Specifically, for every  $\mathbf{v} \in H^1(\Omega, \mathbb{S}^2)$ , we define the bulk DMI energy as

$$\mathcal{H}_\Omega(\mathbf{v}) := \kappa \int_\Omega \mathbf{v} \cdot (\nabla \times \mathbf{v}) \, dx. \quad (3.3)$$

The normalized constant  $\kappa \in \mathbb{R}$  is the so-called DMI constant, and its sign determines the chirality of the system.

The full micromagnetic energy functional we are interested in is then  $\mathcal{E}_\Omega + \mathcal{H}_\Omega + \mathcal{W}_\Omega$ . In particular, with the notation introduced in Section 2, the micromagnetic energy associated with an  $\varepsilon$ -tube  $\Omega_\varepsilon$  along a curve  $\gamma : I \mapsto \mathbb{R}^3$  with cross-sectional shape  $Q$  is given by

$$\mathcal{G}_\varepsilon(\mathbf{v}) := \frac{1}{\varepsilon^2|Q|} \left( \frac{1}{2} \int_{\Omega_\varepsilon} |D\mathbf{v}|^2 \, dx + \kappa \int_{\Omega_\varepsilon} \mathbf{v} \cdot \nabla \times \mathbf{v} \, dx + \frac{1}{2} \int_{\mathbb{R}^3} |\nabla u|^2 \, dx \right), \quad (3.4)$$

We note that in the energy above, we take a form of bulk DMI  $\nabla \times \mathbf{v} \cdot \mathbf{v}$ . To put it in the form (2.2), we define a matrix  $K$  as follows:

$$K(\mathbf{v}) = \kappa \begin{pmatrix} 0 & -v_3 & v_2 \\ v_3 & 0 & -v_1 \\ -v_2 & v_1 & 0 \end{pmatrix}.$$

It is clear that

$$\frac{1}{2} |D\mathbf{v} + K(\mathbf{v})|^2 = \frac{1}{2} |D\mathbf{v}|^2 + D\mathbf{v} : K(\mathbf{v}) + \frac{1}{2} |K(\mathbf{v})|^2,$$

and a simple computation shows that  $D\mathbf{v} : K(\mathbf{v}) = \kappa \mathbf{v} \cdot \nabla \times \mathbf{v}$ , and  $|K(\mathbf{v})|^2 = 2\kappa^2$ . Based on these relations, we can rewrite our energy as follows:

$$\mathcal{G}_\varepsilon(\mathbf{v}) := \frac{1}{\varepsilon^2|Q|} \left( \frac{1}{2} \int_{\Omega_\varepsilon} |D\mathbf{v} + K(\mathbf{v})|^2 \, dx + \frac{1}{2} \int_{\mathbb{R}^3} |\nabla u|^2 \, dx \right) - \kappa^2 \frac{|\Omega_\varepsilon|}{\varepsilon^2|Q|}.$$

Without loss of generality, we assume  $|Q| = 1$  and note that the last constant term in the energy can be dropped as it does not affect critical points or  $\Gamma$ -convergence. Then, using  $\Gamma$ -convergence results from [43] (see also [51]) and (2.7), in the limit as  $\varepsilon \rightarrow 0$  we obtain the following limiting energy

$$\mathcal{E}_C(\mathbf{v}) = \frac{1}{2} \int_I |\partial_s \mathbf{v}(s) + K(\mathbf{v}(s))\mathbf{t}(s)|^2 \, ds + \frac{1}{2} \int_I M\mathbf{v}(s) \cdot \mathbf{v}(s) \, ds, \quad (3.5)$$

where

$$M = -\frac{1}{2\pi} \int_{\partial Q} \int_{\partial Q} \mathbf{n}(\xi) \otimes \mathbf{n}(\eta) \ln |\xi - \eta| \, d\xi \, d\eta. \quad (3.6)$$

We now want to investigate several special cases of the limiting energy and find its minimizers in the form of domain walls.

### 3.1. Straight wire with DMI

In the case of an infinite straight wire with a circular cross-section, the interval  $I$  in (3.5) is not compact, and one cannot directly apply Theorem 2.1. However, it is still possible to prove the result for an infinite straight wire with some standard modifications to account for translation invariance. It is also possible to model this

scenario by considering it as the limit of a sequence of energy functionals defined on straight wires of finite lengths. We have  $\mathbf{t} = \mathbf{e}_1$  and, up to constant terms, the limiting energy can be rewritten as

$$\mathcal{E}_{\mathcal{C}}(\mathbf{v}) = \frac{1}{2} \int_{\mathbb{R}} |\partial_x \mathbf{v} + \kappa(0, v_3, -v_2)|^2 dx + \frac{1}{4\pi} \int_{\mathbb{R}} (1 - v_1^2) dx. \quad (3.7)$$

We note that the global minimum of the energy is achieved for  $\mathbf{v} = \pm \mathbf{e}_1$ .

We are interested in non-trivial local minimizers corresponding to domain walls. Therefore, we can impose boundary conditions  $\mathbf{v} \rightarrow \pm \mathbf{e}_1$  as  $x \rightarrow \pm\infty$  and look for a global minimizer of  $\mathcal{E}_{\mathcal{C}}$  with these boundary conditions. For that, we decompose the first term as

$$|\partial_x \mathbf{v} + \kappa(0, v_3, -v_2)|^2 = |v_1'|^2 + |v_2' + \kappa v_3|^2 + |v_3' - \kappa v_2|^2. \quad (3.8)$$

and represent the magnetization  $\mathbf{v}$  in spherical coordinates

$$\mathbf{v}(x) = (\cos \theta(x), \sin \theta(x) \cos \phi(x), \sin \theta(x) \sin \phi(x)).$$

Plugging it into the energy we obtain

$$\mathcal{E}_{\mathcal{C}}(\theta, \phi) = \frac{1}{2} \int_{\mathbb{R}} |\theta'(x)|^2 + \sin^2 \theta(x) |\phi'(x) - \kappa|^2 dx + \frac{1}{4\pi} \int_{\mathbb{R}} \sin^2 \theta(x) dx. \quad (3.9)$$

The boundary conditions translate into  $\theta(x) \rightarrow 0$  as  $x \rightarrow \infty$ ,  $\theta(x) \rightarrow k\pi$  as  $x \rightarrow -\infty$ , where  $k \in \mathbb{N}$  is odd. We minimize this energy in  $\phi$  to obtain  $\phi(x) = \kappa x$ . The subsequent minimization in  $\theta$  gives us  $k = 1$  and a standard domain wall profile

$$\theta_0(x) = 2 \arctan \left( e^{-\frac{1}{\sqrt{4\pi}} x} \right). \quad (3.10)$$

The minimizing profile is

$$\mathbf{v}_0(x) = (\cos \theta_0(x), \sin \theta_0(x) \cos(\kappa x), \sin \theta_0(x) \sin(\kappa x)). \quad (3.11)$$

It is clear that  $\mathbf{v}_0$  is a global minimizer of the energy (3.7) under domain wall boundary conditions  $\mathbf{v} \rightarrow \pm \mathbf{e}_1$  as  $x \rightarrow \pm\infty$ . Therefore, it follows that it is a local minimizer of  $\mathcal{E}_{\mathcal{C}}$  with respect to  $H^1$  perturbations.

We note that in the intuitive 1D energy for a nanowire sometimes studied in physics literature for soft materials [52]

$$\frac{1}{2} \int_{\mathbb{R}} |\partial_x \mathbf{v}|^2 + \kappa(v_2' v_3 - v_3' v_2) + Q(1 - v_1^2) dx \quad (3.12)$$

the DMI term might lead to an uncontrolled profile winding at the tails to decrease the micromagnetic energy. This phenomenon is always absent in the reduced 3D to 1D model due to the presence of two effective anisotropy terms produced by DMI and stray field energies. Effective anisotropy here globally suppresses conical modulation [53], but modulation occurs along the domain wall (see Fig. 2).

### 3.2. Rings with DMI

If our original domain has a geometry of a ring with radius  $R$  and circular cross-section, we can define  $\mathbf{t}(\phi) = (-\sin \phi, \cos \phi, 0)$ ,  $\mathbf{n}(\phi) = (\cos \phi, \sin \phi, 0)$  and hence

$$K(\mathbf{v}(\phi))\mathbf{t}(\phi) = \kappa(-v_3 \cos \phi, -v_3 \sin \phi, v_2 \sin \phi + v_1 \cos \phi) = -\kappa v_3 \mathbf{n}(\phi) + \kappa \mathbf{v}(\phi) \cdot \mathbf{n}(\phi) \mathbf{e}_3.$$




FIGURE 2. Sketch of the domain wall in the straight wire. The value of the dimensionless parameter characterizing the strength of the DMI is chosen to be  $\kappa = 0.36$ , which is close to that for the B20-type FeGe [18].

It is convenient to rescale the domain to be a unit circle, and hence, we can write the energy rescaled by  $R$  as

$$\mathcal{E}_C(\mathbf{v}) = \frac{1}{2R^2} \int_0^{2\pi} |\partial_\phi \mathbf{v}|^2 d\phi + \frac{\kappa}{R} \int_0^{2\pi} (-\partial_\phi \mathbf{v} \cdot \mathbf{n}v_3 + \mathbf{v} \cdot \mathbf{n}\partial_\phi v_3) d\phi \quad (3.13)$$

$$+ \frac{\kappa^2}{2} \int_0^{2\pi} 1 - (\mathbf{v} \cdot \mathbf{t})^2 d\phi + \frac{1}{4\pi} \int_0^{2\pi} 1 - (\mathbf{v} \cdot \mathbf{t})^2 d\phi. \quad (3.14)$$

When DMI can be neglected, an extensive examination of energy minimizers and the resulting symmetry-breaking phenomena is carried out in [54] (see also [55] for findings on dynamic stability).

Observe that the DMI term produces an additional anisotropy along the wire. It is convenient to introduce local ring coordinates and decompose the magnetization  $\mathbf{v}$  along the Frenet frame  $\mathbf{v} = v_t \mathbf{t} + v_n \mathbf{n} + v_3 \mathbf{b}$ . We now have  $\partial_\phi \mathbf{v} = (v'_t + v_n) \mathbf{t} + (v'_n - v_t) \mathbf{n} + v'_3 \mathbf{b}$  and, therefore, it is possible to rewrite the reduced 1D energy as

$$\begin{aligned} \mathcal{E}_C(\mathbf{v}) = \frac{1}{2R^2} \int_0^{2\pi} & |v'_t + v_n|^2 + |v'_n - v_t - R\kappa v_3|^2 + |v'_3 + R\kappa v_n|^2 d\phi \\ & + \frac{1}{4\pi} \int_0^{2\pi} (1 - v_t^2) d\phi \end{aligned} \quad (3.15)$$

Let us first try to understand the influence of the DMI in the model with only exchange and DMI present

$$\mathcal{E}_C^0(\mathbf{v}) = \frac{1}{2R^2} \int_0^{2\pi} |v'_t + v_n|^2 + |v'_n - v_t - R\kappa v_3|^2 + |v'_3 + R\kappa v_n|^2 d\phi.$$

It is clear that exchange energy wants magnetization to be a constant vector. However, the presence of DMI breaks the symmetry of the problem and leads to novel types of minimizers. Indeed, we can solve the system

$$v'_t + v_n = 0, \quad v'_n - v_t - R\kappa v_3 = 0, \quad v'_3 + R\kappa v_n = 0, \quad |\mathbf{v}| = 1, \quad (3.16)$$

and obtain the following set of solutions:

$$v_t = \frac{A}{\sqrt{1 + R^2 \kappa^2}} \cos(\sqrt{1 + R^2 \kappa^2} \phi + \phi_0) - R\kappa B, \quad (3.17)$$

$$v_n = A \sin(\sqrt{1 + R^2 \kappa^2} \phi + \phi_0), \quad (3.18)$$

$$v_3 = \frac{AR\kappa}{\sqrt{1 + R^2 \kappa^2}} \cos(\sqrt{1 + R^2 \kappa^2} \phi + \phi_0) + B, \quad (3.19)$$




FIGURE 3. Sketch of the minimizer in the circular wire. Spin configuration is drawn according to equation (3.28) with an arbitrarily chosen value of  $\gamma = 2.3$ .

where

$$A^2 + B^2(1 + R^2\kappa^2) = 1. \quad (3.20)$$

We also need to make sure that periodicity conditions on  $\mathbf{v}$  are satisfied and therefore, provided  $A \neq 0$ , we have the following condition

$$(1 + R^2\kappa^2) = n^2, \quad (3.21)$$

where  $n \in \mathbb{N}$ . Note that all these solutions yield zero micromagnetic energy (with absent stray field), and hence, depending on the relation between  $R$  and  $\kappa$ , we might have several minimizers. Two minimizers always exists by taking  $A = 0$  and  $B = \pm \frac{1}{\sqrt{1+R^2\kappa^2}}$  (see Fig. 3).

It is interesting to observe that the introduction of DMI breaks the degeneracy of exchange interaction, and instead of any constant solution being a minimizer of symmetric exchange energy we obtain only two global minimizers of energy  $\mathcal{E}_C^0$  with anti-symmetric exchange

$$\mathbf{v}(\phi) = \mp \frac{R\kappa}{\sqrt{1 + R^2\kappa^2}} \mathbf{t} \pm \frac{1}{\sqrt{1 + R^2\kappa^2}} \mathbf{b}. \quad (3.22)$$

Moreover, when condition (3.21) is satisfied we have a continuum of solutions (*cf.* (3.17)–(3.19)) with  $A$  and  $B$  satisfying (3.20).

An alternative approach to solving the problem sheds light on non-trivial periodic solutions beyond condition (3.21). Let's define an auxiliary unit vector field as follows:

$$\tilde{\mathbf{v}} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(\alpha) & \sin(\alpha) \\ 0 & -\sin(\alpha) & \cos(\alpha) \end{pmatrix} \cdot \begin{pmatrix} \cos(\phi) & \sin(\phi) & 0 \\ -\sin(\phi) & \cos(\phi) & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot \mathbf{v}, \quad (3.23)$$

where  $\alpha = \arctan(R\kappa)$ . The expression for the energy depending on field  $\tilde{\mathbf{v}}$  has the following form:

$$\begin{aligned} \mathcal{E}_C^0(\tilde{\mathbf{v}}) = & \frac{1}{2R^2} \int_0^{2\pi} |\partial_\phi \tilde{\mathbf{v}}|^2 d\phi + \frac{\sqrt{1+R^2\kappa^2}}{R^2} \int_0^{2\pi} (\tilde{v}_1 \partial_\phi \tilde{v}_2 - \tilde{v}_2 \partial_\phi \tilde{v}_1) d\phi \\ & + \frac{1+R^2\kappa^2}{2R^2} \int_0^{2\pi} (1 - \tilde{v}_3^2) d\phi. \end{aligned}$$

The resulting Hamiltonian clearly shows that the system is frustrated between two states: a homogeneous state polarized along the easy axis or a Néel spiral in a perpendicular plane. We note that it is possible to represent  $\tilde{\mathbf{v}}$  in spherical coordinates as follows:

$$\tilde{\mathbf{v}} = (\sin \theta(\phi) \cos \psi(\phi), \sin \theta(\phi) \sin \psi(\phi), \cos \theta(\phi))$$

and rewrite the above energy as

$$\mathcal{E}_C^0(\theta, \psi) = \frac{1}{2R^2} \int_0^{2\pi} |\theta'|^2 + \sin^2 \theta |\psi' - \sqrt{1+R^2\kappa^2}|^2 d\phi. \quad (3.24)$$

We already have information about minimizers of the energy, but using this representation, we can also find special solutions to the Euler–Lagrange equations

$$-\theta'' + \sin \theta \cos \theta |\psi' - \sqrt{1+R^2\kappa^2}|^2 = 0 \quad (3.25)$$

$$-(\sin^2 \theta \psi')' + \sqrt{1+R^2\kappa^2} (\sin^2 \theta)' = 0 \quad (3.26)$$

Here we have to supplement these with boundary conditions  $\theta(0) = \theta(2\pi) + 2k\pi$ ,  $\psi(0) = \psi(2\pi) + 2m\pi$ ,  $\theta'(0) = \theta'(2\pi)$ ,  $\psi'(0) = \psi'(2\pi)$ . We can take  $\theta = \frac{\pi}{2}$  and  $\psi = n\phi + \phi_0$ ,  $n \in \mathbb{N}$  to deduce

$$\tilde{\mathbf{v}} = (\sin(n\phi + \phi_0), \cos(n\phi + \phi_0), 0). \quad (3.27)$$

Solution (3.27) is almost always a saddle point, except when condition (3.21) is satisfied. With this exceptional condition, the unstable mode disappears, and the solution (3.27) coincides with the minimizer.

If we introduce the magnetostatic energy contribution and study energy  $\mathcal{E}_C$ , in general, the minimization problem cannot be solved explicitly anymore, and numerical simulations are needed to investigate the minimizing configurations. However, we can still find an explicit solution

$$\mathbf{v}(\phi) = \mp \frac{\gamma}{\sqrt{1+\gamma^2}} \mathbf{t} \pm \frac{1}{\sqrt{1+\gamma^2}} \mathbf{b}, \quad (3.28)$$

$$\gamma = \frac{\sqrt{4R^2\kappa^2 + (R^2\kappa^2 + \frac{1}{2\pi}R^2 - 1)^2 + R^2\kappa^2 + \frac{1}{2\pi}R^2 - 1}}{2R\kappa}. \quad (3.29)$$

It corresponds to taking  $\theta$  and  $\psi$  as constants and solving an algebraic problem coming from Euler–Lagrange equations for  $\theta$  and  $\psi$ . Our additional numerical simulations indicate that (3.28) is a good candidate for minimizer in a wide range of values  $R$  and  $\kappa$ . Still, the rigorous clarification of this question is beyond the scope of this work.

**Remark 3.1.** We note that our results also apply to some interactions beyond exchange and DMI. Below, we give one example of such a situation – the Ado interaction – highly nonlinear interaction possible in several

systems with broken inversion symmetry [56, 57]. According to [56, 57], for the crystals with the point group symmetry  $T_d$  we have the following simplified energy

$$\mathcal{G}_\varepsilon(\mathbf{v}) := \frac{1}{\varepsilon^2|Q|} \left( \int_{\Omega_\varepsilon} \frac{1}{2} |D\mathbf{v}|^2 + \beta(v_1v_2v_3)\nabla \cdot \mathbf{v} \, dx \right). \quad (3.30)$$

Given that  $K(\mathbf{v})\mathbf{e}_1 \cdot \partial_x \mathbf{v} + K(\mathbf{v})\mathbf{e}_2 \cdot \partial_y \mathbf{v} + K(\mathbf{v})\mathbf{e}_3 \cdot \partial_z \mathbf{v} = \beta(v_1v_2v_3)\nabla \cdot \mathbf{v}$  when  $K(\mathbf{v}) = \beta(v_1v_2v_3)I$  with  $I$  the  $3 \times 3$  identity matrix, using Theorem 2.1 we can obtain the limiting energy for Ado interactions

$$\mathcal{E}_C(\mathbf{v}) = \frac{1}{2} \int_I |\partial_x \mathbf{v} + \beta v_1 v_2 v_3 \mathbf{e}_1|^2 \, dx - \frac{3\beta^2}{2} \int_I v_1^2 v_2^2 v_3^2 \, dx. \quad (3.31)$$

#### ACKNOWLEDGEMENTS

G.DI F. is a member of Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of INdAM. G.DI F. acknowledges partial support from the Italian Ministry of Education and Research through the PRIN2022 project *Variational Analysis of Complex Systems in Material Science, Physics and Biology* No. 2022HKBF5C, and from the Austrian Science Fund (FWF) through the project *Analysis and Modeling of Magnetic Skyrmions* (grant 10.55776/P34609). G.DI F. also thanks TU Wien and MedUni Wien for their hospitality. F.N.R. acknowledges support from the Swedish Research Council (grant 2023-04899).

#### REFERENCES

- [1] H. Du, J.P. DeGrave, F. Xue, D. Liang, W. Ning, J. Yang, M. Tian, Y. Zhang and S. Jin, Highly stable skyrmion state in helimagnetic MnSi nanowires. *Nano Lett.* **14** (2014) 2026–2032.
- [2] N. Mathur, M.J. Stolt, K. Niitsu, X. Yu, D. Shindo, Y. Tokura and S. Jin, Electron holography and magnetotransport measurements reveal stabilized magnetic skyrmions in  $\text{Fe}_{1-x}\text{Co}_x\text{Si}$  nanowires. *ACS Nano* **13** (2019) 7833–7841.
- [3] Y. Liu, S. Vaitiekėnas, S. Martí-Sánchez, C. Koch, S. Hart, Z. Cui, T. Kanne, S.A. Khan, R. Tanta, S. Upadhyay, M.E. Cachaza, C.M. Marcus, J. Arbiol, K.A. Moler and P. Krogstrup, Semiconductor–ferromagnetic insulator–superconductor nanowires: stray field and exchange field. *Nano Lett.* **20** (2020) 456–462.
- [4] S. Vaitiekėnas, Y. Liu, P. Krogstrup and C.M. Marcus, Zero-bias peaks at zero magnetic field in ferromagnetic hybrid nanowires. *Nat. Phys.* **17** (2021) 43–47.
- [5] M. Schöbitz, A. De Riz, S. Martin, S. Bochmann, C. Thirion, J. Vogel, M. Foerster, L. Aballe, T. Mentès, A. Locatelli, F. Genuzio, S. Le-Denmat, L. Cagnon, J. Toussaint, D. Gusakova, J. Bachmann and O. Fruchart, Fast domain wall motion governed by topology and Ørsted fields in cylindrical magnetic nanowires. *Phys. Rev. Lett.* **123** (2019) 217201.
- [6] D. Sanz-Hernández, A. Hierro-Rodríguez, C. Donnelly, J. Pablo-Navarro, A. Sorrentino, E. Pereiro, C. Magén, S. McVitie, J.M. de Teresa, S. Ferrer, P. Fischer and A. Fernández-Pacheco, Artificial double-helix for geometrical control of magnetic chirality. *ACS Nano* **14** (2020) 8084–8092.
- [7] A. Fernández-Pacheco, R. Streubel, O. Fruchart, R. Hertel, P. Fischer and R.P. Cowburn, Three-dimensional nanomagnetism. *Nat. Commun.* **8** (2017) 15756.
- [8] D. Makarov and D. Sheka, *Curvilinear Micromagnetism: From Fundamentals to Applications*. Springer International Publishing (2022).
- [9] W. Huang, C. Gatel, Z.-A. Li and G. Richter, Synthesis of magnetic Fe and Co nano-whiskers and platelets *via* physical vapor deposition. *Mater. Des.* **208** (2021) 109914.
- [10] S. Sahoo, S. Mondal, G. Williams, A. May, S. Ladak and A. Barman, Ultrafast magnetization dynamics in a nanoscale three-dimensional cobalt tetrapod structure. *Nanoscale* **10** (2018) 9981–9986.
- [11] L. Keller, M.K.I. Al Mamoori, J. Pieper, C. Gspan, I. Stockem, C. Schröder, S. Barth, R. Winkler, H. Plank, M. Pohlit, J. Müller and M. Huth, Direct-write of free-form building blocks for artificial magnetic 3D lattices. *Sci. Rep.* **8** (2018) 6160.
- [12] S. Gliga, G. Seniutinas, A. Weber and C. David, Architectural structures open new dimensions in magnetism: magnetic buckyballs. *Mater. Today* **26** (2019) 100–101.

- [13] O.M. Volkov, O.V. Pylypovskiy, F. Porrati, F. Kronast, J.A. Fernandez-Roldan, A. Kkay, A. Kuprava, S. Barth, F.N. Rybakov, O. Eriksson, S. Lamb-Camarena, P. Makushko, M.-A. Mawass, S. Shakeel, O.V. Dobrovolskiy, M. Huth and D. Makarov, Three-dimensional magnetic nanotextures with high-order vorticity in soft magnetic wireframes. *Nat. Commun.* **15** (2024) 2193.
- [14] A. Prohl, Computational micromagnetism, in *Advances in Numerical Mathematics*. B. G. Teubner, Stuttgart (2001).
- [15] M. Kružík and A. Prohl, Recent developments in the modeling, analysis, and numerics of ferromagnetism. *SIAM Rev.* **48** (2006) 439–483.
- [16] C.J. García-Cervera, Numerical micromagnetics: a review. *Bol. Soc. Esp. Mat. Apl. SeMA* **39** (2007) 103–135.
- [17] L. Exl, J. Fischbacher, A. Kovacs, H. Oezelt, M. Gusenbauer and T. Schrefl, Preconditioned nonlinear conjugate gradient method for micromagnetic energy minimization. *Comput. Phys. Commun.* **235** (2019) 179–186.
- [18] F. Zheng, F.N. Rybakov, N.S. Kiselev, D. Song, A. Kovács, H. Du, S. Blügel and R.E. Dunin-Borkowski, Magnetic skyrmion braids. *Nat. Commun.* **12** (2021) 5316.
- [19] R. Cheenikundil and R. Hertel, Switchable magnetic frustration in buckyball nanoarchitectures. *Appl. Phys. Lett.* **118** (2021) 212403.
- [20] G. Gioia and R.D. James, Micromagnetics of very thin films. *Proc. Roy. Soc. Lond. Ser. A: Math. Phys. Eng. Sci.* **453** (1997) 213–223.
- [21] G. Carbou, Thin layers in micromagnetism. *Math. Models Methods Appl. Sci.* **11** (2001) 1529–1546.
- [22] A. DeSimone, R.V. Kohn, S. Müller and F. Otto, A reduced theory for thin-film micromagnetics. *Commun. Pure Appl. Math.* **55** (2002) 1408–1460.
- [23] A. DeSimone, R.V. Kohn, S. Müller, F. Otto and R. Schäfer, Two-dimensional modelling of soft ferromagnetic films. *Roy. Soc. Lond. Proc. Ser. A. Math. Phys. Eng. Sci.* **457** (2001) 2983–2991.
- [24] R.V. Kohn and V.V. Slastikov, Another thin-film limit of micromagnetics. *Arch. Rational Mech. Anal.* **178** (2005) 227–245.
- [25] R. Moser, Boundary vortices for thin ferromagnetic films. *Arch. Rational Mech. Anal.* **174** (2004) 267–300.
- [26] R. Moser, Moving boundary vortices for a thin-film limit in micromagnetics. *Commun. Pure Appl. Math.* **58** (2005) 701–721.
- [27] V. Slastikov, Micromagnetics of thin shells. *Math. Models Methods Appl. Sci.* **15** (2005) 1469–1487.
- [28] G. Di Fratta, C.B. Muratov and V.V. Slastikov, Reduced energies for thin ferromagnetic films with perpendicular anisotropy. *Math. Models Methods Appl. Sci.* **34** (2024) 1861–1904.
- [29] R.V. Kohn and V.V. Slastikov, Effective dynamics for ferromagnetic thin films: a rigorous justification. *Proc. Roy. Soc. Lond. Ser. A. Math. Phys. Eng. Sci.* **461** (2005) 143–154.
- [30] C. Melcher, Thin-film limits for Landau–Lifshitz–Gilbert equations. *SIAM J. Math. Anal.* **42** (2010) 519–537.
- [31] C.J. García-Cervera and E. Weinan, Effective dynamics for ferromagnetic thin films. *J. Appl. Phys.* **90** (2001) 370–374.
- [32] A. Capella, C. Melcher and F. Otto, Wave-type dynamics in ferromagnetic thin films and the motion of Néel walls. *Nonlinearity* **20** (2007) 2519–2537.
- [33] I. Dzyaloshinsky, A thermodynamic theory of “weak” ferromagnetism of antiferromagnetics. *J. Phys. Chem. Solids* **4** (1958) 241–255.
- [34] T. Moriya, Anisotropic superexchange interaction and weak ferromagnetism. *Phys. Rev.* **120** (1960) 91–98.
- [35] A. Fert, N. Reyren and V. Cros, Magnetic skyrmions: advances in physics and potential applications. *Nat. Rev. Mater.* **2** (2017) 1–15.
- [36] E. Davoli, G. Di Fratta, D. Praetorius and M. Ruggeri, Micromagnetics of thin films in the presence of Dzyaloshinskii–Moriya interaction. *Math. Models Methods Appl. Sci.* **32** (2022) 911–939.
- [37] C. Melcher, Chiral skyrmions in the plane. *Proc. Roy. Soc. Lond. Ser. A. Math. Phys. Eng. Sci.* **470** (2014) 20140394.
- [38] C.B. Muratov and V.V. Slastikov, Domain structure of ultrathin ferromagnetic elements in the presence of Dzyaloshinskii–Moriya interaction. *Proc. Roy. Soc. A: Math. Phys. Eng. Sci.* **473** (2017) 20160666.
- [39] C.B. Muratov, V.V. Slastikov, A.G. Kolesnikov and O.A. Tretiakov, Theory of the dzyaloshinskii domain-wall tilt in ferromagnetic nanostrips. *Phys. Rev. B* **96** (2017) 134417.
- [40] G. Di Fratta, Micromagnetics of curved thin films. *Z. Angew. Math. Phys.* **71** (2020) 111.
- [41] G. Di Fratta, C.B. Muratov, F. Rybakov and V.V. Slastikov, Variational principles of micromagnetics revisited. *SIAM J. Math. Anal.* **52** (2020) 3580–3599.



- [42] G. Di Fratta and V. Slastikov, Curved thin-film limits of chiral Dirichlet energies. *Nonlinear Anal.* **234** (2023) 113303.
- [43] V.V. Slastikov and C. Sonnenberg, Reduced models for ferromagnetic nanowires. *IMA J. Appl. Math.* **77** (2012) 220–235.
- [44] W.F. Brown, *Micromagnetics*. Interscience Publishers, London (1963).
- [45] A. Hubert and R. Schäfer, *Magnetic Domains: The Analysis of Magnetic Microstructures*. Springer Science & Business Media (2008).
- [46] L. Landau and E. Lifshitz, On the theory of the dispersion of magnetic permeability in ferromagnetic bodies, in *Perspectives in Theoretical Physics*. Elsevier (1992) 51–65.
- [47] F. Alouges and G. Di Fratta, Homogenization of composite ferromagnetic materials. *Proc. Roy. Soc. A: Math. Phys. Eng. Sci.* **471** (2015) 20150365.
- [48] E. Davoli and G. Di Fratta, Homogenization of chiral magnetic materials: a mathematical evidence of Dzyaloshinskii's predictions on helical structures. *J. Nonlinear Sci.* **30** (2020) 1229–1262.
- [49] E. Davoli, G. Di Fratta and R. Giorgio, A Bourgain–Brezis–Mironescu formula accounting for nonlocal antisymmetric exchange interactions. *SIAM J. Math. Anal.* **56** (2024) 6995–7013.
- [50] L. Landau and E. Lifshitz, *Statistical Physics, Vol. 5*. Elsevier (2013).
- [51] D. Harutyunyan, On the existence and stability of minimizers in ferromagnetic nanowires. *J. Math. Anal. Appl.* **434** (2016) 1719–1739.
- [52] O. Boule, S. Rohart, L.D. Buda-Prejbeanu, E. Jué, I.M. Miron, S. Pizzini, J. Vogel, G. Gaudin and A. Thiaville, Domain wall tilting in the presence of the Dzyaloshinskii–Moriya interaction in out-of-plane magnetized magnetic nanotracks. *Phys. Rev. Lett.* **111** (2013) 217203.
- [53] A.B. Borisov and Y.A. Izyumov, Nonlinear excitation in spiral magnetic structures. *Dokl. Akad. Nauk SSSR* **283** (1985) 859–861.
- [54] G. Di Fratta, A. Fiorenza and V. Slastikov, On symmetry of energy minimizing harmonic-type maps on cylindrical surfaces. *Math. Eng.* **5** (2023) 1–38.
- [55] G. Carbou, M. Moussaoui and R. Rachi, Stability of steady states in ferromagnetic rings. *J. Math. Phys.* **63** (2022) Paper No. 031508, 28.
- [56] F.N. Rybakov, A. Pervishko, O. Eriksson and E. Babaev, Antichiral ferromagnetism. *Phys. Rev. B* **104** (2021) L020406.
- [57] I. Ado, O. Tchernyshyov and M. Titov, Noncollinear ground state from a four-spin chiral exchange in a tetrahedral magnet. *Phys. Rev. Lett.* **127** (2021) 127204.



**Please help to maintain this journal in open access!**

This journal is currently published in open access under the Subscribe to Open model (S2O). We are thankful to our subscribers and supporters for making it possible to publish this journal in open access in the current year, free of charge for authors and readers.

Check with your library that it subscribes to the journal, or consider making a personal donation to the S2O programme by contacting [subscribers@edpsciences.org](mailto:subscribers@edpsciences.org).

More information, including a list of supporters and financial transparency reports, is available at <https://edpsciences.org/en/subscribe-to-open-s2o>.