Journal of Physics A:
Mathematical and
Theoretical

PURPOSE-LED
PUBLISHING™

PAPER • **OPEN ACCESS**

The discrete analogue of the Gaussian

To cite this article: Gautam Chinta *et al* 2025 *J. Phys. A: Math. Theor.* **58** 035203

View the article online for updates and enhancements.

You may also like

- To share and not share a singlet: control qubit and nonclassicality in teleportation
Kornikar Sen, Adithi Ajith, Saronath Halder et al.
- Implications of pinned occupation numbers for natural orbital expansions: I. Generalizing the concept of active spaces
Christian Schilling, Carlos L Benavides-Riveros, Alexandre Lopes et al.
- Exact partition function of the Potts model on the Sierpinski gasket and the Hanoi lattice
P D Alvarez

This content was downloaded from IP address 130.238.172.43 on 10/02/2025 at 15:44

OPEN ACCESS
IOP Publishing
Journal of Physics A: Mathematical and Theoretical
J. Phys. A: Math. Theor. **58** (2025) 035203 (22pp)
https://doi.org/10.1088/1751-8121/ada64b

The discrete analogue of the Gaussian

Gautam Chinta[1,5,6], **Jay Jorgenson**[1,6], **Anders Karlsson**[2,3,7,*] **and Lejla Smajlović**[4]

[1] Department of Mathematics The City College of New York Convent Avenue at 138th Street, New York, NY 10031 United States of America
[2] Section de mathématiques Université de Genève, Case Postale 64, 1211 Genève 4, Switzerland
[3] Matematiska institutionen Uppsala universitet Box 256, 751 05 Uppsala, Sweden
[4] Department of Mathematics and Computer Science University of Sarajevo Zmaja od Bosne 35, 71 000 Sarajevo, Bosnia and Herzegovina

E-mail: anders.karlsson@unige.ch, gchinta@ccny.cuny.edu, jjorgenson@mindspring.com and lejlas@pmf.unsa.ba

Received 25 September 2024; revised 5 December 2024
Accepted for publication 6 January 2025
Published 16 January 2025

Abstract

This paper illustrates the utility of the heat kernel on $\mathbb{Z}$ as the discrete analogue of the Gaussian density function. The heat kernel on $\mathbb{Z}$ is the two-variable function $K_{\mathbb{Z}}(t,x) = \mathrm{e}^{-2t} I_x(2t)$ where $I_x(2t)$ is a Bessel function, with variables $x \in \mathbb{Z}$ and $t \geqslant 0$. Like its classic counterpart, $K_{\mathbb{Z}}(t,x)$ appears in many mathematical and physical contexts and has a wealth of applications. Some of these applications will be reviewed here, and they concern Bessel integrals, trigonometric sums, hypergeometric functions and asymptotics of discrete models appearing in statistical and quantum physics. Moreover, we prove a new local limit theorem for sums of integer-valued random variables, obtain novel special

[5] The first-named author acknowledges support from the Simons Foundation MPS-TSM program.
[6] The first and second-named authors acknowledge grant support from PSC-CUNY Awards 65400-00-53 and 65400-00-55, which are jointly funded by the Professional Staff Congress and The City University of New York.
[7] The third-named author acknowledges grant support by the Swiss NSF Grants 200020-200400, 200021-212864 and the Swedish Research Council Grant 104651320.
[*] Author to whom any correspondence should be addressed.

Original Content from this work may be used under the terms of the Creative Commons Attribution 4.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

values of the spectral zeta function of Bethe lattices, and provide a discussion on how $e^{-2t}I_x(2t)$ could be useful in differential privacy.

Keywords: heat kernels, discrete models, spectral zeta functions, local limit laws, Bethe lattice, Bessel identities, differential privacy

1. Introduction

The *Gaussian density function*, or simply *Gaussian*, is the two-variable function defined by

$$K_{\mathbb{R}}(t,x) := \frac{1}{\sqrt{4\pi t}} e^{-x^2/4t} \quad \text{for} \quad x \in \mathbb{R} \quad \text{and} \quad t \in \mathbb{R}^+. \tag{1}$$

In common language the Gaussian is sometimes referred to as the bell-shaped curve. In probability theory, (1) is the density function of a normal random variable. The universality of the Gaussian is evident from the classical central limit theorem, the theory of Brownian motion, and that (1) serves as the fundamental solution to the heat equation on the real line

$$\left(-\frac{\partial^2}{\partial x^2} + \frac{\partial}{\partial t}\right) f(t,x) = 0. \tag{2}$$

In (2) t is positive time variable while in (1) we have that $t = \sigma^2/2$ where σ^2 is the variance of the corresponding probability distribution.

A main point of the present article is to advocate for the following assertion:
If instead of having $x \in \mathbb{R}$, a continuum, one has that $x \in \mathbb{Z}$, the discrete space of integers, then the discrete analogue of (1) is

$$K_{\mathbb{Z}}(t,x) := e^{-2t} I_x(2t) \tag{3}$$

where $I_x(2t)$ is the I-Bessel function defined either by the integral

$$I_x(z) = \frac{1}{\pi} \int_0^\pi e^{z\cos(\theta)} \cos(x\theta)\, d\theta \tag{4}$$

or by the series representation

$$I_x(z) = \sum_{n=0}^{\infty} \frac{(z/2)^{2n+x}}{n!\Gamma(n+1+x)} \tag{5}$$

for $x \geqslant 0$ and $I_{-x}(z) := I_x(z)$. See figure 1.

The phrase discrete Gaussian is already used in multiple ways in the literature, especially in the field of lattice-based cryptography, and it often means the probability density function associated to a $\mathbb{Z}$-valued random variable W for which

$$\text{Prob}(W = n) = c e^{-n^2/4t} \quad \text{for} n \in \mathbb{Z} \tag{6}$$

with the normalizing constant c necessarily given by

$$c^{-1} = \sum_{k=-\infty}^{\infty} e^{-k^2/4t}; \tag{7}$$



Figure 1. The dots show the discrete Gaussian at $t=1$ as a function of integers x and the curve is the ordinary Gaussian with $t=1$ as a function of real numbers x .

see, for example, [CKS22]. (The constant (7) is known as a Thetanullwerte $\theta(0,i/2t)$ and is an object of extensive mathematical significance; see, for example, [Mu83].) On the other hand, in [CKS22], (6) is recognized as a discretized Gaussian; see the discussion before Definition 1.1 in [CKS22]. Other authors, such as [Li24], distinguish between (6), which they refer to as the sampled Gaussian, and (3), which [Li24] calls the discrete analogue of the Gaussian; see section 2.6 of [Li24]. See also [GK24], where the discretized Gaussian denotes a (continuous) Gaussian quantized on a lattice.
Throughout this article, we will refer to (3) as the discrete Gaussian and to (6) either as the discretized or sampled Gaussian.

We now turn to discussing the many ways the discrete Gaussian satisfies structural results which are analogous to those which are fulfilled by (1).

One immediate justification for calling (3) the discrete Gaussian is that it is the fundamental solution to the heat equation on $\mathbb{Z}$. More precisely, for $x\in\mathbb{Z}$ and $t\in\mathbb{R}^{+}$, we have that

$$\left(\Delta_{\mathbb{Z}}+\frac{\partial}{\partial t}\right)K_{\mathbb{Z}}(t,x)=0 \quad \text{with} \quad K_{\mathbb{Z}}(0,x)=\delta_0(x) \tag{8}$$

and where

$$\Delta_{\mathbb{Z}}f(x)=2f(x)-f(x+1)-f(x-1)\,.$$

As far as we know, (6) does not satisfy such an elementary differential or difference equation. We note that operator $\Delta_{\mathbb{Z}}$ appears in various physics contexts, such as the study of discrete random Schrödinger operators and Anderson localization [An58]. We call the operator $\Delta_{\mathbb{Z}}$ a discrete, or combinatorial, Laplacian because it can be defined in analogy with the usual Laplacian

$$\Delta_{\mathbf{R}}=-\frac{\partial^2}{\partial x^2}$$

see section 2 below. Moreover, $\Delta_{\mathbb{Z}}$ is in fact a discretization of $\Delta_{\mathbf{R}}$.

The claim that $\mathrm{e}^{-2t}I_x(2t)$ satisfies the heat equation (8) follows from the basic relation

$$I_{x+1}(t) + I_{x-1}(t) = 2\frac{\mathrm{d}}{\mathrm{d}t}I_x(t),$$

which in turn can easily be derived from the definitions (4) or (5). The solution of (8) has been discovered and rediscovered over the years; see for example [Ba49, Fe66, GI02, KN06]. Nonetheless, we feel it bears further emphasis, particularly in light of recently emerging applications, some of which we review in this article.

Let us first point out the following rescaled convergence, which makes precise the intuitive statement that the discrete heat kernel approaches the continuous heat kernel as the discretization becomes arbitrarily fine.

Proposition 1. *Let $\{\alpha_n\}$ be any sequence of integers indexed by $n \in \mathbb{Z}^+$ such that*

$$\alpha_n/n \to x \in \mathbb{R} \ \text{ as } \ n \to \infty.$$

Then

$$\lim_{n\to\infty} nK_{\mathbf{Z}}\left(n^2t, \alpha_n\right) = \lim_{n\to\infty} n\mathrm{e}^{-2n^2t}I_{\alpha_n}\left(2n^2t\right) = \frac{1}{\sqrt{4\pi t}}\mathrm{e}^{-x^2/4t}.$$

Proposition 1 was proved by Athreya in [At87]; we provide a different proof in section 3. Alternative asymptotic expressions can be found in Pang [Pa93] and Cowling–Meda–Setti [CMS00]. Both of these works show how estimates for the heat kernel on $\mathbb{Z}$ can be used to derive estimates for heat kernels on more general regular graphs and trees; see section 2.

Note here the following philosophical point. While it is understood from numerical analysis that for a compact space each discrete eigenvalue converges after rescaling to the corresponding continuous one when the mesh size goes to zero, there seems to be no general uniformity result associated to this convergence. Indeed, the discrete problem has only a finite number of eigenvalues while the continuous one has an infinitude (or for unbounded spaces the spectrum may even be purely continuous). The above proposition therefore shows a significant advantage in using the discrete heat kernel which will yield an important application in section 6 below. In general terms, the rescaled discrete Gaussian is a good way of packaging all the spectral information when passing from a lattice to a continuum.

In statistics, as noted in [DH13], practitioners often face a significant inaccuracy when approximating integer-valued ("lattice-valued") random variables by the real Gaussian as suggested by the central limit theorem. The following theorem proved in section 3 responds to this general problem by instead proposing an approximation using the discrete Gaussian.

Theorem 2. *Let $\{X_k\}_{k=1}^{\infty}$ be a sequence of independent, identically distributed integer-valued random variables with finite mean μ and finite variance σ^2, and assume that $\sigma^2 > |\mu|$. Assume further there is no infinite subprogression $a + \ell\mathbb{Z}$ of $\mathbb{Z}$ with $\ell > 1$ such that X_1 takes values in $a + \ell\mathbb{Z}$ almost surely. Let $S_n := \sum_{k=1}^{n} X_k$. Then*

$$\sup_{m\in\mathbb{Z}} \sqrt{n}\left|\mathbf{P}(S_n = m) - \left(\frac{\sigma^2+\mu}{\sigma^2-\mu}\right)^{m/2} \mathrm{e}^{-n\sigma^2} I_m\left(n\sqrt{\sigma^4-\mu^2}\right)\right| \to 0 \ \text{ as } n \to \infty.$$

Spectral zeta functions are relevant in particular for physics, see [El12] and further comments in section 7. Their special values for graphs have received some recent interest in number

theory [FK17, XZZ22, KP23]. We obtain new explicit values for the spectral zeta function of Bethe lattices, meaning the $(q+1)$-regular trees T_{q+1}.

Theorem 3. *For any $q \geqslant 1$, let $\zeta_{T_{q+1}}(s)$ be the spectral zeta function associated to T_{q+1}. Then $\zeta_{T_{q+1}}(0) = 1$ and $\zeta_{T_{q+1}}(-1) = q+1$. Furthermore, for any integer $m > 1$, we have that*

$$\zeta_{T_{q+1}}(-m) = \sum_{k=0}^{m} \binom{m}{k}^2 q^{m-k} - (q-1) \sum_{j=1}^{\lfloor m/2 \rfloor} \sum_{k=0}^{m-2j} \binom{m}{k} \binom{m}{2j+k} q^{m-2j-k}. \quad (9)$$

Moreover,

$$\det{}'\Delta_{T_{q+1}} := \exp\left(-\zeta'_{T_{q+1}}(0)\right) = \begin{cases} \left(1-q^{-2}\right)^{(1-q)/2}, & \text{when } q > 1 \\ 1, & \text{when } q = 1. \end{cases}$$

To close this introduction we highlight some key features of the discrete Gaussian (3) which we explore more thoroughly below.

- (Section 2) The discrete Gaussian is not just the heat kernel on $\mathbb{Z}$, but in fact it serves as the building block for heat kernels on all regular graphs. em (section 3) The discrete Gaussian appears in the limiting density function in the local central limit law for a sequence of identically distributed integer-valued random variables with mean zero and finite variance
- (Section 4) Physical and probabilistic principles apply to the discrete Gaussian which then motivate and make obvious a number of Bessel function identities.
- (Section 5) An application of the Laplace transform to appropriate functionals of the discrete Gaussian leads to explicit evaluation of certain finite trigonometric sums appearing in physics such as the Verlinde formulas, resistance in electrical networks or chiral Potts model. em (section 6) The discrete Gaussian appears in the study of asymptotics of the determinant of the Laplacians in lattices with periodic boundary conditions, of relevance in statistical physics and quantum field theory.
- (Section 7) The spectral zeta function arises from a Mellin transform of the heat kernel. When considering the discrete Gaussian, one obtains a number of identities involving known zeta functions, and some interesting new computations as well.
- (Section 8) The field of differential privacy is a mathematically rigorous methodology by which one measures the output of fixed algorithm $\mathcal{M}$ on a given dataset $\mathcal{D}$ with the goal of preventing one from determining any specific entry in $\mathcal{D}$ through repeated use of $\mathcal{M}$; see [CKS22] and references therein. In section 8 we will comment on the potential usefulness of the discrete Gaussian (3), as opposed to the discretized Gaussian (6), in the analysis in [CKS22]. Going further, we will point out that the analysis of section 3 admits an immediate extension to random variables whose values lie in higher dimensional lattices; see, for example, [AA19].

We think the examples listed above are already a noteworthy variety of applications of the discrete Gaussian, and we are confident that there are more to come.

2. The building block of heat kernels of graphs

Let $X = (VX, EX)$ be a countable graph of bounded degree, where VX and EX denote the sets of vertices and edges, respectively. The *Laplacian* is an operator on $L^2(VX, \mathbf{C})$ defined by

$$(\Delta_X f)(x) = \sum_{y\sim x} (f(x) - f(y)), \tag{10}$$

where the sum is taken over all vertices adjacent to x in X.

A graph is called $(q+1)$-*regular* or *regular* if every vertex has degree $q+1$. The reason for the normalization of '$q+1$' comes form number theory, and we adopt it here as well, because various formulas become slightly more compact when using this convention. Chung and Yau observed in [CY99] that the heat kernel on a regular graph can be expressed in terms of the heat kernel on a regular tree. Building on this, Chinta–Jorgenson–Karlsson [CJK15] proved that the I-Bessel function serves as a crucial ingredient in constructing the heat kernel on any regular graph, similar to the role of the Gaussian $\frac{1}{\sqrt{4\pi t}} e^{-x^2/4t}$ in the heat kernel on homogeneous manifolds. In a related result, Grünbaum and Iliev point out that $e^{-2t} I_x(2t)$ generates, via Darboux transformations, solutions of more complicated differential-difference Equations on the discrete line, again akin to the Gaussian on the real line [GI02]. Let us now be more specific.

Let T_{q+1} be a $(q+1)$-regular tree, also called the *the Bethe lattice*. Chung and Yau [CY99] give an explicit expression for its heat kernel in a radial coordinate r. That is, since the heat kernel is obviously radially symmetric, one first selects a vertex as an origin 0 and then sets $K_{T_{q+1}}(t,x) = K(t,r)$ where r is the distance between 0 and x. In [CY99] it is shown that

$$K(t,r) = \frac{2e^{-(q+1)t}}{\pi q^{r/2-1}} \int_0^\pi \frac{\exp\left(2t\sqrt{q}\cos u\right) \sin u \left(q \sin(r+1)u - \sin(r-1)u\right)}{(q+1)^2 - 4q\cos^2 u} du$$

for $r \geqslant 0$. Cowling–Meda–Setti [CMS00] presented a different formula for $K(t,r)$, namely that

$$K(t,r) = q^{-r/2} e^{-(q+1)t} I_r(2\sqrt{q}t) - (q-1) \sum_{j=1}^{\infty} q^{-(r+2j)/2} e^{-(q+1)t} I_{r+2j}(2\sqrt{q}t). \tag{11}$$

We note that the normalization of the Laplacian in [CMS00] is different than what we use in this article. The formula (11) was rediscovered by different means in [CJK15].

One feature of (11) is that we can think of

$$q^{-n/2} e^{-(q+1)t} I_n(2\sqrt{q}t) \quad \text{for} \quad n \geqslant 0 \tag{12}$$

as a type of building block for the heat kernel on the $(q+1)$-regular tree. More generally, in the case of a $(q+1)$-regular graph, we have a similar expression, as given in the following theorem.

Theorem 4 [CJK15]. *The heat kernel on any $(q+1)$-regular graph X is*

$$K_X(t,x) = e^{-(q+1)t} \sum_{m=0}^{\infty} b_m(x) q^{-m/2} I_m(2\sqrt{q}t),$$

where $b_m(x) = c_m(x) - (q-1)(c_{m-2}(x) + c_{m-4}(x) + \ldots)$ and $c_m(x)$ is the number of geodesics from the origin to x of length $m \geqslant 0$.

It is interesting to compare theorem 4 with known results in the setting of Riemannian geometry, where the heat kernel $K_M(t,x,y)$ on an n-dimensional smooth manifold M with smooth metric has an asymptotic expansion of the form

$$K_M(t,x,y) = \frac{\mathrm{e}^{-d_M^2(x,y)/4t}}{(4\pi t)^{n/2}} \left(\sum_{j=0}^{k} u_j(x,y)\, t^j + O\left(t^{k+1}\right) \right) \quad \text{as } t \to 0$$

for any integer $k > n/2+2$, points $x, y \in M$, distance function d_M on M, and computable functions $\{u_j\}$; see, for example, page 152 of [Ch84]. In certain cases when M is a non-compact symmetric space, the heat kernel can be written as

$$F(r,t) \cdot \mathrm{e}^{-at} \cdot \frac{1}{(4\pi t)^{n/2}} \mathrm{e}^{-r^2/4t} \quad \text{where } r = d_M(x,y)$$

where a is a certain constant depending upon the dimension n of the space, and $F(r,t)$ is an explicitly computable power series; see, for example, [GN98] in the case of hyperbolic spaces $\mathbb{H}^n$.

As a way of motivating the appearance of the Bessel function in (11) and (12) from our perspective, consider the asymmetric Laplacian on $\mathbb{Z}$ defined on L^2 functions $f\colon \mathbb{Z} \to \mathbb{C}$ by

$$\Delta_{p,q} f(x) = (p+q)f(x) - pf(x-1) - qf(x+1) \quad \text{for } p,q > 0 \text{ and } x \in \mathbb{Z}. \tag{13}$$

The corresponding fundamental solution, or heat kernel, can be found in [Fe66] for $p+q=1$, and is given by

$$K_{p,q}(t,x) = \left(\frac{p}{q}\right)^{x/2} e^{-(p+q)t} I_x\left(2\sqrt{pq}t\right). \tag{14}$$

On the other hand, taking $q \in \mathbb{N}$, $x = r > 0$ and $p = 1$ in (13), the Laplacian $\Delta_{1,q}$ becomes the Laplacian $\Delta_{T_{q+1}}$ acting on radially symmetric functions on the tree T_{q+1} when expressed in terms of the radial distance $r = d(y_0, y)$. Here, y_0 is an arbitrarily chosen root and y is any vertex on the tree different from the root. Explicitly,

$$\Delta_{1,q} f(r) = (q+1)f(r) - qf(r+1) - f(r-1).$$

The heat kernel $K_{1,q}(t,r)$ defined by (14) for $r \geqslant 0$ coincides with the "building block" of heat kernels on $(q+1)$-regular graphs. (Note that for $r=0$ we have $\Delta_{T_{q+1}} f(0) = (q+1)f(0) - (q+1)f(1)$, hence $K_{1,q}(t,r)$ is not the heat kernel on the tree T_{q+1}.) Another path-graph interpretation of the "building block" instead via a horospherical transform can be found in [CJK15, section 3.2].

3. Discrete local limit theorem

Let us continue with the setting of the last paragraph with $p,q > 0$ and $p+q=1$. Then the heat kernel $K_{p,q}(2t,x)$ with $x \in \mathbb{Z}$ can be viewed as the probability distribution of the continuous-time random walk on $\mathbb{Z}$ which steps one unit to the left (resp. right) with probability p (resp. $q = 1-p$.) Further, the time interval between steps has an exponential distribution with density $2\mathrm{e}^{-2t}$, see [Fe66] or [Fe70, chapter 7]. In the special case when $p = q = 1/2$, then the function $K_{1/2,1/2}(2t,x) = \mathrm{e}^{-2t} I_x(2t)$ is the heat kernel on $\mathbb{Z}$.

Let us denote by $Y_{p,q,t}$ the integer-valued random variable such that

$$\mathbf{P}\left(Y_{p,q,t}=m\right)=K_{p,q}\left(t,m\right),\quad m\in\mathbb{Z}. \tag{15}$$

The characteristic function $\varphi_{Y_{p,q,t}}$ of $Y_{p,q,t}$ can be computed using the explicit formula for the generating function for Bessel functions. Indeed, from [Fe70, formula (7.8)] we have that

$$\begin{aligned}\varphi_{Y_{p,q,t}}\left(y\right)&=\mathrm{e}^{-(p+q)t}\sum_{x=-\infty}^{\infty}I_x\left(2\sqrt{pq}t\right)\left(\sqrt{\frac{p}{q}}\mathrm{e}^{\mathrm{i}y}\right)^x\\&=\mathrm{e}^{-(p+q)t(1-\cos y)}\mathrm{e}^{\mathrm{i}(p-q)t\sin y}\quad\text{for } y\in\mathbb{R}.\end{aligned} \tag{16}$$

In the following theorem, we prove an analogue of the central limit theorem but with an error bound, or rather a type of Berry–Esseen Theorem, which asserts that under reasonably general conditions the limiting probability distribution for large n for the sum $S_n:=X_1+\ldots+X_n$ of independent identically distributed integer-valued random variables is given by $K_{p,q}(t,x)$, for certain p, q and t which depend on the distribution of X_1.

Theorem 5. *Let $\{X_k\}_{k=1}^{\infty}$ be a sequence of independent, identically distributed integer-valued random variables with finite mean μ and finite variance σ^2, and assume that $\sigma^2>|\mu|$. Assume further there is no infinite subprogression $a+\ell\mathbb{Z}$ of $\mathbb{Z}$ with $\ell>1$ such that X_1 takes values in $a+\ell\mathbb{Z}$ almost surely. Let $S_n:=\sum_{k=1}^{n}X_k$. Then*

$$\sup_{m\in\mathbb{Z}}\sqrt{n}\left|\mathbf{P}\left(S_n=m\right)-\left(\frac{\sigma^2+\mu}{\sigma^2-\mu}\right)^{m/2}\mathrm{e}^{-n\sigma^2}I_m\left(n\sqrt{\sigma^4-\mu^2}\right)\right|\to 0\ \text{ as } n\to\infty.$$

Proof. For positive integers n, let Y_n denote the integer valued random variable $Y_{p,q,t}$ of (15) with $p+q=1$, $t=n\sigma^2$ and $(p-q)t=n\mu$. The assumption that $\sigma^2>|\mu|$ implies that both p and q are non-negative. Upon substituting into (14), we get that

$$\mathbf{P}\left(Y_n=m\right)=\left(\frac{\sigma^2+\mu}{\sigma^2-\mu}\right)^{m/2}\mathrm{e}^{-n\sigma^2}I_m\left(n\sqrt{\sigma^4-\mu^2}\right). \tag{17}$$

Further, the characteristic function φ_{Y_n} is

$$\varphi_{Y_n}\left(y\right)=\mathrm{e}^{-2n\sigma^2\sin^2(y/2)}\mathrm{e}^{\mathrm{i}\mu n\sin y}\quad\text{for } y\in\mathbb{R};$$

hence,

$$\mathbf{P}\left(Y_n=m\right)=\frac{1}{2\pi}\int_{-\pi}^{\pi}\mathrm{e}^{-2n\sigma^2\sin^2(\theta/2)}\mathrm{e}^{\mathrm{i}\mu n\sin\theta}\mathrm{e}^{-\mathrm{i}m\theta}\mathrm{d}\theta.$$

Set $X=X_1$, and let $\varphi_X(\theta)=\mathrm{e}^{\mathrm{i}\mu\theta}\mathbf{E}(\mathrm{e}^{\mathrm{i}\theta(X-\mu)})=\mathrm{e}^{\mathrm{i}\mu\theta}\varphi_{X-\mu}(\theta)$ be the characteristic function of X. Then we have that

$$\mathbf{P}\left(S_n=m\right)=\frac{1}{2\pi}\int_{-\pi}^{\pi}\left(\varphi_{X-\mu}\left(\theta\right)\right)^n\mathrm{e}^{\mathrm{i}\mu n\theta}\mathrm{e}^{-\mathrm{i}m\theta}d\theta.$$

Combining the above two displayed equations, we conclude that

$$\mathbf{P}(S_n = m) - \mathbf{P}(Y_n = m) = \frac{1}{2\pi}\int_{-\pi}^{\pi}\left[(\varphi_{X-\mu}(\theta))^n - \mathrm{e}^{-2n\sigma^2\sin^2(\theta/2)}\mathrm{e}^{\mathrm{i}\mu n(\sin\theta-\theta)}\right]\mathrm{e}^{\mathrm{i}\mu n\theta}\mathrm{e}^{-\mathrm{i}m\theta}\mathrm{d}\theta. \tag{18}$$

A simple change of variables in (18) yields that

$$\begin{aligned}&\sqrt{n}\,|\mathbf{P}(S_n = m) - \mathbf{P}(Y_n = m)|\\ &\leqslant \frac{1}{2\pi}\int_{-\pi\sqrt{n}}^{\pi\sqrt{n}}\left|\left(\varphi_{X-\mu}\left(\theta/\sqrt{n}\right)\right)^n - \mathrm{e}^{-2n\sigma^2\sin^2\left(\theta/\left(2\sqrt{n}\right)\right)}\mathrm{e}^{\mathrm{i}\mu n\left(\sin\left(\theta/\sqrt{n}\right)-\theta/\sqrt{n}\right)}\right|\mathrm{d}\theta.\end{aligned} \tag{19}$$

For any $\theta\in\mathbb{R}$, one can use the Taylor series expansions

$$\varphi_{X-\mu}(\tau) = 1 - \frac{\sigma^2}{2}\tau + o\left(\tau^2\right),\quad \sin\tau = \tau - \frac{\tau^3}{6} + o\left(\tau^3\right)$$

in the neighborhood of $\tau = \frac{\theta}{\sqrt{n}} = 0$ to deduce that

$$\lim_{n\to\infty}\left(\left(\varphi_{X-\mu}\left(\theta/\sqrt{n}\right)\right)^n - \mathrm{e}^{-2n\sigma^2\sin^2\left(\theta/\left(2\sqrt{n}\right)\right)}\mathrm{e}^{\mathrm{i}\mu n\left(\sin\left(\theta/\sqrt{n}\right)-\theta/\sqrt{n}\right)}\right) = 0.$$

If one were to substitute (17) into (19), the proof would follow if one could apply the Lebesgue dominated convergence theorem in (19) and conclude that the right-hand side of (19) is $o(1)$ as $n\to\infty$. In other words, it suffices to show that the function

$$\left|\left(\varphi_{X-\mu}\left(\theta/\sqrt{n}\right)\right)^n - \mathrm{e}^{-2n\sigma^2\sin^2\left(\theta/\left(2\sqrt{n}\right)\right)}\mathrm{e}^{i\mu n\left(\sin\left(\theta/\sqrt{n}\right)-\theta/\sqrt{n}\right)}\right|\chi_{\left[-\pi\sqrt{n},\pi\sqrt{n}\right]}(\theta)$$

is dominated by an integrable function on $\mathbb{R}$. Let us prove this assertion now.

By the Taylor series expansion of $\varphi_{X-\mu}(\theta)$ and $e^{-2\sigma^2\sin^2(\theta/2)}$ in the neighborhood of $\theta = 0$, we have that

$$|\varphi_{X-\mu}(\theta)| \leqslant |1 - \frac{\theta^2\sigma^2}{4}| \quad\text{and}\quad e^{-2\sigma^2\sin^2(\theta/2)} \leqslant |1 - \frac{\theta^2\sigma^2}{4}|.$$

Hence, there exists some $\delta > 0$ such that for all θ such that $|\theta| \leqslant \delta\sqrt{n}$ one has that

$$\left|\left(\varphi_{X-\mu}\left(\theta/\sqrt{n}\right)\right)^n - \mathrm{e}^{-2n\sigma^2\sin^2\left(\theta/\left(2\sqrt{n}\right)\right)}\mathrm{e}^{\mathrm{i}\mu n\left(\sin\left(\theta/\sqrt{n}\right)-\theta/\sqrt{n}\right)}\right| \leqslant 2\mathrm{e}^{-\sigma^2\theta^2/4}.$$

Since the random variable X does not take values in an arithmetic subprogression, we have that $|\varphi_{X-\mu}(\theta)| \neq 1$ whenever $\theta\neq 0$. Indeed, if there were such $\theta\neq 0$, then by the triangle inequality $\mathrm{e}^{\mathrm{i}\theta(X-\mu)}$ had to be constant, or deterministic, as a random variable. This contradicts the subprogression hypothesis. Therefore, $|\varphi_{X-\mu}(\theta)| < 1$ for $0 < |\theta| \leqslant \pi$. By continuity, and hence uniform continuity on closed intervals, we deduce that there exists a constant $0 < C_X < 1$ such that $|\varphi_{X-\mu}(\theta/\sqrt{n})| \leqslant C_X$ for all $\delta\sqrt{n} \leqslant |\theta| \leqslant \pi\sqrt{n}$.

The inequality $\sin^2(\theta/(2\sqrt{n})) \geqslant \theta^2/(\pi n)$ holds true for $\theta\in(-\pi\sqrt{n},\pi\sqrt{n})$, so then

$$\mathrm{e}^{-2n\sigma^2\sin^2\left(\theta/\left(2\sqrt{n}\right)\right)} \leqslant \mathrm{e}^{-2\sigma^2\theta^2/\pi},$$

for all $\delta\sqrt{n} \leqslant |\theta| \leqslant \pi\sqrt{n}$. When combining the above bounds, we get that

$$\begin{aligned}&\left|\left(\varphi_{X-\mu}\left(\theta/\sqrt{n}\right)\right)^n - \mathrm{e}^{-2n\sigma^2\sin^2\left(\theta/\left(2\sqrt{n}\right)\right)}\mathrm{e}^{\mathrm{i}\mu n\left(\sin\left(\theta/\sqrt{n}\right)-\theta/\sqrt{n}\right)}\right| \chi_{\left[-\pi\sqrt{n},\pi\sqrt{n}\right]} \\ &\leqslant \left(C_X^n + \mathrm{e}^{-2\sigma^2\theta^2/\pi}\right)\chi_{\left[-\pi\sqrt{n},\pi\sqrt{n}\right]}(\theta) + 2\mathrm{e}^{-\sigma^2\theta^2/4}. \end{aligned} \tag{20}$$

The right-hand-side of (20) is an integrable function on $\mathbb{R}$, uniformly in n, so then the proof of the theorem is complete. □

In the special case when the mean $\mu = \mathbf{E}(X) = 0$, we get the following corollary.

Corollary 1. *In addition to the assumptions as stated in theorem 5, assume that $\mu = 0$. Then*

$$\sup_{m\in\mathbb{Z}} \sqrt{n}\left|\mathbf{P}(S_n = m) - \mathrm{e}^{-n\sigma^2} I_m\left(n\sigma^2\right)\right| \to 0, \quad \text{as } n \to \infty.$$

When the set of i.i.d. random variables $\{X_k\}$ almost surely takes values in an arithmetic progressions $a + \ell\mathbb{Z}$ with $\ell > 1$, one can prove stronger variants of local limit theorems; see for example the classical books [IL71] or [Pe75] as well as the recent survey [SW23].

Another local limit theorem when assuming the conditions of theorem 5 takes the form that

$$\sup_{m\in\mathbb{Z}} \sqrt{n}\left|\mathbf{P}(S_n = m) - \frac{1}{\sqrt{2\pi n}\sigma}\mathrm{e}^{-(m-n\mu)^2/\left(2n\sigma^2\right)}\right| \to 0 \ \text{ as } n \to \infty; \tag{21}$$

see for example theorem 13 of [Pe75]). As noted above,

$$\sum_{m\in\mathbb{Z}} \frac{1}{\sqrt{2\pi n}\sigma}\mathrm{e}^{-m^2/\left(2n\sigma^2\right)} \neq 1,$$

meaning that the function $\frac{1}{\sqrt{2\pi n}\sigma}\mathrm{e}^{-(m-n\mu)^2/(2n\sigma^2)}$ when viewed as a function of variable $m \in \mathbb{Z}$ is not a probability distribution on integers. However, the heat kernel $K_{p,q}(n\sigma^2, m)$ with $p + q = 1$ and $p - q = \mu/\sigma^2$ *is a probability distribution on* $\mathbb{Z}$. We believe this important point provides significant justification for calling $\mathrm{e}^{-2t}I_x(2t)$, $t \geqslant 0$ with $x \in \mathbb{Z}$ the discrete Gaussian.

Finally, let us note here that theorem 5 combined with the asymptotic (21) for the local limit theorem yields that

$$\sup_{m\in\mathbb{Z}} \sqrt{n}\left|\left(\frac{\sigma^2+\mu}{\sigma^2-\mu}\right)^{m/2} \mathrm{e}^{-n\sigma^2} I_m\left(n\sqrt{\sigma^4-\mu^2}\right) - \frac{1}{\sqrt{2\pi n}\sigma}\mathrm{e}^{-(m-n\mu)/2n\sigma^2}\right| \to 0 \ \text{ as } n \to \infty$$

for $\sigma^2 > |\mu|$. The proof is a direct application of the triangle inequality. By taking $\mu = 0$ in the above inequality, one obtains a alternate proof of proposition 1.

4. Application 1: Bessel identities

Bessel functions are ubiquitous in mathematical physics, perhaps first appearing as a solution to Bessel's differential equations arising from the Laplace equation with cylindrical symmetry. A classic text on Bessel functions is Watson's book [Wa44]. The fact that the I-Bessel function is essentially a heat kernel makes many of the classical identities immediate from physical principles. Note that we continue to view $I_x(t)$ as a function in two variables, where the variable t is continuous and where the variable x is discrete. We will now illustrate how to obtain

some known Bessel function identities, following [KN06, K12, CJKS24]. In fact, a very nice and much earlier discussion of this type can be found in [Fe66]. Though none of the Bessel function identities in this section is new, we wish to illustrate how our perspective informs our understanding of these identities.

To begin with, start with a unit amount of heat at time $t = 0$. The principle of the conservation of heat implies that though the heat diffuses over the integers as time passes, the total amount of heat at all points sums to one at any time. Mathematically, this principle immediately implies that

$$\sum_{j=-\infty}^{\infty} \mathrm{e}^{-2t} I_j(2t) = 1, \tag{22}$$

for all $t \geqslant 0$. We will deduce this identity in an alternate way in an example below. If we change the continuous variable from real t to z, we can rewrite (22) as

$$\sum_{j=-\infty}^{\infty} I_j(z) = \mathrm{e}^{z}. \tag{23}$$

Since (23) holds for any real z, the principle of analytic continuation applies and extends the formula to any complex z. In particular, we can rephrase (23) in terms of the J-Bessel $J_n(z)$ to get that

$$\sum_{j=-\infty}^{\infty} i^{-j} J_j(z) = \mathrm{e}^{\mathrm{i}z},$$

which leads to

$$J_0(z) + 2\sum_{k=1}^{\infty} (-1)^k J_{2k}(z) = \cos(z)$$

and

$$\sum_{k=0}^{\infty} (-1)^k J_{2k+1}(z) = \frac{1}{2}\sin(z).$$

By applying the same reasoning as above to the heat kernel of the asymmetric Laplacian (corresponding to diffusion with a drift) as in the previous section, one gets more generally that

$$\sum_{j=-\infty}^{\infty} J_j(z) x^j = \mathrm{e}^{\frac{z}{2}\left(x - x^{-1}\right)}. \tag{24}$$

This generating series identity (24) is sometimes known as the *Schlömilch formula,* and it is also given in [Fe66], though derived through very different means.

Solutions to differential equations have a semi-group property in time; in probability theory, such an identity is the Chapman-Kolmogorov equation. In our case, the resulting formula is that

$$J_n(t+s) = \sum_{k=-\infty}^{\infty} J_{n-k}(t) J_k(s).$$

This is called *Neumann's identity* for the J-Bessel function and is deduced in a very different way in [Wa44].

As in [KN06], we can write the heat kernel on $\mathbb{Z}/n\mathbb{Z}$ in two ways, once through its spectral expansion and once by periodizing the heat kernel on $\mathbb{Z}$, also known as the method of images. Since the heat kernel is unique, the two expressions are equal. As such, we obtain a discrete analogue of the Poisson summation formula, namely that

$$\sum_{j=-\infty}^{\infty} \mathrm{e}^{-2t} I_{x+jn}(2t) = \frac{1}{n} \sum_{k=0}^{n-1} \mathrm{e}^{-4\sin^2(k\pi/n)t} \mathrm{e}^{2\pi i k x/n}. \tag{25}$$

We refer to [ADG02] for a derivation when $x = 0$ without the heat kernel perspective. In the case when $n = 1$ and $x = 0$, (25) gives that

$$\sum_{j=-\infty}^{\infty} \mathrm{e}^{-2t} I_j(2t) = 1,$$

thus reproving (22). That is, the consideration of heat diffusion on the space consisting of exactly one point gives rise to a nontrivial Bessel function summation identity.

We conclude this section with some examples of Bessel convolution identities. Define the convolution of two functions on $(0, \infty)$ by

$$f * g(x) = \int_0^x f(t)\, g(x-t) \mathrm{d}t.$$

Continuing to use the heat kernel formalism as above, we can deduce that

$$J_l * J_n(x) = (-1)^{\frac{l+n+1}{2}} \left(\cos x + J_0(x) - 2 \sum_{k=0}^{(l+n-1)/2} (-1)^k J_{2k}(x) \right)$$

provided the sum $l + n$ of the positive integers l and n is odd; see [K12] for details. For example, with $l = 0$ and $n = 1$, one gets that

$$\int_0^x J_0(t) J_1(x-t)\, \mathrm{d}t = J_0(x) - \cos(x).$$

Further integral formulas are deduced in [K12] such as

$$\int_0^x J_{2l+1}(t)\, \mathrm{d}t = 1 - J_0(x) - 2 \sum_{k=1}^{l} J_{2k}(x).$$

As another example, for positive integers n and m, one has that

$$I_{n+m}(t) = \sum_{k=0}^{\infty} \frac{(-1)^k}{2^{k+1}} \left(I_1^{*k} * I_{n-1} * I_m \right)(t),$$

where $*k$ denotes the k-fold convolution. This identity is proved in [CJKS24], to which we refer for further discussion and context.

5. Application 2: finite trigonometric sums

Finite sums of powers of trigonometric functions of multiple angles, such as

$$\sum_{j=1}^{m-1} \frac{e^{2\pi i rj/m}}{\sin^n(j\pi/m)} \quad \text{for} m \in \mathbb{N} \text{ and } r \in \{0,1,\ldots,m-1\}$$

appear in many contexts, including Verlinde formulas, chiral Potts models, expressions for resistance in electrical network, and modeling angles in proteins and circular genomes; see [Do92, JKS23] for references. Let us now describe the methodology developed in [JKS23] to evaluate these sums, which is based on analysis of the resolvent kernel as derived from the heat kernel on the discrete torus.

For r and m as above and $\beta \in \mathbb{R} \setminus \mathbb{Z}$, define

$$C_{m,r}(\beta,n) = \frac{1}{m}\sum_{j=0}^{m-1} \frac{1}{\sin^{2n}((j+\beta)\pi/m)} e^{2\pi irj/m},$$

as well as the generating function

$$f(s,\beta,r) := \sum_{n=0}^{\infty} C_{m,r}(\beta,n+1)\, s^n \tag{26}$$

of the sequence $\{C_{m,r}(\beta,n)\}_{n=1}^{\infty}$. As shown in [JKS23], the series (26) is related to the resolvent kernel, or the Green's function, with an additive twist on the discrete circle $\mathbb{Z}/m\mathbb{Z}$ when viewed as a Cayley graph C_m of degree two. Specifically, from the spectral theory of the circle graph C_m, the resolvent kernel, evaluated at two points $x, y \in C_m$ such that $r = |x-y| \pmod m$ can be expressed as

$$G_{m,\beta}(r,s) = \frac{1}{m}\sum_{j=0}^{m-1} \frac{1}{s + 2\sin^2\left(\pi\frac{j+\beta}{m}\right)} \exp\left(2\pi i \frac{j+\beta}{m} r\right),$$

for all complex s for which the right-hand side is well-defined. Hence

$$f(s,\beta,r) = 2G_{m,\beta}(r,-2s)\, e^{-2\pi i\beta r/m}.$$

On the other hand, the twisted resolvent kernel, for sufficiently large $\mathrm{Re}(s)$ is the Laplace transform of the twisted heat kernel, which, by the method of imaging, can be expressed as a rapidly convergent sum of discrete Gaussians $e^{-t}I_x(t)$, namely that

$$K_{X_m,\chi_\beta}(r;t) = \sum_{k\in\mathbb{Z}} e^{-2\pi i\beta k} e^{-t} I_{r+km}(t).$$

Note: In [JKS23] the authors employed a different normalization of the heat kernel than the one given in (10). In effect, the edge-difference is given weight $1/2$ rather than 1, resulting in a different time scale in the heat kernel. To be consistent with the formulas in [JKS23], we will follow the normalization employed there.

By computing the Laplace transform of the heat kernel (5), the following result is proved in [JKS23].

Theorem 6. *([JKS23]) For sufficiently small complex s,*

$$\sum_{n=0}^{\infty} C_{m,r}(\beta, n+1)\, s^n = 2\mathrm{e}^{-2\pi i\beta r/m} \frac{U_{m-r-1}(1-2s) + \mathrm{e}^{2\pi i\beta} U_{r-1}(1-2s)}{T_m(1-2s) - \cos 2\pi\beta},$$

where T_n and U_n denote the Chebyshev polynomials of the first and second kind, with the convention that $U_{-1}(1-2s) \equiv 0$.

The right-hand-side of the identity in theorem 6 is a rational function in s with well-known coefficients. Therefore, one can obtain an evaluation for each of the coefficients in the right-hand-side. For example, as shown in [JKS23], for any $k > 0$ we have that

$$\sum_{j=1}^{3k-1} \frac{1}{\sin^4(j\pi/3k)} \cos(2\pi j/3) = -\frac{1}{45}\left(39k^4 + 30k^2 + 11\right),$$

which, as far as we know, is a new evaluation of the above classical trigonometric sum.

Previous results of this type have been studied by many authors; see the references in [Do92, JKS23] for a start. We find our approach, ultimately arising from the discrete Gausssian, to be structurally pleasing, and it also gives new and general formulas for the exact evaluation of sums of this type.

To illustrate the universality of this idea, if one starts instead with the untwisted discrete Gaussian, one obtains in the same way (see [K12]) the celebrated special values

$$\zeta(k) = \sum_{n=1}^{\infty} \frac{1}{n^k}$$

for k even, such as $\zeta(2) = \pi^2/6$, $\zeta(4) = \pi^4/90$ *etc* due to Euler. Furthermore, in [JKS23] we were able to answer a question posed in [XZZ22] about certain related trigonometric sums, that, thanks to the intriguing formulas in [XZZ22] linking the discrete and continuous, provides a new approach to obtain special values of Dirichlet L-functions.

6. Application 3: asymptotics by passing from the discrete to the continuous

Consider a discrete tori in d dimensions, or equivalently, a rectangular box in the standard lattice $\mathbb{Z}^d$ equipped with periodic boundary conditions, see figure 2. The question is how does the determinant of the (finite) graph Laplacian grows as the rectangular box becomes larger. This invariant counts the number of rooted spanning trees. (We note that arbitrary tori can also be treated; see [CJK12].) In thermodynamics one speaks of the infinite volume limit and the continuum limit. These are both visible in our asymptotics which extends what was known in dimension 2 by work of Kasteleyn, Barber, Duplantier-David [Ka61, Ba70, DD88]. In short, among the key ideas in our approach are, first, to use a discrete version of the Poisson summation formula in order to encode the spectral data of the graph in terms of discrete Gaussians, then second, in the asymptotic analysis to relate the discrete to the continuous via proposition 1 above. The following is one of the main results from [CJK10, CJK12].

Theorem 7. *([CJK10, CJK12]) Let A_n be a sequence of integral $d \times d$ matrices with $\det A_n \to \infty$ and $A_n/(\det A_n)^{1/d} \to A \in \mathrm{SL}(d,\mathbb{R})$. As $n \to \infty$,*

$$\log\det \Delta_{\mathbb{Z}^d/A_n\mathbb{Z}^d} = \log\det \Delta_{\mathbb{Z}^d} \det A_n + \frac{2}{d}\log\det A_n + \log\det \Delta_{\mathbb{R}^d/A\mathbb{Z}^d} + o(1).$$



Figure 2. Spectral asymptotics of a discretized torus as the mesh goes to zero.

In the above theorem, $\Delta_{\mathbb{Z}^d/A_n\mathbb{Z}^d}$ is the Laplacian on the discrete torus; $\Delta_{\mathbb{Z}^d}$ is the Laplacian on the lattice $\mathbb{Z}^d$ and $\Delta_{\mathbb{R}^d/A\mathbb{Z}^d}$ is the Laplacian on the (continuous) torus $\mathbb{R}^d/A\mathbb{Z}^d$. The determinant of the Laplacian on the discrete torus is a finite determinant, in the sense of linear algebra. However, the determinant of the Laplacian on the continuous torus is obtained through zeta regularization, which adds an entirely new level of complexity to the matter.

Let us briefly explain how the discrete Gaussian comes into play in this setting. Following the usual formalism, $\log\det\Delta_{\mathbb{Z}^d/A_n\mathbb{Z}^d}$ is related to the Mellin transform of the trace of the heat kernel. The trace of the heat kernel is the theta function, which in our discrete setting involves a sum of products of discrete Gaussians $\prod_{j=1}^{d} \mathrm{e}^{-2t} I_{n_j k_j}(2t)$. A careful analysis of the asymptotics of the discrete Gaussian is the key ingredient in the proof of theorem 7.

Theorem 7 is of interest in combinatorics, statistical physics and quantum field theory (QFT). Several recent papers in mathematical physics, notably [HK20, IK22, Gr23], have extended theorem 7. A problem is raised in [RV15] in connection with their approach to a combinatorial QFT which theorem 7 addresses just a part, aside from the 2D case that was already known as just mentioned.

We point out that in higher dimensions $d > 2$ the lead terms in the asymptotics of theorem 7 were established in [SW00, SS01, Ly05]; see also their bibliographies for further references. In particular, the formula of Sokal-Starinets involves the I-Bessel function in the same way as in [CJK10], but Sokal-Starinets have another way of interpreting the appearance of the I-Bessel, presumably not as a discrete Gaussian. Detailed asymptotics for more general lattice domains in 2D are obtained in the work of Kenyon, see [Ke00].

For other contexts in physics where recent similar considerations of discrete heat kernels appear, see [KS18, KS23].

7. Spectral zeta functions

7.1. Spectral zeta function of $\mathbb{Z}$

The zeta function regularization method in physics started in the 1970 s with papers by Dowker-Critchley and by Hawking. For a survey of applications to mathematical physics of this procedure, see [V03, El12, DEK12]. The zeta function in question is defined by a integral transform, specifically the Mellin transform, of the heat kernel. The case of the circle essentially gives rise to the Riemann zeta function. It is much less standard is to take the Mellin integral transform of $\mathrm{e}^{-2t} I_n(2t)$, such as

$$\zeta_{\mathbb{Z}}(s) = \frac{1}{\Gamma(s)} \int_0^{\infty} \mathrm{e}^{-2t} I_0(2t)\, t^s \frac{\mathrm{d}t}{t},$$

for $0 < \mathrm{Re(s)} < 1/2$ and then extend by meromorphic continuation, as was done in [FK17]. Is this function also relevant for physics? There are some indications that this the case. First, using the method of [CJK10] the asymptotics of the spectral zeta function of discrete tori were determined in [FK17]. Note that in the zeta function regularization method the determinant of the Laplacian discussed above corresponds to the special value $\zeta'(0)$ of the corresponding zeta function.

Second, here is a different context motivated by understanding the Casimir energy in QFT. The paper [NP00] compared the spectral zeta functions of a circular disk, denote by $\zeta_{\mathrm{cir}}(s)$, and that of the solid cylinder, denote by $\zeta_{\mathrm{cyl}}(s+1)$, and obtained the following relationship:

$$\zeta_{\mathrm{cir}}(s) = 2\sqrt{\pi}\frac{\Gamma\left((s+1)/2\right)}{\Gamma(s/2)}\zeta_{\mathrm{cyl}}(s+1).$$

The factor appearing here is the same independently of whether one considers Dirichlet or Neumann boundary conditions in the circular disk and solid cylinder.

It turns out from calculations in [FK17, Du19] that

$$\zeta_{\mathbb{Z}}(s) = \frac{1}{\sqrt{\pi}4^s}\frac{\Gamma(1/2-s)}{\Gamma(1-s)} = \begin{pmatrix} -2s \\ -s \end{pmatrix},$$

which essentially is an extension of the Catalan numbers. Therefore, the factor appearing in [NP00]

$$\frac{2\sqrt{\pi}\Gamma\left((1+s)/2\right)}{\Gamma(s/2)}$$

equals

$$\frac{2^s}{\zeta_{\mathbb{Z}}\left((1-s)/2\right)}.$$

Note that, via the functional symmetry given in [FK17], $\zeta_{\mathbb{Z}}((1-s)/2)$ is essentially equal to $\zeta_{\mathbb{Z}}(s/2)$ similarly to what is the case for the Riemann zeta function. More precisely, with $\xi_{\mathbb{Z}}(s) := 2^s\cos(\pi s/2)\zeta_{\mathbb{Z}}(s/2)$, it holds that

$$\xi_{\mathbb{Z}}(s) = \xi_{\mathbb{Z}}(1-s)$$

for all $s \in \mathbb{C}$. Is there a spectral explanation for the appearance of $\zeta_{\mathbb{Z}}(s)$ in [NP00] and if so, how does it generalize?

Finally, we mention the paper [KP23] which argues that special values of $\zeta_{\mathbb{Z}}(s)$ are related to the volume of spheres.

7.2. Application 4: spectral zeta function of regular trees and hypergeometric identities

The spectral zeta function $\zeta_{T_{q+1}}$ on the $(q+1)$-regular tree T_{q+1} was computed in [FK17, theorem 1.4] by representing it as an integral over the spectral measure on T_{q+1}. Note that the case $q=1$ was discussed in the previous subsection. It was proved that for $q>1$ one has that

$$\zeta_{T_{q+1}}(s) = \frac{q(q+1)}{(q-1)^2\left(\sqrt{q}-1\right)^{2s}}F_1\left(\frac{3}{2}, s+1, 1, 3; -\frac{4\sqrt{q}}{\left(\sqrt{q}-1\right)^2}, \frac{4\sqrt{q}}{\left(\sqrt{q}+1\right)^2}\right). \tag{27}$$

On a different front, the spectral zeta function can be expressed in terms of a Mellin transform of the heat kernel on T_{q+1}. Namely, for $0 < \mathrm{Re}(s) < 1/2$ we have that

$$\zeta_{t_{q+1}}(s) = \frac{1}{\Gamma(s)} \int_0^\infty K(t,0)\, t^s \frac{\mathrm{d}t}{t},$$

where $K(t,r)$ is the heat kernel on the tree T_{q+1}, as defined by (11). The Mellin transform of the rescaled discrete Gaussian (12) that appears in (11) can be evaluated, for any $n \geqslant 0$ and $0 < \mathrm{Re}(s) < 1/2$ as follows. For any integer $n \geqslant 0$, we can write that

$$\mathcal{M}\left(q^{-n/2}\mathrm{e}^{-(q+1)t} I_n\left(2\sqrt{q}t\right)\right)(s) = q^{-n/2}\frac{1}{\pi}\int_0^\pi \left(\int_0^\infty \mathrm{e}^{-\left(1-2\sqrt{q}\cos\theta+q\right)t} t^{s-1}\mathrm{d}t\right)\cos(n\theta)\,\mathrm{d}\theta.$$

where the application of Fubini–Tonelli theorem is justified by assumptions on s and the fact that $1 - 2\sqrt{q}\cos\theta + q > 0$ for $q > 1$. Therefore,

$$\mathcal{M}\left(q^{-n/2}\mathrm{e}^{-(q+1)t} I_n\left(2\sqrt{q}t\right)\right)(s) = q^{-n/2}\frac{1}{2\pi}\int_{-\pi}^\pi \frac{\cos(n\theta)}{\left(1-2\sqrt{q}\cos\theta+q\right)^s}\mathrm{d}\theta.$$

Applying a simple manipulation of the above integral in combination with [GR07], formula 9.112 (defining the Gauss hypergeometric function F) one gets that

$$\mathcal{M}\left(q^{-n/2}\mathrm{e}^{-(q+1)t} I_n\left(2\sqrt{q}t\right)\right)(s) = q^{-n-s}\frac{\Gamma(n+s)}{\Gamma(n+1)} F\left(s, s+n; n+1; \frac{1}{q}\right).$$

Therefore, the zeta function on T_{q+1} can be expressed as

$$\zeta_{T_{q+1}}(s) = q^{-s} F\left(s,s,1;\frac{1}{q}\right) - (q-1)\sum_{j=1}^\infty q^{-s-2j}\frac{\Gamma(s+2j)}{\Gamma(s)\Gamma(2j+1)} F\left(s, s+2j; 2j+1; \frac{1}{q}\right), \tag{28}$$

where the termwise integration of the series on the right-hand side of (11) is justified by exponential decay of the building blocks (12). When $q = 1$, then (28) becomes

$$\zeta_{T_2}(s) = F(s,s,1;1) = \frac{\Gamma(1-2s)}{\Gamma(1-s)^2} = \zeta_{\mathbb{Z}}(s).$$

Comparing (27) with (28) yields a new identity for the Appell hypergeometric function F_1.

Moreover, the expression (28) is suitable for computing special values of the spectral zeta function at zero and negative integers. We have the following proposition.

Theorem 8. *For any $q \geqslant 1$, we have that*

$$\zeta_{T_{q+1}}(0) = 1 \quad \textit{and} \quad \zeta'_{T_{q+1}}(0) = \begin{cases} \frac{q-1}{2}\log\left(1-q^{-2}\right), & \textit{when } q > 1 \\ 0, & \textit{when } q = 1. \end{cases}$$

Also, $\zeta_{T_{q+1}}(-1) = 1 + q$. Furthermore, for integers $m \geqslant 2$, we have that

$$\zeta_{T_{q+1}}(-m) = \sum_{k=0}^m \binom{m}{k}^2 q^{m-k} - (q-1)\sum_{j=1}^{\lfloor m/2\rfloor}\sum_{k=0}^{m-2j}\binom{m}{k}\binom{m}{2j+k} q^{m-2j-k}. \tag{29}$$

Before we proceed with the proof, let us note an interesting curiosity. It is immediate from (29) that the values of $\zeta_{T_{q+1}}$ at negative integers $-m$ and are degree m polynomials in q. This is reminiscent of the property of the Hurwitz zeta function

$$\zeta_H(z,q) := \sum_{n=0}^{\infty} \frac{1}{(n+q)^z}, \quad \mathrm{Re}(z) > 1$$

that the values of its meromorphic continuation at negative integers $-m$ are $\zeta_H(-m,q) = -\frac{B_{m+1}(q)}{m+1}$, where B_{m+1} are the Bernoulli polynomials, which are degree $m+1$ polynomials in q.

Proof. Fix $q > 0$. In order to evaluate $\zeta_{T_{q+1}}$ and its derivative at $s = 0$ we use the following asymptotic expressions:

$$F\left(s,s,1;\frac{1}{q}\right) = 1 + O\left(s^2\right) \quad \text{as } s \to 0,$$

$$\frac{\Gamma(s+2j)}{\Gamma(s)\Gamma(2j+1)} = \frac{s}{2j} + O\left(s^2\right) \quad \text{as } s \to 0,$$

and

$$q^{-s-2j} F\left(s,s+2j;2j+1;\frac{1}{q}\right) = q^{-2j}\left(1 + O(s)\right) \quad \text{as } s \to 0.$$

From (28) we get the following asymptotic expansion as $s \to 0$:

$$\zeta_{T_{q+1}}(s) = 1 - (q-1)s \sum_{j=1}^{\infty} \frac{q^{-2j}}{2j} + O\left(s^2\right),$$

which yields that $\zeta_{T_{q+1}}(0) = 1$. When $q = 1$, trivially $\zeta'_{T_2}(0) = \left(\frac{\Gamma(1-2s)}{\Gamma(1-s)^2}\right)'\Big|_{s=0} = 0$, while for $q > 1$ we get

$$\zeta'_{T_{q+1}}(0) = -\frac{q-1}{2} \sum_{j=1}^{\infty} \frac{\left(q^{-2}\right)^j}{j} = \frac{q-1}{2} \log\left(1 - q^{-2}\right).$$

The evaluation of $\zeta_{T_{q+1}}(-1)$ is immediate because

$$\frac{\Gamma(s+2j)}{\Gamma(s)\Gamma(2j+1)}\bigg|_{s=-1} = 0 \quad \text{for all integers } j \geqslant 1.$$

Hence,

$$\zeta_{T_{q+1}}(-1) = qF\left(-1,-1;1;\frac{1}{q}\right) = 1 + q.$$

Finally, let $m \geqslant 2$. Then, for any $1 \leqslant j \leqslant \lfloor m/2 \rfloor$ we have that

$$\frac{\Gamma(s+2j)}{\Gamma(s)\Gamma(2j+1)}\bigg|_{s=-m} = \lim_{z \to 0} \frac{\Gamma(z+2j-m)}{\Gamma(z-m)\Gamma(2j+1)} = \frac{m!}{(2j)!\,(m-2j)!}.$$

Also, for $2j > m$ we have that

$$\left.\frac{\Gamma(s+2j)}{\Gamma(s)\Gamma(2j+1)}\right|_{s=-m} = 0.$$

Therefore,

$$\zeta_{T_{q+1}}(-m) = q^m F\left(-m,-m,1;\frac{1}{q}\right) - (q-1)\sum_{j=1}^{\lfloor m/2\rfloor}\frac{m!q^{m-2j}}{(2j)!(m-2j)!}F\left(-m,2j-m;2j+1;\frac{1}{q}\right). \tag{30}$$

From the definition of the hypergeometric function, it is trivial to deduce that

$$F\left(-m,-m,1;\frac{1}{q}\right) = \sum_{k=0}^{m}\binom{m}{k}^2 q^{-k}$$

and that

$$\frac{m!}{(2j)!(m-2j)!}F\left(-m,2j-m;2j+1;\frac{1}{q}\right) = \sum_{k=0}^{m-2j}\binom{m}{k}\binom{m}{2j+k}q^{-k}.$$

Substituting these evaluations into (30) proves (29). □

8. Potential applications to differential privacy

Differential privacy is a mathematically rigorous criterion by which an algorithm that acts on a dataset can be assessed. We refer to the article [CKS22] and references therein for an excellent presentation. In practice, as described in [CKS22], one alters the output of a $\mathbb{Z}$-valued algorithm $\mathcal{M}$ which produces results from a given dataset $\mathcal{D}$ by adding a random amount. The main results of [CKS22] are that one can use a discretized Gaussian random (6) (using our terminology) to create algorithms which satisfy differential privacy, and that one can effectively and efficiently sample from a discretized Gaussian random variable.

It would be interesting to determine if the discrete Gaussian (3) also satisfies the criterion established in [CKS22] for random 'noise' as required by differential privacy. It should be noted that other authors have pointed out the potential usefulness of (3) and distinguished the difference between (3) and (6); see section 2.6 of [Li24] who refers to earlier work by the same author in [Li90]. One can reasonably expect that, indeed, (3) will fulfill the conditions needed for differential privacy, if for no reason other than the triangle inequality, the results from [CKS22] and Proposition 1. However, it would be better to establish differential privacy associated to (3) without appealing to [CKS22]. We will leave this study for a later time.

From the point of view of probability, one can state several advantages that (3) has over (1). First, we note that the (quite impressive) analysis of [CKS22] includes, for example, estimates for the moment generating function and variance of the discretized Gaussian (6); see lemma 8 and corollary 9 of [CKS22]. By contrast, note that the characteristic function of (3) is explicitly computed; see (16). As such, one has elementary closed-form expressions for all moments of (3). See also [AGMP24]. Second, the sum of two independent discretized Gaussian random variables (6) is close to, but not equal to, a discretized Gaussian random variable; see the

comment after remark 3.7 in [AA19] and theorem 1.1 of [AR24]. By comparison, the sum of two independent discrete Gaussians (3) is a discrete Gaussian, and the convolution of the probability density functions is, in effect, computed in the above-cited Chapman–Kolmogorov equation; see section 4. Finally, let us point out that one application of theorem 5 is a means by which one can sample from a discrete Gaussian is a manner similar to a sampling method for a continuous Gaussian random variable. In doing so, one has, at least the beginning of, an analogue to the sampling algorithms developed in [CKS22]. Other sampling methods could be developed based on numerical evaluations of Bessel functions; to this end, we found the articles [Am74] and [GS81] to be interesting.

Finally, let us note that the articles [AA19] and [AR24] consider discretized Gaussian random variables on any d-dimensional lattice Λ in $\mathbb{R}^d$. Certainly, our point of view is amenable to such a setting; see, for example, [CJK10] and more generally [CJK12]. The probability density function is simply the heat kernel on Λ, and we assert that one can prove an analogue of theorem 5. Again, these considerations will be undertaken elsewhere.

Data availability statement

No new data were created or analysed in this study.

ORCID iDs

Gautam Chinta https://orcid.org/0000-0002-2672-3819
Jay Jorgenson https://orcid.org/0000-0002-9765-0944
Anders Karlsson https://orcid.org/0000-0002-1886-6780
Lejla Smajlović https://orcid.org/0000-0002-2709-5535

References

[AGMP24] Abadías L, Gonzalez-Camus J, Miana P and Pozo J 2021 Large time behaviour for the heat equation on $\mathbb{Z}$, moments and decay rates *J. Math. Anal. Appl.* **500** 125137
[AR24] Aggarwal D and Regev O A note on discrete Gaussian combinations of lattice vectors (arXiv:1308.2405)
[AA19] Agostini D and Amendola C 2019 Discrete Gaussian distributions via theta functions *SIAM J. Appl. Algebra Geom.* **3** 1–30
[Am74] Amos D E 1974 Computation of modified Bessel functions and their ratios *Math. Comp.* **28** 239–51
[An58] Anderson P W 1958 Absence of diffusion in certain random lattices *Phys. Rev.* **109** 1492–505
[ADG02] Al-Jarrah A, Dempsey K M and Glasser M L 2002 Generalized series of Bessel functions *J. Comput. Appl. Math.* **143** 1–8
[At87] Athreya K B 1987 Modified Bessel function asymptotics via probability *Stat. Probab. Lett.* **5** 325–7
[Ba70] Barber M N 1970 Asymptotic results for self-avoiding walks on a manhattan lattice *Physica* **48** 237–41
[Ba49] Bateman H 1943 Some simple differential difference equations and the related functions *Bull. Am. Math. Soc* **49** 494–512
[CKS22] Canonne C, Kamath G and Steinke T 2022 The discrete Gaussian for differential privacy *J. Privacy Confidentiality* **12** 784
[Ch84] Chavel I 1984 *Eigenvalues in Riemannian Geometry* (Academic)
[CJK10] Chinta G, Jorgenson J and Karlsson A 2010 Zeta functions, heat kernels and spectral asymptotics on degenerating families of discrete tori *Nagoya Math. J.* **198** 121–72

[CJK12] Chinta G, Jorgenson J and Karlsson A 2012 Complexity and heights of tori. Dynamical systems and group actions *Contemp. Math.* vol 567 (American Mathematical Society) pp 89–98
[CJK15] Chinta G, Jorgenson J and Karlsson A 2015 Heat kernels on regular graphs and generalized Ihara zeta function formulas Monatsh. *Math.* **178** 171–90
[CJKS24] Chinta G, Jorgenson J, Karlsson A and Smajlović L 2024 The parametrix construction of the heat kernel on a graph *Complex Anal. Oper. Theory* **18** 176
[CY99] Chung F and Yau S-T 1999 Coverings, heat kernels and spanning trees *Electron. J. Combin.* **6** 21
[CMS00] Cowling M, Meda S and Setti A 2000 Estimates for functions of the Laplace operator on homogeneous trees *Trans. Am. Math. Soc.* **352** 4271–93
[DH13] Decrouez G and Hall P 2013 Normal approximation and smoothness for sums of means of lattice-valued random variables *Bernoulli* **19** 1268–93
[Do92] Dowker J S 1992 On Verlinde's formula for the dimensions of vector bundles on moduli spaces *J. Phys.* A **25** 2641–8
[DEK12] Dowker F, Elizalde E and Kirsten K 2012 Applications of zeta functions and other spectral functions in mathematics and physics: a special issue in honour of Stuart Dowker's 75th birthday *J. Phys. A* **45** 370301
[Du19] Dubout J Spectral zeta functions of graphs, their symmetries and extended Catalan numbers (arXiv:1909.01659)
[DD88] Duplanticr B and David F 1988 Exact partition functions and correlation functions of multiple Hamiltonian walks on the Manhattan lattice *J. Stat. Phys.* **51** 327–434
[El12] Elizalde E 2012 Ten physical applications of spectral zeta functions *Lecture Notes in Physics* vol 855, 2nd edn (Springer)
[Fe66] Feller W 1966 Infinitely divisible distributions and Bessel functions associated with random walks *SIAM J. Appl. Math.* **14** 864–75
[Fe70] Feller W 1970 An Introduction to Probability Theory and Its Applications *Vol II*, 2dn edn (Wiley)
[FK17] Friedli F and Karlsson A 2017 Spectral zeta functions of graphs and the Riemann zeta function in the critical strip *Tohoku Math. J.* **69** 585–610
[GS81] Gatto M A and Seery J B 1981 Numerical evaluation of the modified Bessel functions *I* and *K Comput. Math. Appl.* **7** 203–9
[GK24] Gavalakis L and Kontoyiannis I 2024 Entropy and the discrete central limit theorem *Stochastic Process. Appl.* **170** 10
[GR07] Gradshteyn I S and Ryzhik I M 2007 *table of Integrals, Series and Products* (Elsevier Academic)
[Gr23] Greenblatt R L 2023 Discrete and zeta-regularized determinants of the Laplacian on polygonal domains with Dirichlet boundary conditions *J. Math. Phys.* **64** 043301
[GN98] Grigor'yan A and Noguchi M 1998 The heat kernel on hyperbolic space *Bull. London Math. Soc.* **30** 643–50
[GI02] Grünbaum F A and Iliev P 2002 Heat kernel expansions on the integers *Math. Phys. Anal. Geom* **5** 183–200
[HK20] Hou Y and Kandel S 2020 Asymptotic analysis of determinant of discrete Laplacian *Lett. Math. Phys.* **110** 259–96
[IL71] Ibragimov I A Yu. I A and Linnik V 1971 *Independent and Stationary Sequences of Random Variables* (Wolters-Noordhoff)
[IK22] Izyurov K and Khristoforov M 2022 Asymptotics of the determinant of discrete Laplacians on triangulated and quadrangulated surfaces *Commun. Math. Phys.* **394** 531–72
[JKS23] Jorgenson J, Karlsson A and Smajlović L 2024 The resolvent kernel on the discrete circle and twisted cosecant sums *J. Math. Anal. Appl.* **538** 23
[KS18] Kan N and Shiraishi K 2018 Free energy on a cycle graph and trigonometric deformation of heat kernel traces on odd spheres *J. Phys.* A **51** 23
[KS23] Kan N and Shiraishi K 2023 Discrete time heat kernel and UV modified propagators with dimensional deconstruction *J. Phys.* A **56** 16
[K12] Karlsson A 2012 Applications of Heat Kernels on Abelian Groups: $\zeta z(2n)$, Quadratic Reciprocity, Bessel Integrals *Number Theory, Analysis and Geometry* (Springer) pp 307–20

[KN06] Karlsson A and Neuhauser M 2006 Heat kernels, theta identities and zeta functions on cyclic groups *Topological and Asymptotic Aspects of Group Theory* vol 394 (American Mathematical Society) pp 177–89

[KP23] Karlsson A and Pallich M 2023 Volumes of spheres and special values of zeta functions of $\mathbb{Z}$ and $\mathbb{Z}/n\mathbb{Z}$ *Acta Arith.* **208** 161–70

[Ka61] Kasteleyn P W 1961 The statistics of dimers on a lattice, I. The number of dimer arrangements on a quadratic lattice *Physica* **27** 1209–25

[Ke00] Kenyon R 2000 Long-range properties of spanning trees *J. Math. Phys.* **41** 1338–63

[Li90] Lindeberg T 1990 Scale-space for discrete signals *IEEE Trans. Pattern Anal. Mach. Intell.* **12** 234–54

[Li24] Lindeberg T Discrete approximations of Gaussian smoothing and Gaussian derivatives (arXiv:2311.11317)

[Ly05] Lyons R 2005 Asymptotic enumeration of spanning trees *Combin. Probab. Comput.* **14** 491–522

[Mu83] Mumford D 1983 Tata Lectures on Theta I, With the Assistance *Progr. Math.* vol 28 C Musili, M Nori, E Previato and M Stillman (Birkhäuser Boston, Inc) p xiii+ 235

[NP00] Nesterenko V V and Pirozhenko I G 2000 Spectral zeta functions for a cylinder and a circle *J. Math. Phys.* **41** 4521–31

[Pa93] Pang M M H 1993 Heat kernels of graphs *J. London Math. Soc.* **47** 50–64

[Pe75] Petrov V V 1975 *Sums of Independent Random Variables* (Springer)

[RV15] Reshetikhin N and Vertman B 2015 Combinatorial quantum field theory and gluing formula for determinants *Lett. Math. Phys.* **105** 309–40

[SW00] Shrock R and Wu F Y 2000 Spanning trees on graphs in d dimensions *J. Phys. A* **33** 3881–902

[SS01] Sokal A D and Starinets A O 2001 Pathologies of the large-N limit for RP^{N-1}, CP^{N-1}, QP^{N-1} and mixed isovector/isotensor σ-models *Nucl. Phys.* B **601** 425–502

[SW23] Szewczak Z and Weber M 2023 Classical and almost sure local limit theorems *Dissertationes Math.* **97** 589

[V03] Vassilevich D V 2003 Heat kernel expansion: user's manual *Phys. Rep.* **388** 279

[Wa44] Watson G N 1995 *A Treatise on the Theory of Bessel Functions* 2nd edn (Cambridge University Press) p viii+804

[XZZ22] Xie B, Zhao Y and Zhao Y 2024 Special values of spectral zeta functions of graphs and Dirichlet L-functions *J. Number Theory* **256** 136–59