

Mental health status and quality of life after an acute myocardial infarction with non-obstructive coronary arteries or takotsubo syndrome: a systematic review

Philip Leissner ^{1*}, Erik M.G. Olsson ¹, Elisabet Rondung ², Runa Sundelin ³, Jonas Spaak ⁴, Anders Ulvenstam ⁵, Anna Nordenskjöld⁶, Lena Kövamees⁷, Patrik Lyngå ³, Claes Held ^{8,9}, Per Tornvall ³, and Sophia Humphries ¹⁰

¹Department of Women's and Children's Health, Uppsala University, MTC-huset, Dag Hammarskjölds väg 14B, Akademiska sjukhuset, Uppsala 752 37, Sweden; ²Department of Psychology and Social Work, Mid Sweden University, Östersund 831 25, Sweden; ³Department of Clinical Science and Education Södersjukhuset, Karolinska Institutet, Stockholm, Sweden; ⁴Department of Clinical Sciences, Danderyd Hospital, Karolinska Institutet, Stockholm, Sweden; ⁵Department of Public Health and Clinical Medicine, Umeå University, Umeå, Sweden; ⁶Department of Cardiology, Faculty of Medicine and Health, Örebro University, Örebro, Sweden; ⁷Patient Representative, Swedish Heart and Lung Association, Stockholm, Sweden; ⁸Uppsala Clinical Research Center, Uppsala University, Uppsala, Sweden; ⁹Department of Medical Sciences, Cardiology, Uppsala Clinical Research Center, Uppsala University, Uppsala, Sweden; and ¹⁰Department of Neurobiology, Care Science and Society, Karolinska Institute, Stockholm, Sweden

Received 13 November 2024; revised 6 January 2025; accepted 16 February 2025; online publish-ahead-of-print 25 February 2025

Aims

Myocardial infarction with non-obstructive coronary arteries (MINOCA) and takotsubo syndrome (TS) are both characterized by lack of significant coronary artery stenoses and a higher prevalence of mental health disorders preceding the event. Currently, little is known about their pathological aetiologies and subsequent treatment plans, giving cause for concern among those affected. The objective of this review is to provide a comprehensive overview of mental health status and quality of life (QoL) in MINOCA and TS patients after the acute event, compared with both cardiac and non-cardiac populations, and over time.

Methods and results

A systematic search was conducted via Cochrane Library, CINAHL, PsycINFO, PubMed, ASSIA, Web of Science, Scopus, and Embase from inception to May 2024. The review was registered in PROSPERO and methods, and results were reported in accordance with the PRISMA guidelines. Quality assessment and risk of bias were evaluated using the Newcastle–Ottawa Scale for cross-sectional and cohort studies. Sample sizes ranged from 13 to 5322 participants. The risk of bias was high in 18/28, medium in 7/28, and low in 3/28 studies. Across the symptoms assessed, MINOCA and TS patients reported worse mental health status or QoL than non-cardiac groups in 10/13 studies and cardiac groups in 10/20 studies. Investigating change over time, 1/5 studies found deteriorating mental health status, 3/5 reported improved mental health status or QoL, and 1/5 reported no change in MINOCA and TS patients.

Conclusion

Patients with MINOCA or TS seem to have worse mental health status and QoL after the acute event than non-cardiac individuals, but it is yet difficult to conclude whether mental distress and QoL are equal or worse compared with coronary heart disease patients. There is no convincing evidence that mental health status or QoL of MINOCA and TS patients naturally improve over time after the acute event. Among the studies evaluated, risk of bias was high. More high-quality studies are needed, investigating mental health status and QoL among MINOCA and TS patients.

* Corresponding author. Tel: +46 734697318, Email: philip.leissner@uu.se

© The Author(s) 2025. Published by Oxford University Press on behalf of the European Society of Cardiology.


This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Lay summary

This is a systematic review of all papers reporting data on post-event mental health status and quality of life in individuals following either a MINOCA or takotsubo syndrome.

- Individuals who have suffered a MINOCA or takotsubo syndrome more often report worse mental health status and quality of life than healthy controls and at least as often or more often than other cardiac patients.
- Much of the available data had a high risk of bias, demonstrating a lack of high-quality investigation of the mental well-being of these patients, and more attention is warranted.

Graphical Abstract



Keywords

MINOCA • Takotsubo syndrome • Mental health • Quality of life • Systematic Review • After event

Introduction

Cardiovascular disease (CVD), including myocardial infarction (MI), continues to be a leading cause of morbidity and mortality worldwide.¹ For those surviving an MI, it can be a highly disturbing event, and symptoms of psychological distress are common.² Eighty-five years ago, MI with non-obstructive coronary arteries (MINOCA) was first noted,^{3,4} and over the last 20 years, it has been increasingly recognized by the frequent use of coronary angiography. However, the current understanding of MINOCA is limited, and the availability of evidence-based treatment is lacking.^{5,6} While this may cause further psychological suffering, little is still known regarding these patients' mental health status and quality of life (QoL), after the acute event.

MINOCA is defined as a 'working diagnosis' and may be caused by several underlying pathological mechanisms, presenting unique

diagnostic and therapeutic challenges.⁷ One pathophysiological explanation that was long classified as a cause of MINOCA is takotsubo syndrome (TS).⁷ While TS no longer is considered a type of MI,⁸ MINOCA and TS still share common clinical features and aetiologies.⁹ Compared with MI with obstructive coronary artery disease (MI-CAD), MINOCA and TS affect women to a greater extent, the patients are younger, and they have a higher prevalence of previous psychological distress or psychiatric illness.^{5,6,10} Additionally, TS is often reported as triggered by intense physical or emotional stress.⁵

Around 5–6% of all MI cases are categorized as MINOCA,⁶ and TS makes up around 2–3% of the cases presenting with a suspected acute coronary syndrome (ACS).⁵ Comparing MINOCA and TS, one study found that TS patients more often were women and older than the MINOCA patients and had a higher prevalence of mental health disorders and more in-hospital complications, but with a better long-term

prognosis.⁹ Although lacking obstructive CAD, both patient groups have a worse prognosis and more functional impairment than the general population.^{5,6}

Previous reviews have summarized somatic clinical features, mechanisms, treatments, and prognosis of MINOCA patients,^{11–13} but as of yet, there is no review of the mental health status or QoL of MINOCA patients after the acute event. Takotsubo syndrome has been reviewed regarding psychological factors before the acute event,^{14,15} but for mental health status after the acute event, there has only been a synthesis and review of case studies.¹⁶ The aim of this review is therefore to fill this knowledge gap and provide a comprehensive overview of mental health status and QoL in MINOCA and TS patients after the acute event, compared with both cardiac and non-cardiac populations, and over time.

Methods

This review was registered in the international Prospective Register of Systematic Reviews (PROSPERO 20202020 CRD42020160778). Methods and results were reported in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.¹⁷

Search strategy

An extensive systematic literature search was carried out in the following databases: the Cochrane Library, CINAHL, PsycINFO, PubMed, ASSIA, Web of Science, Scopus, and Embase from inception to May 2024. Search terms were tailored to each database and included terms for the exposure (MINOCA and Takotsubo) and outcomes (mental health status and QoL or associated symptoms). Search terms were based on Medical Subject Headings where available and complemented by free text searches. The full search strategy used for the CINAHL database is provided as an example in the appendix (see [Supplementary material online, Table S4](#)). The list of references from eligible papers was manually screened to additionally detect any potential studies not picked up by the database searches.

Study selection and eligibility criteria

Two authors independently screened the title and abstract of the retrieved studies, applying the same inclusion and exclusion criteria to determine study relevance. Full-text articles were assessed for eligibility after the initial screening. Disparities around inclusion criteria were resolved by discussion between the two reviewers and through consulting with a third independent reviewer, if necessary. Studies were eligible for inclusion if they (i) included adult patients diagnosed with MINOCA or TS or a diagnosis synonymous with MINOCA or TS; (ii) assessed the mental health status or QoL after the diagnosis; and (iii) utilized standardized and validated questionnaires or diagnostic interviews for assessment relating to the outcome of interest. Studies were excluded if they (i) were not written in English; (ii) were not peer-reviewed; (iii) did not have full-text availability; and (iv) followed case study design.

There is certain overlap between psychological states and psychological traits. As the purpose of this paper was to assess mental health status after the acute event, which may be due to the event itself, studies or results assessing psychological traits or personality were not included.

Data extraction

After full-text screening, data from the included studies were independently extracted by the two reviewers. This was done using an extraction form in which the following information was retrieved: general study characteristics (author, year, country), clinical characteristics (diagnosis, study group, sample size, time since event, control group characteristics), and outcomes (assessment tool or diagnosis). Quantitative data on severity (median or mean

scores) and frequency (prevalence or incidence) of adverse mental health outcomes were extracted for each group or time point, as well as the accompanying comparison between them (e.g. *P*-value, odds ratio, relative risk).

Quality appraisal

Risk of bias was assessed in all studies that matched inclusion and exclusion criteria with the Newcastle–Ottawa Scale (NOS) for cohort studies and an adapted version for cross-sectional studies which is modified accordingly. The NOS has been recommended for use in assessing studies on psychiatric conditions.¹⁸ The scales have three dimensions for which quality is assessed: selection, comparability, and outcome. Both versions have a maximum of four stars for selection and two stars for comparability. The cohort version has a maximum of three stars for outcome while the cross-sectional has two. It is possible to get a total of nine stars for cohort studies and eight stars for cross-sectional studies. The cut-off for high risk of bias is usually set at five stars or less,^{18,19} medium risk for bias at seven to six stars for cohort studies and six stars for cross-sectional, and low risk of bias at seven to eight for cross-sectional and eight to nine for cohort studies.¹⁹

Included studies may have followed case-control, cohort, or cross-sectional design. However, for the purpose of this systematic review, the studies were categorized based on our research question, i.e. with MINOCA/TS as exposure and mental health status or QoL as outcome. For example, a case-control study investigating differences in exposures between MINOCA patients and healthy controls, but that also collected data on mental health status after the diagnosis, would be described and assessed as a cross-sectional study; and a cohort study investigating cardiovascular outcomes, but with psychosocial measurements at baseline, would be considered a cross-sectional study if they had no additional psychosocial outcomes at follow-up.

Data analysis


The results of included studies are presented as a narrative synthesis, and no statistical comparisons or pooling of data was conducted. These results are presented in detail in [Supplementary material online, Table S1](#) and [Table S2](#). For the purpose of readability, the same results are presented synoptically in simplified tables and elaborated more comprehensibly in text. To make reading and comparing studies easier, some categorizations were made.

The included studies assessed mental health status and QoL at varying elapsed times since diagnosis of MINOCA or TS, as shown in [Supplementary material online, Table S1](#) and [Table S2](#). To maintain a more parsimonious language in the narrative synthesis of results and in the simplified tables, we categorized these as (i) acute phase, defined as assessment made during hospital admission and before discharge; (ii) short-term follow-up, defined as after discharge up to 3 months after the event; and (iii) long-term follow-up, defined as more than 3 months after the acute event.

Referring to QoL, some studies only reported a composite score, consisting of both mental and physical aspects of QoL, and other studies reported specific sub-scale scores. If only the composite was available, we reported this score, but if separate sub-scale scores were available, we only reported the ones pertaining to mental QoL.

MINOCA and TS were compared with various diagnoses related to coronary heart disease (CHD). These include MI, acute MI, ACS, coronary artery disease, and non-ST-elevation MI. The specific diagnoses are specified in the [Supplementary material online, Table S1](#) and [Table S2](#) and are referred to as CHD in the narrative synthesis of results and in the simplified tables.

Studies comparing MINOCA and TS with non-cardiac populations used varying matching methods and criteria, which are described in detail in the [Supplementary material online, Table S1](#) and [Table S2](#). However, in the narrative synthesis of results and simplified tables, these will be categorized as CVD risk controls, if matched on CVD risk factors, or healthy controls, if not matched on CVD risk factors.



In the narrative synthesis, results are categorized and presented under ordered headings. The first level of headings describes one of the following: (i) MINOCA patients, (ii) TS patients, or (iii) comparisons within and between MINOCA and TS patients. Under the headings of MINOCA and TS patients, the results are further categorized in groups of (i) change over time within the group, (ii) change over time compared with other groups, (iii) cross-sectional comparisons with non-cardiac groups, and (iv) cross-sectional comparisons with cardiac groups. The studies of comparisons within and between MINOCA and TS patients are few and do not have any specific sub-headings. The non-cardiac groups are defined as healthy controls, CVD risk controls, and patients with non-cardiac chest pain (NCCP). The cardiac groups are defined as patients with either CHD or heart failure (HF).

Results

Search results

A total of 9330 publications were found, with 2322 being eliminated during the initial screening process due to duplicates. Further exclusions based on title and abstract screening resulted in the removal of 6930

articles. Subsequently, 78 full-text articles underwent eligibility assessment, resulting in the exclusion of 50. After comprehensive full-text screening, 28 studies were found to meet the inclusion criteria. The inclusion and exclusion criteria are shown in the PRISMA flowchart (Figure 1).

Study characteristics

Table 1 provides a summary of the characteristics of the studies. There were six studies describing MINOCA patients, 20 studies describing TS patients, and 2 studies describing both groups. Out of the studies describing MINOCA patients, one study explicitly states that TS patients were excluded, three studies explicitly state that TS patients were included, and two studies do not mention whether TS patients were excluded or included.

The sample sizes of the 28 studies ranged from 13 to 5322 participants, totalling 15 572 participants. Out of these, 1241 were MINOCA patients and 898 were TS patients (total 2193). The mean age of these patients was 60 years (mean range: 46–72.7), with females (1717) constituting 80% of the samples. These studies were conducted

Table 1 Main characteristics of the included studies and population

Study characteristics	n (%)
Study design, n = studies	
Cohort	10 (36)
Cross-sectional	18 (64)
Diagnosis-related inclusion criteria, n = studies	
Takotsubo	20 (71)
MINOCA	6 (21)
Mixed (MINOCA/takotsubo/other)	2 (7)
Total sample size, n = studies	
<50	7 (25)
50–99	5 (18)
100–150	4 (14)
>150	12 (43)
MINOCA sample characteristics	
Mean/median age, n = studies	
<55 years	2
55–65 years	2
>65 years	1
Sample characteristics	
Total	668
Female	586 (88)
Takotsubo sample characteristics	
Mean/median age, n = studies	
<55 years	
55–65 years	8
>65 years	12
Sample characteristics	
Total	856
Female	754 (88)
Mixed MINOCA/takotsubo and other	
non-obstructive CVD sample characteristics	
Mean/median age	
<55 years	
55–65 years	3
>65 years	
Sample characteristics	
Total	615
Female	377 (61)

CVD, cardiovascular disease; MINOCA, myocardial infarction with non-obstructive coronary arteries

in 13 different countries: Canada (2), Denmark (1), France (1), Germany (2), Iran (1), Italy (5), New Zealand (1), Poland (1), Spain (1), Sweden (6), Switzerland (1), The Netherlands (1), and the USA (5).

Quality appraisal

Tables 2 and 3 displays the assessment of risk of bias in the included studies. Among the cross-sectional studies (range: 2–7, average: 4.6), the risk of bias was deemed low in 1 study, medium in 1 study, and high in 16 studies. Among the cohort studies (range: 5–9, average: 6.6), the risk of bias was deemed low in two studies, medium in six studies, and high in two studies.

Assessment tools

Across the 28 studies, mental health status and QoL were assessed with 36 different standardized questionnaires, and mental health disorders were diagnosed using the Mini International Neuropsychiatric Interview (for an overview, see [Supplementary material online, Table S3](#)).

Simplified summaries of cross-sectional and longitudinal comparisons of MINOCA- and TS-patients, together with the overall bias rating score from the Newcastle-Ottawa quality assessment scale are presented in [Tables 4–6](#).

Mental health and quality of life of myocardial infarction with non-obstructive coronary arteries patients

Change over time within the group

Three studies investigated change over time among MINOCA patients.^{20–22}

Two of the studies both found that self-reported QoL improved over the first year following the event.^{20,21} On the other hand, two of these studies also assessed self-reported depression and found that symptoms remained unchanged after 2 months²² and 1 year after the event.²¹ Similarly, one of these studies also found that stress symptoms remained unchanged after 2 months.²²

Change over time compared with other groups

One study followed MINOCA patients over the first 2 months after the acute event and found no differences in change of depressive and stress symptoms compared with CHD patients.²²

Cross-sectional comparisons with non-cardiac groups

Mental health status and QoL among MINOCA patients were compared with healthy controls in three studies.^{20,23,24}

All studies reported that MINOCA patients had worse mental health status compared with healthy controls. In the acute phase, patients reported lower emotional role functioning and social functioning,²⁰ and at short-term follow-up, they reported lower levels of mental QoL and higher levels of anxiety and depression.^{23,24}

Cross-sectional comparisons with cardiac groups

Mental health status and QoL among MINOCA patients were compared with cardiac groups in six studies.^{22–27}

Three studies reported that, in the acute phase, at short-term follow-up, and at long-term follow-up, MINOCA patients reported higher levels of anxiety and depression and lower mental QoL, compared with CHD patients.^{23–25} Four studies reported no differences regarding anxiety, insomnia, or stress between MINOCA and CHD patients, either in the acute phase, at short-term follow-up, or at long-term follow-up.^{22,24,26,27} One of the studies also found that MINOCA patients reported lower levels of stress in the acute phase and lower prevalence of high stress compared with CHD patients at short-term follow-up.²²

Mental health and quality of life of takotsubo patients

Change over time within the group

Three studies investigated change over time in mental health status among TS patients.^{28–30} While one study found no difference in

Table 2 Newcastle–Ottawa quality assessment scale, cross-sectional studies

Study	Total score	Representativeness of the sample		Selection		Ascertainments of exposure		Comparability		Outcome	
		sample	size	Sample size	Non-respondents	Ascertainments of exposure	Based on design and analysis	Assessment of outcome	Statistical test		
Christensen et al. (2016)	5	*			*	*				*	*
Compare et al. (2018)	5	*				*		*		*	*
Daniel et al. (2018)	5	*			*	*				*	*
Daniel et al. (2017)	5	*			*	*				*	*
Delmas et al. (2013)	5	*			*	*				*	*
Goh et al. (2016)	6	*			*	*		**		*	*
Gorini et al. (2022)	4	*				*				*	*
Izquierdo Coronel et al. (2023)	5	*			*	*				*	*
Kastaun et al. (2014)	4	*				*				*	*
Lazzeroni et al. (2022)	3					*				*	*
Naegele et al. (2016)	5	*		*		*				*	*
Olliges et al. (2020)	4	*				*				*	*
Sabisz et al. (2016)	2					*				*	*
Salmoraigo-Blotcher et al. (2016)	5	*				*		*		*	*
Sancassiani et al. (2018)	4	*				*				*	*
Smeijers et al. (2016)	4					*		*		*	*
Wallström et al. (2016)	7	*			*	*		**		*	*
Zarifeh et al. (2012)	4	*			*	*				*	*

A * signifies having met the quality assessment criteria for the Newcastle–Ottawa quality assessment scale. A maximum of one star is possible for each section, except for the section “Comparability”, where a maximum of two stars is possible.

Table 3 Newcastle–Ottawa quality assessment scale, cohort studies

Study	Total score	Selection			Demonstration that outcome was not present at the start of the study	Comparability		Outcome	
		Representativeness of the exposed cohort	Selection of non-exposed	Ascertainment of exposure		Based on design and analysis	Assessment of outcome	Length of follow-up	Adequacy of follow-up
Auger <i>et al.</i> (2019)	9	*	*	*	*	**	*	*	*
Berg <i>et al.</i> (2023)	5	*		*	*		*	*	*
Compare <i>et al.</i> (2014)	7	*	*	*		*	*	*	*
Grodzinsky <i>et al.</i> (2015)	7	*	*	*	*		*	*	*
Hausvater <i>et al.</i> (2023)	5	*	*	*	*		*	*	*
Parvand <i>et al.</i> (2022)	6	*	N/A	*	*		*	*	*
Saffari <i>et al.</i> (2017)	6	*	*	*	*		*	*	*
Safdar <i>et al.</i> (2018)	6	*	*	*	*	**	*	*	*
Sundelin <i>et al.</i> (2020)	6	*	N/A	*	*		*	*	*
Waldenbourg <i>et al.</i> (2011)	6	*	N/A	*	*		*	*	*

N/A, selection of non-exposed not applicable due to sample only consisting of exposed.

A * signifies having met the quality assessment criteria for the Newcastle–Ottawa quality assessment scale. A maximum of one star is possible for each section, except for the section “Comparability”, where a maximum of two stars is possible.

Table 4 MINOCA patients—cross-sectional comparison (simplified)

Study	n	Quality rating	Follow-up	Comparator	Symptom	Direction
Safdar et al. (2018)	2690	9	Acute phase	CHD	Depression, perceived stress	+/-
Grodzinsky et al. (2015)	5539	7	Acute phase	CHD	Depression	-
Daniel et al. (2018)	296	5	Short-term	Healthy controls	Depression, anxiety	-
				CHD	Depression	+/-
					Anxiety	-
Daniel et al. (2017)	270	5	Short-term	Healthy controls	Mental and physical QoL	-
				CHD	Mental QoL	-
					Physical QoL	+/-
				TS ^a	Mental health	+
					Mental and physical QoL	+/-
Berg et al. (2023)	459	5	Short-term	TS ^a	Emotional well-being, social functioning, fatigue	+
					Emotional role functioning, bodily pain	+/-
			Long-term	Healthy controls	Social and emotional role functioning	-
					Emotional well-being, fatigue, and bodily pain	+/-
				TS ^a	Social and emotional role functioning and bodily pain	+
					Emotional well-being and fatigue	+/-
Hausvater et al. (2023)	486	5	Acute phase	CHD	Stress, depression	+/-
			Short-term	CHD	Stress	+
					Depression	+/-
Izquierdo Coronel et al. (2023)	533	5	Acute phase	CHD	Anxiety and insomnia	+/-

Risk of bias: high (1–5), medium (6), and low (7–8). Direction refers to the outcome in MINOCA patients compared with the comparator. ‘-’ denotes worse, ‘+/-’ denotes non-different, and ‘+’ denotes better mental health or quality of life.

^aComparison of non-TS MINOCA with TS MINOCA.

symptoms of post-traumatic stress and depression between the acute phase and 3 months,²⁸ another noted reductions in self-reported stress at 6 months that remained at 1 year after the acute event.²⁹ The third study found that TS patients reported worse emotional well-being and higher prevalence of depression 1 year after the acute event, but no difference with regard to anxiety.³⁰

Change over time compared with other groups

Change in mental health status compared with other groups was assessed in two studies.^{30,31} The first study found deteriorations of emotional well-being and depression 1 year after the event that were worse for TS patients than for CHD patients, but found no differences in change of anxiety.³⁰ The second study followed TS patients over 15 years and found they had a higher risk of rehospitalization for stress and anxiety disorders, but not for mood disorders, compared with healthy controls.³¹ On the other hand, the same study found no difference in risk for rehospitalization over 15 years when comparing TS patients with CHD patients.

Cross-sectional comparisons with non-cardiac groups

Mental health status and QoL among TS patients was compared with non-cardiac groups in nine studies.^{32–40} Two studies found that TS patients had higher levels of psychological distress at short-term follow-up³⁵ and lower QoL at long-term follow-up³⁶ compared with healthy controls. On the other hand, two studies found no differences in

QoL, neither compared with healthy controls at short-term follow-up,³² nor compared with CVD risk controls at long-term follow-up.³³ Similarly, two studies found no differences at long-term follow-up, neither regarding psychological distress compared with CVD risk controls,³³ nor psychiatric symptoms compared with healthy controls.³⁴

Higher levels of anxiety for TS patients compared with healthy controls were observed in three studies, both in the acute phase³⁶ and at long-term follow-up.^{37,38} Two studies observed no differences regarding anxiety at long-term follow-up, neither in anxiety levels,³⁹ nor in lifetime prevalence of anxiety disorders³² as compared with healthy controls. Similarly, one study found no differences in anxiety at short-term follow-up, compared with NCCP patients.⁴⁰

Three studies found that levels of depression were higher in TS patients compared with healthy controls, both in the acute phase³⁶ and at long-term follow-up.^{34,38} Similarly, one study found that TS patients had a higher lifetime prevalence of major depressive disorders than healthy controls, assessed at short-term follow-up, but no difference in prevalence of bipolar disorder.³² On the other hand, one study found no difference in levels of depression for TS patients, when compared with NCCP patients.⁴⁰

One study found higher levels of stress in TS patients compared with healthy controls at short-term follow-up.³⁵ However, no difference was observed in two studies regarding stress at long-term follow-up.^{34,38}

While one study found TS patients to report higher levels of post-traumatic stress than healthy controls at short-term follow-up,³⁵ two other studies reported no differences at short-term follow-up, neither

Table 5 Takotsubo patients—cross-sectional comparison (simplified)

Study	n	Quality rating	Follow-up	Comparator	Symptom	Direction
Auger <i>et al.</i> (2019)	3650	9	Long-term	CHD	Stress, anxiety	+/-
				Healthy controls	Stress, anxiety	-
Compare <i>et al.</i> (2014)	74	7	Acute phase	CHD	Anxiety, depression, QoL	+/-
			Long-term	CHD	Psychological distress	-
Wallström <i>et al.</i> (2016)	127	7	Short-term	CHD	Anxiety, depression, stress, sleep disturbance	+/-
Goh <i>et al.</i> (2016)	184	6	Long-term	CHD	Anxiety	-
					Depression	+/-
Saffari <i>et al.</i> (2017)	188	6	Acute phase	CHD	Sexual functioning, anxiety, depression	-
			Long-term	CHD	Sexual functioning	-
Christensen <i>et al.</i> (2016)	173	5	Long-term	CHD	Well-being, anxiety, depression	-
				Healthy controls	Well-being, depression	+/-
					Anxiety	-
Delmas <i>et al.</i> (2013)	95	5	Acute phase	CHD	Psychiatric diagnoses	-
Neagele <i>et al.</i> (2016)	43	5	Long-term	Matched controls	QoL	+/-
Salmoirago-Blotcher (2016)	107	5	Acute phase	CHD	Psychological distress, stress, post-traumatic stress	-
				Healthy controls		
Kastaun <i>et al.</i> (2014)	59	4	Long-term	CHD	Psychiatric symptoms, depression, stress	+/-
				Healthy controls	Psychiatric symptoms, stress	
					Depression, chronic worry	-
Olliges <i>et al.</i> (2020)	136	4	Long-term	Healthy controls	Anxiety, depression, QoL	-
				CHD	Anxiety, depression, QoL	+/-
Sancassiani <i>et al.</i> (2018)	95	4	Short-term	Healthy controls	QoL, anxiety disorders, post-traumatic stress disorder	+/-
					Major depressive disorder	-
Smeijers <i>et al.</i> (2016)	56	4	Long-term	Heart failure	Anxiety, depression, stress	+/-
				Healthy controls	Anxiety, depression	-
					Stress	+/-
Zarifeh <i>et al.</i> (2012)	17	4	Short-term	NCCP	Anxiety, Depression, post-traumatic stress	+/-
				CHD		
Gorini <i>et al.</i> (2022)	40	4	Short-term	CHD	Anxiety, post-traumatic stress	+/-
					Reassurance seeking	-
					Fear of death, interference	+
Lazzeroni <i>et al.</i> (2022)	19	3	Long-term	Healthy controls	Depression, anxiety	+/-
Sabisz <i>et al.</i> (2016)	26	2	Long-term	Norm values	Anxiety	-
					Depression	+/-

Risk of bias: high (1–5), medium (6), and low (7–8). Direction refers to the outcome in MINOCA patients compared with the comparator. ‘-’ denotes worse, ‘+/-’ denotes non-different, and ‘+’ denotes better mental health or quality of life.

CHD, coronary heart disease; QoL, quality of life; NCCP, non-cardiac chest pain.

with regard to self-reported levels nor in lifetime prevalence of post-traumatic stress disorders.^{32,40} It was also reported that TS patients had higher sexual distress and more sexual dysfunction compared with healthy controls in the acute phase,³⁶ and they reported more symptoms of chronic worry at long-term follow-up.³⁴

Cross-sectional comparisons with cardiac groups

Mental health status and QoL among TS patients was compared with HF or CHD patients in 11 studies.^{30,34–36,38,41–46} Compared with CHD patients, in the acute phase, one study found higher psychological distress for TS patients³⁵ and two studies found lower emotional well-

being at long-term follow-up.^{30,42} Two other studies reported no differences in QoL compared with CHD patients, in the acute phase,³⁶ or in psychiatric symptoms at long-term follow-up.³⁴

Regarding self-reported anxiety, four studies found TS patients to have higher levels compared with CHD patients, both in the acute phase³⁶ and at long-term follow-up.^{42,44,45} On the other hand, three studies found no differences in anxiety levels compared with either HF or CHD patients, neither in the acute phase,³⁰ at short-term follow-up,⁴³ nor at long-term follow-up.^{30,38} Additionally, one study found TS patients to report more fear of death and reassurance seeking related to health anxiety but no difference regarding state anxiety compared with CHD patients at short-term follow-up.⁴⁶

Table 6 MINOCA and takotsubo patients—longitudinal comparisons (simplified)

Study	n	Quality rating	Follow-up	Comparator	Symptom	Direction
Safdar et al. (2018)	2690	9	3 years	CHD	Depression and perceived stress	+/-
Compare et al. (2014)	74	7	12 months	Within group	Depression and emotional well-being	-
				CHD	Anxiety	+/-
Sundelin et al. (2020)	20	6	6 months	Within group	Depression and emotional well-being	-
			12 months		Anxiety	+/-
Waldenbourg et al. (2011)	13	6	3 months	Within group	Stress	+
Parvand et al. (2022)	154	6	12 months	Within group	Post-traumatic stress, depression	+/-
Hausvater et al. (2023)	486	5	2 months	CHD	QoL, mental health	+
Berg et al. (2023)	459	5	6 months	Within group	Stress, depression	+/-
			12 months	Within group	Bodily pain, fatigue, social and emotional role functioning, emotional well-being	+
					Fatigue, emotional well-being	+
					Bodily pain, social and emotional role functioning	+/-

Risk of bias: high (1–5), medium (6–7), and low (8–9). Comparator indicates either only change over time within the group or the difference in change over time compared with another group. Direction refers to the outcome in MINOCA patients compared with the comparator. ‘-’ denotes worse, ‘+/-’ denotes non-different, and ‘+’ denotes better mental health or quality of life.

CHD, coronary heart disease; QoL, quality of life; TS, takotsubo syndrome.

Regarding self-reported depression, three studies found higher levels for TS patients compared with CHD patients, both during the acute phase³⁶ and at long-term follow-up.^{42,45} Similarly, one study found the prevalence of major depressive disorder to be higher in the acute phase in TS patients than in CHD patients.⁴¹ Conversely, there were five studies that found no differences in depression between TS patients and CHD patients, neither in the acute phase,³⁰ at short-term follow-up^{43,46} nor at long-term follow-up.^{38,44}

One study found TS patients to have more symptoms of stress than CHD patients in the acute phase.³⁵ However, in follow-up assessments, none of the three studies identified any differences in stress symptoms, neither compared with CHD patients at short-term follow-up,^{43,46} nor HF patients at long-term follow-up.³⁸ Similarly, one of these studies did not find any differences in symptoms of mental fatigue nor sleep disturbance at short-term follow-up⁴³ compared with CHD patients.

One study found that in the acute phase, TS patients had higher symptoms of post-traumatic stress compared with CHD patients.³⁵ However, at short-term follow-up, another study found no differences in post-traumatic stress compared with CHD patients.⁴⁶ One study found that TS patients reported more sexual distress and experienced more symptoms of sexual dysfunction compared with CHD patients, both in the acute phase, and at long-term follow-up.³⁶

Comparisons within and between myocardial infarction with non-obstructive coronary arteries and takotsubo syndrome patients

Two studies compared MINOCA patients with TS patients and found that patients with TS had lower emotional and social well-being and vitality in the acute phase, worse mental health status at short-term follow-up, and more impairment of emotional and social role functioning at long-term follow-up compared with MINOCA patients.^{20,23}

Another study found that TS patients who reported an emotional trigger to the event had higher levels of depression compared with TS patients that did not report an emotional trigger at long-term follow-up.⁴⁷ The same study also reported that CHD patients who had reported an emotional trigger had higher levels of depression than TS patients that did not report an emotional trigger. No differences were found between CHD and TS patients that both reported emotional triggers.⁴⁷

Discussion

Summary of findings

In this paper, studies of mental health status and QoL in MINOCA and TS patients after cardiac event have been systematically reviewed, compared with healthy controls, CVD risk controls, and cardiac patients, both cross-sectionally and in change over time. Generally, MINOCA and TS patients report worse mental health status and QoL than non-cardiac groups and at least as poor mental health status and QoL compared with cardiac patients. Results of longitudinal change were inconsistent, showing improved, worsened, or unchanged mental health status and QoL over time for MINOCA and TS patients.

Quality assessment and study-specific discussion

The quality assessment of included studies varied, with many studies demonstrating a high risk of bias (18/28). Only three studies were deemed at low risk of bias and should be primarily considered when evaluating the evidence regarding mental health status and QoL of MINOCA and TS patients, after the acute event. Among these studies, results were not completely congruent.

The first study compared MINOCA patients with CHD patients and investigated the prevalence of angina compared with patients with coronary artery disease.²⁵ Additionally, they collected data on QoL using

the SF-12 during the follow-ups and on depression using the PHQ-9 at baseline. They found that during the acute phase, MINOCA patients reported both higher levels of depression and worse mental QoL compared with the CHD patients. Additionally, at the 12-month follow-up, MINOCA patients still reported worse mental QoL.

The second study also compared MINOCA patients with CHD patients, with the purpose of characterizing the socio-demographic, psychosocial, and clinical presentations of these patients, as well as predicting clinical outcomes and QoL.²⁷ They used the SAQ to collect angina-related QoL, the PHQ-9 for depressive symptoms and the PSS for symptoms of stress. Compared with the first study, they found no differences in symptoms of stress or depression during the acute phase. Additionally, while unadjusted analyses showed that CHD patients reported worse angina-related QoL at the 12-month follow-up, this association did not remain in adjusted analyses. On the other hand, at the 1- and 12-month follow-ups, ratings of stress and depression were numerically higher for the MINOCA patients, but no statistical tests of difference of these assessments were conducted.

The third study compared TS patients both with CHD patients and healthy controls, aiming to investigate differences in mortality, morbidity, and future hospitalizations.³¹ Mental disorders [grouped as (i) schizophrenia, (ii) stress and anxiety, and (iii) mood disorder] were collected through hospital registries. They found that TS patients had a higher risk for stress and anxiety disorders, but not schizophrenia or mood disorders compared with healthy controls. And although the hazard ratios were higher for TS patients for rehospitalization for any mental disorder, they were not statistically significant compared with CHD patients.

These three studies reflect the general findings, with results indicating that MINOCA and TS patients have worse mental health status and QoL than healthy controls and either worse or non-different compared with CHD patients.

Between group comparisons in mental health and quality of life

In the comparisons of MINOCA and TS patients with CHD patients, all but one of the included studies showed that there was either no difference in mental health status or QoL or that it was worse for MINOCA and TS patients. Only one study²² found that CHD patients reported higher stress and depression than their MINOCA counterparts. However, this study had a high risk of bias, and differences reported were minimal and most likely not clinically relevant. While it is possible that in certain cases, MINOCA patients experience better mental health status and QoL than CHD patients, there is little evidence currently supporting this. Current conclusions based on empirical findings are heavily inclined towards MINOCA and TS patients having at least as bad mental health status and QoL compared with CHD patients after the acute event.

Only two of the included studies compared MINOCA patients with TS patients.^{20,23} These studies found that TS patients either had comparable mental health status and QoL or worse than MINOCA patients. It is known that around one-third of TS cases are reported to be triggered by emotional stress.⁵ It is possible that the difference in mental health and QoL between MINOCA and TS may be driven by that TS patients have experienced an emotional trigger to a greater extent. Indeed, one of the studies compared TS patients with an emotional trigger vs. those without and found that those who experienced an emotional trigger suffered from worse mental health status in the aftermath of the event.⁴⁷ Yet, we must emphasize that studies comparing these patients are few and more research is needed.

Change over time in mental health and quality of life

When scrutinizing studies investigating longitudinal change of mental health status and QoL, it was hard to discern any consistent trends. Some symptoms were reported to improve over time,^{20,21,29} while other symptoms were not.^{20,27,28,30} One study even found depression and emotional well-being to become worse over time.³⁰ This suggests that, in many cases, mental health status and QoL in MINOCA and TS patients do not improve spontaneously. For CHD patients, it is commonly reported that in about 50% of cases, mental health status and QoL do not naturally improve after the acute event.^{48–51} Among these patients, predictors for maintained poor mental health status are history of mental distress, worse mental health status at baseline, poor socioeconomic status, and younger age.^{50–52} Perhaps, there are similarities between the CHD patients with persistently worse mental health status and QoL and MINOCA and TS patients. We know, for example, that MINOCA and TS patients are on average younger and more often have previously identified mental health disorders, compared with CHD patients.^{5,6,10} While more research is needed to clarify risk factors and prognosis for mental health status and QoL in MINOCA and TS patients, the results presented in this review indicate that there are persistent mental health issues among these patients that need to be addressed. A previous study found that these patients valued psychological support and that the conduct of a psychological intervention RCT was feasible.⁵³ We believe that based on these findings, interventional studies for improving mental health status and QoL, after the acute event, could prove highly beneficial.

Mechanisms of association between diagnosis and mental health and quality of life

Given the results presented in this review, the association between mental health status and QoL and the diagnosis of MINOCA or TS seem undeniable, but it is at this point difficult to draw any firm conclusions regarding a potential causal relationship. The review is based only on observational data, and as none of the included studies used causal diagrams to guide their analyses, and many only presented unadjusted analyses, still much uncertainty exists regarding this question.

However, if the associations reflect causal relationships, it is still unclear to what role mental health status plays. Even though the results presented in this study only concern assessments of mental health status and QoL after the acute event, we cannot exclude their presence preceding the event. On the one hand, a common hypothesis for TS is that it is triggered by emotional stress,⁵ and it is possible that poor mental health increases the risk for MINOCA or TS through behavioural or physiological factors.⁵⁴ On the other hand, in addition to that a cardiac event may be life-threatening and potentially traumatic,⁵⁵ the uncertainty regarding cause, treatment, and prognosis of MINOCA and TS is likely a further burden to the patients. Indeed, patients have described themselves as having an increased stress sensitivity after the event.²⁹ Perhaps, what is most likely is a combination of the two, acting bi-directionally. While more research is needed to clarify potential confounders, and the nature and potential direction of this association, one thing remains clear: after the acute event, many MINOCA and TS patients suffer from worse mental health status and reduced QoL.

Methodological discussion

Including many studies with small sample sizes and data that has a high risk of bias may seem superfluous. However, as this is, to our knowledge, the first review of mental health status and QoL among

MINOCA and TS patients after the acute event, there is a value in reporting all conducted studies to provide an overview of the field. It is not only important to discuss the outcomes assessed in the included studies but also the quality of available data. Initially, our wish was to conduct a meta-analysis, pooling the results of included studies. However, at the present, it is difficult to conduct a meta-analysis for this research question, due to the high heterogeneity in study designs and high risk of bias for these outcomes.

We would again like to remind the reader that the bias assessment was made based on the data and analyses corresponding to our research question. This means that while the data of mental health status and QoL of MINOCA and TS patients have a high risk of bias, the study itself might not have been poorly conducted in regards to its primary aim. The most common issues with bias were lacking descriptions of sample size calculation, descriptions of non-respondents, and adjustment for confounding in the analyses (see [Tables 2](#) and [3](#)). Lacking sample size calculations may lead to a higher risk of Type 2 errors. As the number of non-responders is unknown, the observed estimates might be biased either towards or away from the null, depending on the characteristics of these subjects. Lacking adjustment for confounding, either by study design or statistical models, increases the risk of observing spurious associations between variables. This lack of information hinders making solid claims regarding the mental health status and QoL among MINOCA and TS patients after the acute event, and more high-quality studies are needed, specifically addressing these questions.

A growing evidence base of high-quality studies will also facilitate investigation of specific symptom clusters. Currently, even when assessing the same construct, there are variations regarding the assessment procedure, such as which tools were used, if data were self-reported or collected by a clinician, length of follow-up, if data were reported at symptom levels or as clinical diagnoses, and if it was on a graded scale or dichotomous. For example, two of the studies with low risk of bias demonstrated seemingly contradictory results regarding depression in MINOCA patients.^{25,27} This could be due to procedures of data handling. Helpfully, both studies used the PHQ-9 to assess depression, but one dichotomized the variable (cut-off ≥ 10),²⁷ and the other reported the variable on a continuous scale.²⁵ It is known that dichotomization increases the risk of Type 2 errors,⁵⁶ and it is possible that the *P*-value (which was $P = 0.06$) would have been deemed statistically significant if PHQ-9 was analysed continuously.

Finally, a limitation is the shifting definitions of diagnoses and updates of diagnostic procedures of MINOCA and TS. Depending on the year of publication, the definition and diagnostic procedures of MINOCA may vary. For example, in 2018, both myocarditis and TS were excluded from the definition of an MI,⁸ perhaps another explanation for the contradictory results of Safdar *et al.*²⁷ and Grodzinsky *et al.*²⁵ as the latter did not mention any exclusion of TS patients. Furthermore, recent studies have shown that for example using cardiovascular magnetic resonance imaging greatly improves correct diagnosis of MINOCA patients.⁵⁷ As such, although we describe and narratively compare studies of MINOCA and TS patients, uncertainties remain whether the different studies actually refer to the same patients.

Taken together, these issues remain an obstacle in making firm conclusions regarding specific symptoms and trends of mental health and QoL, in MINOCA and TS patients, and more high-quality research is needed.

Conclusion

In this systematic review, we have reviewed mental health status and QoL in MINOCA and TS patients after the acute event. The current literature suggests that these patients more often have worse mental health status and QoL than non-cardiac groups. Compared with CHD patients, it is yet

difficult to conclude whether mental distress and QoL are equal or worse for MINOCA and TS patients. Studies investigating longitudinal change were few and showed no coherent trends, but gave concerns regarding persistent poor mental health status and reduced QoL. Many of the studies had a high risk of bias and suffered from small sample sizes calling for a need for more high-quality research, investigating this specific question.

Supplementary material

Supplementary material is available at *European Journal of Preventive Cardiology*.

Acknowledgements

The authors would like to send their most sincere thanks to *The Bergman Estate on Fårö Foundation* for their grant of residency. This serene and secluded location gave inspiration and peace of mind for the work with this review. We would also like to thank the U-CARE intern Stephanie who helped with title and abstract screening.

Author contribution

C.H., E.M.G.O., E.R., P.Le., P.T., and S.H. contributed to the conception or design of the work. S.H. contributed to the acquisition of data for the work. P.Le. and S.H. contributed to the analysis and interpretation of the data for this work. P.Le. and S.H. drafted the manuscript. A.N., A.U., C.H., E.M.G.O., E.R., J.S., L.K., P.Le., P.Ly., P.T., and S.H. critically revised the manuscript. All gave final approval and agreed to be accountable for all aspects of work ensuring integrity and accuracy.

Funding

This article was funded by the Swedish Research Council (grant 2018-02655), the Swedish Heart Lung Foundation (grant 20200221), and the Swedish Heart and Lung Association (grant 2019:41). It was further sponsored by the Swedish Research Council also as part of the strategic research environment U-CARE (2009-1093).

Conflict of interest: none declared.

Data availability

All relevant data are within the manuscript and its supporting information files.

References

1. World Health Organization. *Global Health Estimates for 2020: deaths by Cause, age, sex, by Country and by Region, 2000–2019*. Geneva, Switzerland: WHO; 2020. p. 1–59.
2. Cohen BE, Edmondson D, Kronish IM. State of the art review: depression, stress, anxiety, and cardiovascular disease. *Am J Hypertens* 2015;**28**:1295–1302.
3. Gross H, Sternberg WH. Myocardial infarction without significant lesions of coronary arteries. *Arch Intern Med* 1939;**64**:249–267.
4. Miller RD, Burchell HB, Edwards JE. Myocardial infarction with and without acute coronary occlusion: a pathologic study. *AMA Arch Intern Med* 1951;**88**:597–604.
5. Singh T, Khan H, Gamble DT, Scally C, Newby DE, Dawson D. Takotsubo syndrome: pathophysiology, emerging concepts, and clinical implications. *Circulation* 2022;**145**:1002–1019.
6. Tamis-Holland JE, Jneid H, Reynolds HR, Agewall S, Brilakis ES, Brown TM, et al. Contemporary diagnosis and management of patients with myocardial infarction in the absence of obstructive coronary artery disease: a scientific statement from the American Heart Association. *Circulation* 2019;**139**:E891–E908.
7. Agewall S, Beltrame JF, Reynolds HR, Niessner A, Rosano G, Caforio ALP, et al. ESC working group position paper on myocardial infarction with non-obstructive coronary arteries. *Eur Heart J* 2017;**38**:143–153.
8. Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA, et al. Fourth universal definition of myocardial infarction (2018). *J Am Coll Cardiol* 2018;**72**:2231–2264.

9. Lopez-Pais J, Izquierdo Coronel B, Raposeiras-Roubin S, Álvarez Rodríguez L, Vedia O, Almendro-Delia M, et al. Differences between takotsubo and the working diagnosis of myocardial infarction with nonobstructive coronary arteries on behalf of the RETAKO investigators. *Front Cardiovasc Med* 2022;**9**:742010.
10. Abdu FA, Mohammed A-Q, Liu L, Xu Y, Che W. Myocardial infarction with nonobstructive coronary arteries (MINOCA): a review of the current position. *Rev Artic Cardiol* 2020;**145**:543–552.
11. La S, Beltrame J, Tavella R. Sex-specific and ethnicity-specific differences in MINOCA. *Nat Rev Cardiol* 2024;**21**:192–202.
12. Herling de Oliveira LL, Correia VM, Nicz PFG, Soares PR, Scudeler TL. MINOCA: one size fits all? Probably not—A review of etiology, investigation, and treatment. *J Clin Med* 2022;**11**:5497.
13. Cheema AN, Yanagawa B, Verma S, Bagai A, Liu S. Myocardial infarction with nonobstructive coronary artery disease (MINOCA): a review of pathophysiology and management. *Curr Opin Cardiol* 2021;**36**:589–596.
14. Galli F, Bursi F, Carugo S. Traumatic events, personality and psychopathology in takotsubo syndrome: a systematic review. *Front Psychol* 2019;**10**:2742.
15. Nayeri A, Rafla-Yuan E, Krishnan S, Ziaiean B, Cadeiras M, McPherson JA, et al. Psychiatric illness in takotsubo (stress) cardiomyopathy: a review. *Psychosomatics* 2018;**59**:220–226.
16. Carroll AJ, Goergen J, Wafford QE, Flaherty JD, Grady KL, Feingold KL. Psychiatric conditions in patients presenting with takotsubo syndrome: a systematic review and synthesis of case studies. *Gen Hosp Psychiatry* 2020;**65**:54–63.
17. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg* 2010;**8**:336–341.
18. Luchini C, Stubbs B, Solmi M, Veronese N. Assessing the quality of studies in meta-analyses: advantages and limitations of the Newcastle Ottawa scale. *World J Meta Anal* 2017;**5**:80–84.
19. Ribeiro CM, Beserra BTS, Silva NG, Lima CL, Rocha PRS, Coelho MS, et al. Exposure to endocrine-disrupting chemicals and anthropometric measures of obesity: a systematic review and meta-analysis original research. *BMJ Open* 2020;**10**:e033509.
20. Berg E, Agewall S, Brolin EB, Caidahl K, Cederlund K, Collste O, et al. Health-related quality-of-life up to one year after myocardial infarction with non-obstructive coronary arteries. *Eur Hear J Qual Care Clin Outcomes* 2023;**9**:639–644.
21. Parvand M, Cai L, Ghadiri S, Humphries KH, Starovoytov A, Daniele P, et al. One-year prospective follow-up of women with INOCA and MINOCA at a Canadian women's heart centre. *Can J Cardiol* 2022;**38**:1600–1610.
22. Hausvater A, Spruill TM, Xia Y, Smilowitz NR, Arabadjian M, Shah B, et al. Psychosocial factors of women presenting with myocardial infarction with or without obstructive coronary arteries. *J Am Coll Cardiol* 2023;**82**:1649–1658.
23. Daniel M, Agewall S, Caidahl K, Collste O, Ekenbäck C, Frick M, et al. Effect of myocardial infarction with nonobstructive coronary arteries on physical capacity and quality-of-life. *Am J Cardiol* 2017;**120**:341–346.
24. Daniel M, Agewall S, Berglund F, Caidahl K, Collste O, Ekenbäck C, et al. Prevalence of anxiety and depression symptoms in patients with myocardial infarction with non-obstructive coronary arteries. *Am J Med* 2018;**131**:1118–1124.
25. Grodzinsky A, Arnold SV, Gosch K, Spertus JA, Foody JM, Beltrame J, et al. Angina frequency after acute myocardial infarction in patients without obstructive coronary artery disease. *Eur Heart J Qual Care Clin Outcomes* 2015;**1**:92–99.
26. Izquierdo Coronel B, López Pais J, Nieto Ibáñez D, Olsen Rodríguez R, Galán Gil D, Perela Álvarez C, et al. Prevalence and prognosis of anxiety, insomnia, and type D personality in patients with myocardial infarction: a Spanish cohort. *Cardiol J* 2024;**31**:261–270.
27. Safdar B, Spatz ES, Dreyer RP, Beltrame JF, Lichtman JH, Spertus JA, et al. Presentation, clinical profile, and prognosis of young patients with myocardial infarction with nonobstructive coronary arteries (MINOCA): results from the VIRGO study. *J Am Heart Assoc* 2018;**7**:e009174.
28. Waldenborg M, Soholat M, Kähäri A, Emilsson K, Fröbert O. Multidisciplinary assessment of tako tsubo cardiomyopathy: a prospective case study. *BMC Cardiovasc Disord* 2011;**11**:14.
29. Sundelin R, Bergsten C, Tornvall P, Lyngå P. Self-rated stress and experience in patients with takotsubo syndrome: a mixed methods study. *Eur J Cardiovasc Nurs* 2020;**19**:740–747.
30. Compare A, Grossi E, Bigli R, Proietti R, Shonin E, Orrego PS, et al. Stress-induced cardiomyopathy and psychological wellbeing 1 year after an acute event. *J Clin Psychol Med Settings* 2014;**21**:81–91.
31. Auger N, Paradis G, Healy-Profitts J, He S, Potter BJ. Outcomes of takotsubo syndrome at 15 years: a matched cohort study. *Am J Med* 2020;**133**:627–634.e4.
32. Sancassiani F, Carta MG, Montisci R, Preti A, Machado S, Moro MF, et al. Takotsubo syndrome is associated with mood disorders and antidepressants use, not with anxiety and impairment of quality of life due to the psychiatric disorder. *Clin Pract Epidemiol Ment Heal* 2018;**14**:26–32.
33. Naegele M, Flammer AJ, Enseleit F, Roas S, Frank M, Hirt A, et al. Endothelial function and sympathetic nervous system activity in patients with takotsubo syndrome. *Int J Cardiol* 2016;**224**:226–230.
34. Kastaun S, Schwarz NP, Juenemann M, Yeniguen M, Nef HM, Moellmann H, et al. Cortisol awakening and stress response, personality and psychiatric profiles in patients with takotsubo cardiomyopathy. *Heart* 2014;**100**:1786–1792.
35. Salmoiraghi-Blotcher E, Rosman L, Wittstein IS, Dunsiger S, Swales HH, Aurigemma GP, et al. Psychiatric history, post-discharge distress, and personality characteristics among incident female cases of takotsubo cardiomyopathy: a case-control study. *Heart Lung* 2016;**45**:503–509.
36. Saffari M, Lin C-Y, Broström A, Mårtensson J, Malm D, Burri A, et al. Investigating sexual problems, psychological distress and quality of life in female patients with takotsubo cardiomyopathy: a prospective case-control study. *Eur J Cardiovasc Nurs* 2017;**16**:614–622.
37. Sabisz A, Treder N, Fijałkowska M, Sieminski M, Fijałkowska J, Naumczyk P, et al. Brain resting state functional magnetic resonance imaging in patients with takotsubo cardiomyopathy an inseparable pair of brain and heart. *Int J Cardiol* 2016;**224**:376–381.
38. Smeijers L, Szabó BM, Kop VJ. Psychological distress and personality factors in takotsubo cardiomyopathy. *Netherlands Hear J* 2016;**24**:530–537.
39. Lazzeroni D, Ciraci C, Sommaruga M, Oggioni C, Saccò M, Ziveri V, et al. Perceived anxiety, coping, and autonomic function in takotsubo syndrome long after the acute event. *Life* 2022;**12**:1376.
40. Zarifeh JA, Mulder RT, Kerr AJ, Chan CW, Bridgman PG. Psychology of earthquake-induced stress cardiomyopathy, myocardial infarction and non-cardiac chest pain. *Intern Med J* 2012;**42**:369–373.
41. Delmas C, Lairez O, Mulin E, Delmas T, Boudou N, Dumonteil N, et al. Anxiodepressive disorders and chronic psychological stress are associated with tako-tsubo cardiomyopathy: new physiopathological hypothesis. *Circ J* 2013;**77**:175–180.
42. Christensen TE, Bang LE, Holmvang L, Hasbak P, Kjær A, Bech P, et al. Neuroticism, depression and anxiety in takotsubo cardiomyopathy. *BMC Cardiovasc Disord* 2016;**16**:118.
43. Wallström S, Ulin K, Omerovic E, Ekman I. Self-reported symptoms 8 weeks after discharge: a comparison of takotsubo syndrome and myocardial infarction. *Int J Cardiol* 2016;**224**:348–352.
44. Goh ACH, Wong S, Zaroff JG, Shafee N, Lundstrom RJ. Comparing anxiety and depression in patients with takotsubo stress cardiomyopathy to those with acute coronary syndrome. *J Cardiopulm Rehabil Prev* 2016;**36**:106–111.
45. Olliges E, Burgdorf C, Ladwig KH, Moeller C, Deftu-Kloes D, Pohl S, et al. Psychosocial and physical long-term outcome in patients with a history of takotsubo cardiomyopathy or myocardial infarction—a multi-centered case control study. *Psychol Heal Med* 2020;**25**:989–1003.
46. Gorini A, Galli F, Giuliani M, Pierobon A, Werba JP, Adriano EP, et al. Psychological characteristics of patients with takotsubo syndrome and patients with acute coronary syndrome: an explorative study toward a better personalized care. *J Pers Med* 2022;**12**:38.
47. Compare A, Brugnera A, Spada MM, Zarbo C, Tasca GA, Sassaroli S, et al. The role of emotional competence in takotsubo cardiomyopathy. *Psychosom Med* 2018;**80**:377–384.
48. Grace SL, Abbey SE, Irvine J, Shnek ZM, Stewart DE. Prospective examination of anxiety disorder and its relationship to cardiac symptoms and recurrent cardiac events. *Psychother Psychosom* 2004;**73**:344–352.
49. Schleifer SJ, Macari-Hinson MM, Coyle DA, Slater WR, Kahn M, Gorlin R, et al. The nature and course of depression following myocardial infarction. *Arch Intern Med* 1989;**149**:1785–1789.
50. Hanssen TA, Nordrehaug JE, Eide GE, Bjelland I, Rokne B. Anxiety and depression after acute myocardial infarction: an 18-month follow-up study with repeated measures and comparison with a reference population. *Eur J Cardiovasc Prev Rehabil* 2009;**16**:651–659.
51. Murphy B, Le Grande M, Alvarenga M, Worcester M, Jackson A. Anxiety and depression after a cardiac event: prevalence and predictors. *Front Psychol* 2020;**10**:3010.
52. Bjerkeset O, Nordahl HM, Mykletun A, Holmen J, Dahl AA. Anxiety and depression following myocardial infarction: gender differences in a 5-year prospective study. *J Psychosom Res* 2005;**58**:153–161.
53. Rondung E, Humphries SM, Olsson EM, Sundelin R, Norlund F Held C, et al. Reducing stress and anxiety in patients with myocardial infarction with non-obstructive coronary arteries or takotsubo syndrome: a non-randomized feasibility study. *Internet Interv* 2022;**29**:100562.
54. Levine GN, Cohen BE, Commodore-Mensah Y, Fleury J, Huffman JC, Khalid U, et al. Psychological health, well-being, and the mind-heart-body connection: a scientific statement from the American Heart Association. *Circulation* 2021;**143**:E763–E783.
55. Jacquet-Smailovic M, Tarquinio C, Alla F, Denis I, Kirche A, Tarquinio C, et al. Posttraumatic stress disorder following myocardial infarction: a systematic review. *J Trauma Stress* 2021;**34**:190–199.
56. MacCallum RC, Zhang S, Preacher KJ, Rucker DD. On the practice of dichotomization of quantitative variables. *Psychol Methods* 2002;**7**:19–40.
57. Sörensson P, Ekenbäck C, Lundin M, Agewall S, Bacvosics Brolin E, Caidahl K, et al. Early comprehensive cardiovascular magnetic resonance imaging in patients with myocardial infarction with nonobstructive coronary arteries. *JACC Cardiovasc Imaging* 2021;**14**:1774–1783.